

Math 415 - Lecture 27

An application of QR -decomposition, Change of basis

Friday October 30th 2015

Textbook reading: Chapter 3.4, Chapter 2.6

Textbook reading: Chapter 3.4, Chapter 2.6

Suggested practice exercises: Chapter 2.6: Exercises 36, 37, 38,39,
40,43

Textbook reading: Chapter 3.4, Chapter 2.6

Suggested practice exercises: Chapter 2.6: Exercises 36, 37, 38,39,
40,43

Khan Academy video: Change of basis

Textbook reading: Chapter 3.4, Chapter 2.6

Suggested practice exercises: Chapter 2.6: Exercises 36, 37, 38,39,
40,43

Khan Academy video: Change of basis

Strang lecture: Change of basis; image compression

Review

Theorem (QR decomposition)

Let A be a $m \times n$ matrix of rank n with linear independent columns. There is an orthogonal matrix $m \times n$ -matrix Q and an upper triangular $n \times n$ invertible matrix R such that

$$A = QR.$$

Theorem (QR decomposition)

Let A be a $m \times n$ matrix of rank n with linear independent columns. There is an orthogonal matrix $m \times n$ -matrix Q and an upper triangular $n \times n$ invertible matrix R such that

$$A = QR.$$

Theorem

Let A be a matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Theorem (QR decomposition)

Let A be a $m \times n$ matrix of rank n with linear independent columns. There is an orthogonal matrix $m \times n$ -matrix Q and an upper triangular $n \times n$ invertible matrix R such that

$$A = QR.$$

Theorem

Let A be a matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Theorem (QR decomposition)

Let A be a $m \times n$ matrix of rank n with linear independent columns. There is an orthogonal matrix $m \times n$ -matrix Q and an upper triangular $n \times n$ invertible matrix R such that

$$A = QR.$$

Theorem

Let A be a matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T \mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Theorem (QR decomposition)

Let A be a $m \times n$ matrix of rank n with linear independent columns. There is an orthogonal matrix $m \times n$ -matrix Q and an upper triangular $n \times n$ invertible matrix R such that

$$A = QR.$$

Theorem

Let A be a matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.


Then the solution of $R\mathbf{x} = Q^T \mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.


An application in wireless communication

In multiple-input and multiple-output (short: MIMO) systems, a transmitter sends multiple streams by multiple transmit antennas. Let us suppose there are n transmitters and m receivers. This can be modelled using Linear Algebra:


$$\underbrace{\begin{bmatrix} y_1 \\ \vdots \\ y_m \end{bmatrix}}_{\text{received vector } \mathbf{y}} = \underbrace{\begin{bmatrix} h_{1,1} & \dots & h_{1,n} \\ \vdots & \ddots & \vdots \\ h_{m,1} & \dots & h_{m,n} \end{bmatrix}}_{\text{channel matrix } H} \underbrace{\begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}}_{\text{transmitted vector } \mathbf{x}}.$$

Suppose that the channel matrix H is known both to person A who is sending information and to person B who is receiving the information.






Let us try and understand the engineering meaning of some of the linear algebra of the matrix H and the equation $y = Hx$.



Let us try and understand the engineering meaning of some of the linear algebra of the matrix H and the equation $y = Hx$.

Remember: the x vector describes what the transmitter is sending out and y is the vector describing what is received.



Let us try and understand the engineering meaning of some of the linear algebra of the matrix H and the equation $y = Hx$.

Remember: the x vector describes what the transmitter is sending out and y is the vector describing what is received.

We want to understand

$$y = Hx.$$

We want to understand

$$y = Hx.$$

- What is the first column of H ?

We want to understand

$$y = Hx.$$

- What is the first column of H ?

We want to understand

$$y = Hx.$$

- What is the first column of H ?
- What is $\text{Nul}(H)$? If the signal x belongs to the nullspace, what signal y will be received?

We want to understand

$$y = Hx.$$

- What is the first column of H ?
- What is $\text{Nul}(H)$? If the signal x belongs to the nullspace, what signal y will be received?

We want to understand

$$y = Hx.$$

- What is the first column of H ?
- What is $\text{Nul}(H)$? If the signal x belongs to the nullspace, what signal y will be received?
- In a well designed system you want $\text{Dim}(\text{Nul}(H)) = ?$

We want to understand

$$y = Hx.$$

- What is the first column of H ?
- What is $\text{Nul}(H)$? If the signal x belongs to the nullspace, what signal y will be received?
- In a well designed system you want $\text{Dim}(\text{Nul}(H)) = ?$

We want to understand

$$y = Hx.$$

- What is the first column of H ?
- What is $\text{Nul}(H)$? If the signal x belongs to the nullspace, what signal y will be received?
- In a well designed system you want $\text{Dim}(\text{Nul}(H)) = ?$
- What is $\text{Col}(H)$?

When B receives the signal, she wants to reconstruct the vector \mathbf{x} .
Optimally, she would just solve the linear system

$$H\mathbf{x} = \mathbf{y}.$$

Unfortunately, almost always B received $\mathbf{y} + \epsilon$ instead of \mathbf{y} , where $\epsilon \in \mathbb{R}^m$ is noise.

So B would try to solve

$$H\mathbf{x} = \mathbf{y} + \epsilon.$$

instead. However, that system might not have a solution. So B has to find the least square solution! Because B receives many messages from A, she will have to find the least square solution many times. Luckily, H does not change, and has independent columns ($\text{Nul}(H) = 0$). So B determines the QR -decomposition of H

$$H = QR,$$

once, and then just solves

$$R\mathbf{x} = Q^T(\mathbf{y} + \epsilon)$$

each time she receives a new message. This is easy to do, since R is upper triangular.

Linear transformation revisited

Recall the notion of coordinate vectors.

Recall the notion of coordinate vectors. If $\mathcal{B} = \{b_1, b_2\}$ is a basis for \mathbb{R}^2 , and x some vector then the coordinate vector of x is $x_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$ precisely if $x = c_1 b_1 + c_2 b_2$.

Recall the notion of coordinate vectors. If $\mathcal{B} = \{b_1, b_2\}$ is a basis for \mathbb{R}^2 , and x some vector then the coordinate vector of x is

$x_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$ precisely if $x = c_1 b_1 + c_2 b_2$. We want to understand

how to relate coordinate vectors $x_{\mathcal{B}}$ and $x_{\mathcal{C}}$ for different bases \mathcal{B} and \mathcal{C} .

Recall the notion of coordinate vectors. If $\mathcal{B} = \{b_1, b_2\}$ is a basis for \mathbb{R}^2 , and x some vector then the coordinate vector of x is

$x_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$ precisely if $x = c_1 b_1 + c_2 b_2$. We want to understand

how to relate coordinate vectors $x_{\mathcal{B}}$ and $x_{\mathcal{C}}$ for different bases \mathcal{B} and \mathcal{C} . We will see that there is for every two bases a matrix $I_{\mathcal{C}, \mathcal{B}}$ so that

$$x_{\mathcal{C}} = I_{\mathcal{C}, \mathcal{B}} x_{\mathcal{B}}.$$

Remember Theorem 1 of Lecture 17? Here it is again.

Remember Theorem 1 of Lecture 17? Here it is again.

Theorem

Let \mathcal{B} be a basis of \mathbb{R}^m and \mathcal{C} be a basis of \mathbb{R}^n and let $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear transformation. Then there is a $n \times m$ matrix $T_{\mathcal{C},\mathcal{B}}$ such that for every $\mathbf{v} \in \mathbb{R}^m$

$$T(\mathbf{v})_{\mathcal{C}} = T_{\mathcal{C},\mathcal{B}}\mathbf{v}_{\mathcal{B}}.$$

and

$$T_{\mathcal{C},\mathcal{B}} = [T(\mathbf{v}_1)_{\mathcal{C}} \quad T(\mathbf{v}_2)_{\mathcal{C}} \quad \dots \quad T(\mathbf{v}_m)_{\mathcal{C}}]$$

where $\mathcal{B} = (\mathbf{v}_1; \dots; \mathbf{v}_m)$.

Remember Theorem 1 of Lecture 17? Here it is again.

Theorem

Let \mathcal{B} be a basis of \mathbb{R}^m and \mathcal{C} be a basis of \mathbb{R}^n and let $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear transformation. Then there is a $n \times m$ matrix $T_{\mathcal{C},\mathcal{B}}$ such that for every $\mathbf{v} \in \mathbb{R}^m$

$$T(\mathbf{v})_{\mathcal{C}} = T_{\mathcal{C},\mathcal{B}}\mathbf{v}_{\mathcal{B}}.$$

and

$$T_{\mathcal{C},\mathcal{B}} = [T(\mathbf{v}_1)_{\mathcal{C}} \quad T(\mathbf{v}_2)_{\mathcal{C}} \quad \dots \quad T(\mathbf{v}_m)_{\mathcal{C}}]$$

where $\mathcal{B} = (\mathbf{v}_1; \dots; \mathbf{v}_m)$.

We will use this first in the special case $T = I$, where $I(v) = v$ (seemingly boring!).

Example

Consider $\mathcal{E} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{B} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$. Let $I : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation (the Identity!)

$$I\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x \\ y \end{bmatrix}.$$

Find the matrix $I_{\mathcal{E}, \mathcal{B}}$ that represents I with respect to the input basis \mathcal{B} and output basis \mathcal{E} .

Example

Consider $\mathcal{E} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{B} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$. Let $I : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation (the Identity!)

$$I\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x \\ y \end{bmatrix}.$$

Find the matrix $I_{\mathcal{E}, \mathcal{B}}$ that represents I with respect to the input basis \mathcal{B} and output basis \mathcal{E} .

Solution

By definition the matrix $I_{\mathcal{E}, \mathcal{B}}$ has as first column b_1 expressed in the standard basis, and as second column b_2 also expressed in the standard basis.

Example

Consider $\mathcal{E} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{B} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$. Let $I : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation (the Identity!)

$$I\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x \\ y \end{bmatrix}.$$

Find the matrix $I_{\mathcal{E}, \mathcal{B}}$ that represents I with respect to the input basis \mathcal{B} and output basis \mathcal{E} .

Solution

By definition the matrix $I_{\mathcal{E}, \mathcal{B}}$ has as first column b_1 expressed in the standard basis, and as second column b_2 also expressed in the standard basis. But for any vector $x \in \mathbb{R}^n$ we have $x_{\mathcal{E}} = x$! So

Example

Consider $\mathcal{E} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{B} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$. Let $I : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation (the Identity!)

$$I\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x \\ y \end{bmatrix}.$$

Find the matrix $I_{\mathcal{E}, \mathcal{B}}$ that represents I with respect to the input basis \mathcal{B} and output basis \mathcal{E} .

Solution

By definition the matrix $I_{\mathcal{E}, \mathcal{B}}$ has as first column b_1 expressed in the standard basis, and as second column b_2 also expressed in the standard basis. But for any vector $x \in \mathbb{R}^n$ we have $x_{\mathcal{E}} = x$! So

$$I_{\mathcal{E}, \mathcal{B}} = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} = [b_1 \quad b_2].$$

Example

Given $\mathbf{v} \in \mathbb{R}^2$ what is $I_{\mathcal{E}, \mathcal{B}} \mathbf{v}_{\mathcal{B}}$?

Example

Given $\mathbf{v} \in \mathbb{R}^2$ what is $l_{\mathcal{E}, \mathcal{B}} \mathbf{v}_{\mathcal{B}}$?

Solution

Let $\mathbf{v}_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$. Then

$$l_{\mathcal{E}, \mathcal{B}} \mathbf{v}_{\mathcal{B}} = \begin{bmatrix} b_1 & b_2 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = c_1 b_1 + c_2 b_2 = v!$$

Suppose \mathbf{v} is a vector in \mathbb{R}^n , and we have two bases in \mathbb{R}^n . so that we get two coordinate vectors \mathbf{v}_C and \mathbf{v}_B .

Suppose \mathbf{v} is a vector in \mathbb{R}^n , and we have two bases in \mathbb{R}^n . so that we get two coordinate vectors \mathbf{v}_C and \mathbf{v}_B . How are they related?

Suppose \mathbf{v} is a vector in \mathbb{R}^n , and we have two bases in \mathbb{R}^n . so that we get two coordinate vectors \mathbf{v}_C and \mathbf{v}_B . How are they related?

Theorem

Let \mathcal{B} be a basis of \mathbb{R}^n and \mathcal{C} be another basis of \mathbb{R}^n and let $I : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the linear transformation such that $I(\mathbf{v}) = \mathbf{v}$ for every $\mathbf{v} \in \mathbb{R}^n$. Then

$$\mathbf{v}_C = I_{C,B} \mathbf{v}_B.$$

Suppose \mathbf{v} is a vector in \mathbb{R}^n , and we have two bases in \mathbb{R}^n . so that we get two coordinate vectors \mathbf{v}_C and \mathbf{v}_B . How are they related?

Theorem

Let \mathcal{B} be a basis of \mathbb{R}^n and \mathcal{C} be another basis of \mathbb{R}^n and let $I : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the linear transformation such that $I(\mathbf{v}) = \mathbf{v}$ for every $\mathbf{v} \in \mathbb{R}^n$. Then

$$\mathbf{v}_C = I_{C,B} \mathbf{v}_B.$$

We call the matrix $I_{C,B}$ a **change of base matrix**, it transforms coordinate vectors from the \mathcal{B} to the \mathcal{C} basis.

Example

Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{B} be another basis of \mathbb{R}^n . What is $I_{\mathcal{E},\mathcal{B}}$?

Solution

The columns of $I_{\mathcal{E},\mathcal{B}}$ are the basic vectors b_1, b_2, \dots expressed in the standard basis. So

$$I_{\mathcal{E},\mathcal{B}} = [b_1 \quad b_2 \quad \dots \quad b_n]$$

Example

Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{B} be another basis of \mathbb{R}^n . What is $I_{\mathcal{E},\mathcal{B}}$?

Solution

The columns of $I_{\mathcal{E},\mathcal{B}}$ are the basic vectors b_1, b_2, \dots expressed in the standard basis. So

$$I_{\mathcal{E},\mathcal{B}} = [b_1 \quad b_2 \quad \dots \quad b_n]$$

So this is the **easy** change of basis matrix: you just write down the \mathcal{B} basis as columns of your matrix.

Example

Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{B} be another basis of \mathbb{R}^n . What is $I_{\mathcal{E},\mathcal{B}}$?

Solution

The columns of $I_{\mathcal{E},\mathcal{B}}$ are the basic vectors b_1, b_2, \dots expressed in the standard basis. So

$$I_{\mathcal{E},\mathcal{B}} = [b_1 \quad b_2 \quad \dots \quad b_n]$$

So this is the **easy** change of basis matrix: you just write down the \mathcal{B} basis as columns of your matrix. It has the property that

$$v = v_{\mathcal{E}} = I_{\mathcal{E},\mathcal{B}} v_{\mathcal{B}}$$

Example

Let \mathcal{B} be a basis of \mathbb{R}^n and \mathcal{C} be a basis of \mathbb{R}^n . What is $I_{\mathcal{C},\mathcal{B}}^{-1}$?

Example

Let \mathcal{B} be a basis of \mathbb{R}^n and \mathcal{C} be a basis of \mathbb{R}^n . What is $I_{\mathcal{C},\mathcal{B}}^{-1}$?

Solution

$I_{\mathcal{C},\mathcal{B}}$ is the matrix with columns the \mathcal{B} basis vectors expressed in the \mathcal{C} basis, and $I_{\mathcal{C},\mathcal{B}}^{-1}$ is the inverse of this matrix. These matrices have the property that

$$v_{\mathcal{C}} = I_{\mathcal{C},\mathcal{B}} v_{\mathcal{B}}, \quad v_{\mathcal{B}} = I_{\mathcal{C},\mathcal{B}}^{-1} v_{\mathcal{C}}.$$

Example

As before, let $\mathcal{E} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{B} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$. What is $I_{\mathcal{B}, \mathcal{E}}$?

Example

As before, let $\mathcal{E} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{B} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$. What is $I_{\mathcal{B},\mathcal{E}}$?

Solution

We know what $I_{\mathcal{E},\mathcal{B}}$ is, it is just $\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$. Then $I_{\mathcal{B},\mathcal{E}}$ is the transition matrix going the other way, so it is the inverse of the **easy** matrix, so

$$I_{\mathcal{B},\mathcal{E}} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}.$$

Example

As before, let $\mathcal{E} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{B} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$. What is $I_{\mathcal{B},\mathcal{E}}$?

Solution

We know what $I_{\mathcal{E},\mathcal{B}}$ is, it is just $\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$. Then $I_{\mathcal{B},\mathcal{E}}$ is the transition matrix going the other way, so it is the inverse of the **easy** matrix, so

$$I_{\mathcal{B},\mathcal{E}} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}.$$

Example

Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{C} be a orthonormal basis of \mathbb{R}^n . Then $I_{\mathcal{C},\mathcal{E}} = I_{\mathcal{E},\mathcal{C}}^T$. Why?

Example

Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{C} be a orthonormal basis of \mathbb{R}^n . Then $I_{\mathcal{C},\mathcal{E}} = I_{\mathcal{E},\mathcal{C}}^T$. Why?

Solution

$I_{\mathcal{E},\mathcal{C}}$ the matrix with orthonormal columns, so it is an orthogonal matrix.

Example

Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{C} be a orthonormal basis of \mathbb{R}^n . Then $I_{\mathcal{C},\mathcal{E}} = I_{\mathcal{E},\mathcal{C}}^T$. Why?

Solution

$I_{\mathcal{E},\mathcal{C}}$ the matrix with orthonormal columns, so it is an orthogonal matrix. $I_{\mathcal{C},\mathcal{E}}$ is the inverse.

Example

Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{C} be a orthonormal basis of \mathbb{R}^n . Then $I_{\mathcal{C},\mathcal{E}} = I_{\mathcal{E},\mathcal{C}}^T$. Why?

Solution

$I_{\mathcal{E},\mathcal{C}}$ the matrix with orthonormal columns, so it is an orthogonal matrix. $I_{\mathcal{C},\mathcal{E}}$ is the inverse. But the inverse of an orthogonal matrix is easy, just the transpose.

Theorem

Let $\mathcal{B} := (\mathbf{u}_1, \dots, \mathbf{u}_n)$ be an orthonormal basis of \mathbb{R}^n and $U = [\mathbf{u}_1 \ \dots \ \mathbf{u}_n]$. Then for every $\mathbf{v} \in \mathbb{R}^n$

$$v_{\mathcal{B}} = U^T \mathbf{v}.$$

Change of basis

Theorem

Let $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear transformation and \mathcal{A} and \mathcal{B} be two bases of \mathbb{R}^m and \mathcal{C}, \mathcal{D} be two bases of \mathbb{R}^n . Then

$$T_{\mathcal{C}, \mathcal{A}} = I_{\mathcal{C}, \mathcal{D}} T_{\mathcal{D}, \mathcal{B}} I_{\mathcal{B}, \mathcal{A}}.$$

$$\begin{array}{ccc} (\mathbb{R}^m, \mathcal{A}) & \xrightarrow{\text{apply } T_{\mathcal{C}, \mathcal{A}}} & (\mathbb{R}^n, \mathcal{C}) \\ I_{\mathcal{B}, \mathcal{A}} \downarrow & & \uparrow I_{\mathcal{C}, \mathcal{D}} \\ (\mathbb{R}^m, \mathcal{B}) & \xrightarrow{\text{apply } T_{\mathcal{D}, \mathcal{B}}} & (\mathbb{R}^n, \mathcal{D}) \end{array}$$

Example

Consider $\mathcal{B} := \mathcal{D} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{A} := \mathcal{C} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ as before. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be again the linear transformation that

$$\begin{bmatrix} x \\ y \end{bmatrix} \mapsto A \begin{bmatrix} x \\ y \end{bmatrix}, \text{ where } A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}.$$

Determine $T_{\mathcal{C}, \mathcal{A}}$.

Example

Consider $\mathcal{B} := \mathcal{D} := \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{A} := \mathcal{C} := \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ as before. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be again the linear transformation that

$$\begin{bmatrix} x \\ y \end{bmatrix} \mapsto A \begin{bmatrix} x \\ y \end{bmatrix}, \text{ where } A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}.$$

Determine $T_{\mathcal{C}, \mathcal{A}}$.

Solution

Example

Let \mathcal{E} be the standard basis of \mathbb{R}^n , let $\mathcal{B} := (\mathbf{u}_1, \dots, \mathbf{u}_n)$ be an orthonormal basis of \mathbb{R}^n and $U = [\mathbf{u}_1 \ \dots \ \mathbf{u}_n]$. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation. Then

$$T_{\mathcal{B}, \mathcal{B}} = U^T T_{\mathcal{E}, \mathcal{E}} U.$$

Why?

Solution