

# Math 415 - Lecture 1

## Introduction

Monday August 24 2015

- Textbook: Chapter 1.3, Chapter 2.2 (just the pages 78 and 79)

- Textbook: Chapter 1.3, Chapter 2.2 (just the pages 78 and 79)
- Suggested Practice Exercise: in Chapter 1.3, Exercise 1, 3, 5, 6, 11

- Textbook: Chapter 1.3, Chapter 2.2 (just the pages 78 and 79)
- Suggested Practice Exercise: in Chapter 1.3, Exercise 1, 3, 5, 6, 11
- Khan Academy Video: Matrices: Reduced Row Echelon Form  
1

## Systems of Linear Equations

## Definition

A **linear equation** is a equation of the form

$$a_1x_1 + \dots + a_nx_n = b$$

where  $a_1, \dots, a_n, b$  are numbers and  $x_1, \dots, x_n$  are variables.

## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$$4x_1 - 5x_2 + 2 = x_1$$

## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$$4x_1 - 5x_2 + 2 = x_1 \quad | \quad 3x_1 - 5x_2 = -2 \quad \text{Linear.}$$



## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$$\begin{array}{l|l} 4x_1 - 5x_2 + 2 = x_1 & 3x_1 - 5x_2 = -2 \\ x_2 = 2(\sqrt{6} - x_1) + x_3 & \end{array} \quad \text{Linear.}$$

## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$$\begin{array}{l|l} 4x_1 - 5x_2 + 2 = x_1 & 3x_1 - 5x_2 = -2 \quad \text{Linear.} \\ x_2 = 2(\sqrt{6} - x_1) + x_3 & 2x_1 + x_2 - x_3 = 2\sqrt{6} \quad \text{Linear.} \end{array}$$

## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$$\begin{array}{l|l} 4x_1 - 5x_2 + 2 = x_1 & 3x_1 - 5x_2 = -2 \quad \text{Linear.} \\ x_2 = 2(\sqrt{6} - x_1) + x_3 & 2x_1 + x_2 - x_3 = 2\sqrt{6} \quad \text{Linear.} \\ 4x_1 - 6x_2 = x_1x_2 & \end{array}$$

## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$4x_1 - 5x_2 + 2 = x_1$	$3x_1 - 5x_2 = -2$	<i>Linear.</i>
$x_2 = 2(\sqrt{6} - x_1) + x_3$	$2x_1 + x_2 - x_3 = 2\sqrt{6}$	<i>Linear.</i>
$4x_1 - 6x_2 = x_1x_2$	$4x_1 - 6x_2 = \underline{x_1x_2}$	<i>Not linear.</i>

## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$$4x_1 - 5x_2 + 2 = x_1$$

$$x_2 = 2(\sqrt{6} - x_1) + x_3$$

$$4x_1 - 6x_2 = x_1x_2$$

$$x_2 = 2\sqrt{x_1} - 7$$

$$3x_1 - 5x_2 = -2$$

$$2x_1 + x_2 - x_3 = 2\sqrt{6}$$

$$4x_1 - 6x_2 = \underline{x_1x_2}$$

*Linear.*

*Linear.*

*Not linear.*

## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$$4x_1 - 5x_2 + 2 = x_1$$

$$x_2 = 2(\sqrt{6} - x_1) + x_3$$

$$4x_1 - 6x_2 = x_1x_2$$

$$x_2 = 2\sqrt{x_1} - 7$$

$$3x_1 - 5x_2 = -2$$

$$2x_1 + x_2 - x_3 = 2\sqrt{6}$$

$$4x_1 - 6x_2 = \underline{x_1x_2}$$

$$x_2 = \underline{2\sqrt{x_1}} - 7$$

*Linear.*

*Linear.*

*Not linear.*

*Not linear.*

## Example

Which of the following equations are linear equations (or can be rearranged to become linear equations)?

$$4x_1 - 5x_2 + 2 = x_1$$

$$x_2 = 2(\sqrt{6} - x_1) + x_3$$

$$4x_1 - 6x_2 = x_1x_2$$

$$x_2 = 2\sqrt{x_1} - 7$$

$$3x_1 - 5x_2 = -2$$

$$2x_1 + x_2 - x_3 = 2\sqrt{6}$$

$$4x_1 - 6x_2 = \underline{x_1x_2}$$

$$x_2 = \underline{2\sqrt{x_1}} - 7$$

*Linear.*

*Linear.*

*Not linear.*

*Not linear.*

This course will focus on linear equations.

## Definition

A **system of linear equations** (or a **linear system**) is a collection of one or more linear equations involving the same set of variables, say,  $x_1, x_2, \dots, x_n$ .



## Definition

A **system of linear equations** (or a **linear system**) is a collection of one or more linear equations involving the same set of variables, say,  $x_1, x_2, \dots, x_n$ .

### Definition

A **system of linear equations** (or a **linear system**) is a collection of one or more linear equations involving the same set of variables, say,  $x_1, x_2, \dots, x_n$ .

### Definition

A **solution** of a linear system is a list  $(s_1, s_2, \dots, s_n)$  of numbers that makes each equation in the system true when the values  $s_1, s_2, \dots, s_n$  are substituted for  $x_1, x_2, \dots, x_n$ , respectively.

### Definition

A **system of linear equations** (or a **linear system**) is a collection of one or more linear equations involving the same set of variables, say,  $x_1, x_2, \dots, x_n$ .

### Definition

A **solution** of a linear system is a list  $(s_1, s_2, \dots, s_n)$  of numbers that makes each equation in the system true when the values  $s_1, s_2, \dots, s_n$  are substituted for  $x_1, x_2, \dots, x_n$ , respectively.

### Definition

A **system of linear equations** (or a **linear system**) is a collection of one or more linear equations involving the same set of variables, say,  $x_1, x_2, \dots, x_n$ .

### Definition

A **solution** of a linear system is a list  $(s_1, s_2, \dots, s_n)$  of numbers that makes each equation in the system true when the values  $s_1, s_2, \dots, s_n$  are substituted for  $x_1, x_2, \dots, x_n$ , respectively.

### Definition

The **solution set** of a system of linear equations is the set of all possible solutions of a linear system.

## Example

Two equations in two variables:

$$x_1 + x_2 = 1$$

$$-x_1 + x_2 = 0.$$

What is a solution for this system of linear equations?

## Example

Two equations in two variables:

$$x_1 + x_2 = 1$$

$$-x_1 + x_2 = 0.$$

What is a solution for this system of linear equations?

Add them.  $2x_2 = 1 \Rightarrow x_2 = \frac{1}{2}$

### Example

Two equations in two variables:

$$x_1 + x_2 = 1$$

$$-x_1 + x_2 = 0.$$

What is a solution for this system of linear equations?

Add them.  $2x_2 = 1 \Rightarrow x_2 = .5$

Plug into first equation.  $x_1 + .5 = 1 \Rightarrow x_1 = .5$

### Example

Two equations in two variables:

$$x_1 + x_2 = 1$$

$$-x_1 + x_2 = 0.$$

What is a solution for this system of linear equations?

Add them.  $2x_2 = 1 \Rightarrow x_2 = .5$

Plug into first equation.  $x_1 + .5 = 1 \Rightarrow x_1 = .5$

$(x_1, x_2) = (.5, .5)$  is the only solution.



### Example

Does every system of linear equation have a solution?

$$x_1 - 2x_2 = -3$$

$$2x_1 - 4x_2 = 8.$$

### Example

Does every system of linear equation have a solution?

$$x_1 - 2x_2 = -3$$

$$2x_1 - 4x_2 = 8.$$

Multiply first equation by 2.  $2x_1 - 4x_2 = -6$

### Example

Does every system of linear equation have a solution?

$$x_1 - 2x_2 = -3$$

$$2x_1 - 4x_2 = 8.$$

Multiply first equation by 2.  $2x_1 - 4x_2 = -6$

Subtract from second equation.  $0 = 14$

### Example

Does every system of linear equation have a solution?

$$x_1 - 2x_2 = -3$$

$$2x_1 - 4x_2 = 8.$$

Multiply first equation by 2.  $2x_1 - 4x_2 = -6$

Subtract from second equation.  $0 = 14$

The equation  $0 = 14$  is always false, so no solutions exist.

### Example

How many solutions are there to the following system?

$$x_1 + x_2 = 3$$

$$-2x_1 - 2x_2 = -6$$

### Example

How many solutions are there to the following system?

$$x_1 + x_2 = 3$$

$$-2x_1 - 2x_2 = -6$$

Multiply first equation by 2.  $2x_1 + 2x_2 = 6$

### Example

How many solutions are there to the following system?

$$x_1 + x_2 = 3$$

$$-2x_1 - 2x_2 = -6$$

Multiply first equation by 2.  $2x_1 + 2x_2 = 6$

Add to second equation.  $0 = 0$

### Example

How many solutions are there to the following system?

$$x_1 + x_2 = 3$$

$$-2x_1 - 2x_2 = -6$$

Multiply first equation by 2.  $2x_1 + 2x_2 = 6$

Add to second equation.  $0 = 0$

Any value of  $x_1$  works.  $x_2 = 3 - x_1$ . Infinitely many solutions.



## Theorem

*This is all there is:*

## Theorem

*This is all there is:*

## Theorem

*This is all there is: A linear system has either*

**one unique solution** *or*

## Theorem

*This is all there is: A linear system has either*

**one unique solution or no solution or**

## Theorem

*This is all there is: A linear system has either*

**one unique solution or no solution or infinitely many solutions.**

Can you draw the set of solutions of the above equations?

$$x_1 + x_2 = 1$$

$$-x_1 + x_2 = 0.$$

$$x_1 - 2x_2 = -3$$

$$2x_1 - 4x_2 = 8.$$

$$x_1 + x_2 = 3$$

$$-2x_1 - 2x_2 = -6$$

Can you draw the set of solutions of the above equations?

$$x_1 + x_2 = 1$$


$$-x_1 + x_2 = 0.$$

$$x_1 - 2x_2 = -3$$

$$2x_1 - 4x_2 = 8.$$


$$x_1 + x_2 = 3$$

$$-2x_1 - 2x_2 = -6$$




Can you draw the set of solutions of the above equations?

$$\begin{aligned}x_1 + x_2 &= 1 \\ -x_1 + x_2 &= 0.\end{aligned}$$




$$\begin{aligned}x_1 - 2x_2 &= -3 \\ 2x_1 - 4x_2 &= 8.\end{aligned}$$






Can you draw the set of solutions of the above equations?


$$\begin{aligned}x_1 + x_2 &= 1 \\ -x_1 + x_2 &= 0.\end{aligned}$$



$$\begin{aligned}x_1 - 2x_2 &= -3 \\ 2x_1 - 4x_2 &= 8.\end{aligned}$$




$$\begin{aligned}x_1 + x_2 &= 3 \\ -2x_1 - 2x_2 &= -6.\end{aligned}$$




Can you draw the set of solutions of the above equations?


$$\begin{aligned}x_1 + x_2 &= 1 \\ -x_1 + x_2 &= 0.\end{aligned}$$



$$\begin{aligned}x_1 - 2x_2 &= -3 \\ 2x_1 - 4x_2 &= 8.\end{aligned}$$



$$\begin{aligned}x_1 + x_2 &= 3 \\ -2x_1 - 2x_2 &= -6.\end{aligned}$$



(The numbers in the graphs are not quite right.)

oooooooooo

**Take away:** Whenever you have a linear system with  $n$  equations, then the set of solutions of this system is precisely the intersection of the sets of solutions of each of the  $n$  equations on its own.

## Strategies for solving systems of linear equations

## Definition

Two systems are **equivalent** if they have the same solution set.

The general strategy is to replace one system with an equivalent system that is easier to solve.

### Example

Consider

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array}$$

The general strategy is to replace one system with an equivalent system that is easier to solve.

### Example

Consider

$$\begin{array}{rclcrcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array}$$

$$R2 \rightarrow R2 + R1 \quad \begin{array}{rclcrcl} x_1 & - & 2x_2 & = & -1 \\ 0 & + & x_2 & = & 2 \end{array}$$



The general strategy is to replace one system with an equivalent system that is easier to solve.

### Example

Consider

$$\begin{array}{rclcrcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array}$$

$$R2 \rightarrow R2 + R1 \quad \begin{array}{rclcrcl} x_1 & - & 2x_2 & = & -1 \\ 0 & + & x_2 & = & 2 \end{array}$$

$x_2 = 2$ , so  $x_1 = 3$ .

## Matrix Notation

## Matrix Notation

From a system of equations, we can get:

### Coefficient Matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array}$$

From a system of equations, we can get:

### Coefficient Matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \quad \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$

From a system of equations, we can get:

### Coefficient Matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \quad \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$

### Augmented matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array}$$

From a system of equations, we can get:

### Coefficient Matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \qquad \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$

### Augmented matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \qquad \left[ \begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right]$$

From a system of equations, we can get:

### Coefficient Matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \quad \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$

### Augmented matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \quad \left[ \begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right]$$

### $R2 \rightarrow R2 + R1$

$$\left[ \begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 1 & 2 \end{array} \right]$$



From a system of equations, we can get:

### Coefficient Matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \quad \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$

### Augmented matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \quad \left[ \begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right]$$

### $R2 \rightarrow R2 + R1$

$$\left[ \begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 1 & 2 \end{array} \right]$$

### $R1 \rightarrow R1 + 2R2$

$$\left[ \begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \end{array} \right]$$

From a system of equations, we can get:

### Coefficient Matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \quad \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$

### Augmented matrix

$$\begin{array}{rclcl} x_1 & - & 2x_2 & = & -1 \\ -x_1 & + & 3x_2 & = & 3 \end{array} \quad \left[ \begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right]$$

### $R2 \rightarrow R2 + R1$

$$\left[ \begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 1 & 2 \end{array} \right]$$

### $R1 \rightarrow R1 + 2R2$

$$\left[ \begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \end{array} \right]$$

**Solution:**  $x_1 = 3, x_2 = 2$

## Definition

An **elementary row operation** is one of the following

## Definition

An **elementary row operation** is one of the following

(Replacement) Add a multiple of one row to another row,

## Definition

An **elementary row operation** is one of the following

(Replacement) Add a multiple of one row to another row,

(Interchange) Interchange two rows, or

## Definition

An **elementary row operation** is one of the following

(Replacement) Add a multiple of one row to another row,

(Interchange) Interchange two rows, or

(Scaling) Multiply all entries in a row by a nonzero constant.

## Definition

An **elementary row operation** is one of the following

(Replacement) Add a multiple of one row to another row,

(Interchange) Interchange two rows, or

(Scaling) Multiply all entries in a row by a nonzero constant.

## Definition

An **elementary row operation** is one of the following

(Replacement) Add a multiple of one row to another row,

(Interchange) Interchange two rows, or

(Scaling) Multiply all entries in a row by a nonzero constant.

## Definition

Two matrices are **row equivalent**, if one matrix can be transformed into the other matrix by a sequence of elementary row operations.



## Definition

An **elementary row operation** is one of the following

(Replacement) Add a multiple of one row to another row,

(Interchange) Interchange two rows, or

(Scaling) Multiply all entries in a row by a nonzero constant.

## Definition

Two matrices are **row equivalent**, if one matrix can be transformed into the other matrix by a sequence of elementary row operations.

## Definition

An **elementary row operation** is one of the following

(Replacement) Add a multiple of one row to another row,

(Interchange) Interchange two rows, or

(Scaling) Multiply all entries in a row by a nonzero constant.

## Definition

Two matrices are **row equivalent**, if one matrix can be transformed into the other matrix by a sequence of elementary row operations.

## Theorem

*If the augmented matrices of two linear systems are row equivalent, then the two systems have the same solution set.*

## Example

Solve the following system (or show there is no solution):

$$\begin{aligned}x_1 - 2x_2 + x_3 &= 0 \\2x_2 - 8x_3 &= 8 \\-4x_1 + 5x_2 + 9x_3 &= -9\end{aligned}$$

$$\begin{aligned}x_1 - 2x_2 + x_3 &= 0 \\2x_2 - 8x_3 &= 8 \\-4x_1 + 5x_2 + 9x_3 &= -9\end{aligned} \quad \left[ \begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ -4 & 5 & 9 & -9 \end{array} \right]$$

---


$$\begin{aligned}x_1 - 2x_2 + x_3 &= 0 \\2x_2 - 8x_3 &= 8 \\-3x_2 + 13x_3 &= -9\end{aligned} \quad \left[ \begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ 0 & -3 & 13 & -9 \end{array} \right]$$


---

$$\begin{array}{rcl}
 x_1 - 2x_2 + x_3 & = & 0 \\
 x_2 - 4x_3 & = & 4 \\
 -3x_2 + 13x_3 & = & -9
 \end{array}
 \quad
 \left[ \begin{array}{ccc|c}
 1 & -2 & 1 & 0 \\
 0 & 1 & -4 & 4 \\
 0 & -3 & 13 & -9
 \end{array} \right]$$


---

$$\begin{array}{rcl}
 x_1 - 2x_2 + x_3 & = & 0 \\
 x_2 - 4x_3 & = & 4 \\
 x_3 & = & 3
 \end{array}
 \quad
 \left[ \begin{array}{ccc|c}
 1 & -2 & 1 & 0 \\
 0 & 1 & -4 & 4 \\
 0 & 0 & 1 & 3
 \end{array} \right]$$


---

$$\begin{array}{rcl}
 x_1 - 2x_2 & = & -3 \\
 x_2 & = & 16 \\
 x_3 & = & 3
 \end{array}
 \quad
 \left[ \begin{array}{ccc|c}
 1 & -2 & 0 & -3 \\
 0 & 1 & 0 & 16 \\
 0 & 0 & 1 & 3
 \end{array} \right]$$


---

$$\begin{array}{rcl}
 x_1 & = & 29 \\
 x_2 & = & 16 \\
 x_3 & = & 3
 \end{array}
 \quad
 \left[ \begin{array}{ccc|c}
 1 & 0 & 0 & 29 \\
 0 & 1 & 0 & 16 \\
 0 & 0 & 1 & 3
 \end{array} \right]$$

**Solution:**  $(29, 16, 3)$

**Check:** Is  $(29, 16, 3)$  a solution of the **original** system?

$$\begin{array}{rcccccccl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & 2x_2 & - & 8x_3 & = & 8 \\ -4x_1 & + & 5x_2 & + & 9x_3 & = & -9 \end{array}$$

**Solution:**  $(29, 16, 3)$

**Check:** Is  $(29, 16, 3)$  a solution of the **original** system?

$$\begin{array}{rcccccccl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & 2x_2 & - & 8x_3 & = & 8 \\ -4x_1 & + & 5x_2 & + & 9x_3 & = & -9 \end{array}$$

$$\begin{array}{rcccccccl} 29 & - & 32 & + & 3 & = & 0 & \checkmark \\ & & 32 & - & 24 & = & 8 & \checkmark \\ -116 & + & 80 & + & 27 & = & -9 & \checkmark \end{array}$$

## Two Fundamental Questions (Existence and Uniqueness)

## Two Fundamental Questions (Existence and Uniqueness)



There are two fundamental question about linear equation:

(1) Is the system consistent? (I.e. does a solution **exist**?)

There are two fundamental question about linear equation:

- (1) Is the system consistent? (I.e. does a solution **exist**?)
- (2) If a solution exists, is it **unique**? (I.e. is there one only one solution?)

### Example

Is this system consistent? If so, is the solution unique?

$$\begin{array}{rcccccccl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & 2x_2 & - & 8x_3 & = & 8 \\ -4x_1 & + & 5x_2 & + & 9x_3 & = & -9 \end{array}$$

In the last example, this system was reduced to the triangular form:

$$\begin{array}{rcccccl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & x_2 & - & 4x_3 & = & 4 \\ & & & & x_3 & = & 3 \end{array} \quad \left[ \begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right]$$

This is sufficient to see that the system is consistent and unique.  
Why?

In the last example, this system was reduced to the triangular form:

$$\begin{array}{rclclcl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & x_2 & - & 4x_3 & = & 4 \\ & & & & x_3 & = & 3 \end{array} \quad \left[ \begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right]$$

This is sufficient to see that the system is consistent and unique. Why?

- The last row determines  $x_3$  uniquely.

In the last example, this system was reduced to the triangular form:

$$\begin{array}{rclclcl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & x_2 & - & 4x_3 & = & 4 \\ & & & & x_3 & = & 3 \end{array} \quad \left[ \begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right]$$

This is sufficient to see that the system is consistent and unique. Why?

- The last row determines  $x_3$  uniquely.
- Knowing  $x_3$ , the second row determines  $x_2$  uniquely.

In the last example, this system was reduced to the triangular form:

$$\begin{array}{rrcrcl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & x_2 & - & 4x_3 & = & 4 \\ & & & & x_3 & = & 3 \end{array} \quad \left[ \begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right]$$

This is sufficient to see that the system is consistent and unique. Why?

- The last row determines  $x_3$  uniquely.
- Knowing  $x_3$ , the second row determines  $x_2$  uniquely.
- Knowing  $x_2$  and  $x_3$ , the first row determines  $x_1$  uniquely.
- So, exactly one possible solution  $(x_1, x_2, x_3)$ .

### Example

Is this system consistent?

$$\begin{array}{rcl} 3x_2 - 6x_3 & = & 8 \\ x_1 - 2x_2 + 3x_3 & = & -1 \\ 5x_1 - 7x_2 + 9x_3 & = & 0 \end{array} \quad \left[ \begin{array}{ccc|c} 0 & 3 & -6 & 8 \\ 1 & -2 & 3 & -1 \\ 5 & -7 & 9 & 0 \end{array} \right]$$



## Solution:

**Solution:**

$$\xrightarrow{R1 \leftrightarrow R2} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 5 & -7 & 9 & 0 \end{array} \right]$$

**Solution:**

$$\xrightarrow{R1 \leftrightarrow R2} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 5 & -7 & 9 & 0 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R3 - 5R1} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 3 & -6 & 5 \end{array} \right]$$

**Solution:**

$$\xrightarrow{R1 \leftrightarrow R2} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 5 & -7 & 9 & 0 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R3 - 5R1} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 3 & -6 & 5 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R3 - R2} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 0 & 0 & -3 \end{array} \right]$$

**Solution:**

$$\xrightarrow{R1 \leftrightarrow R2} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 5 & -7 & 9 & 0 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R3 - 5R1} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 3 & -6 & 5 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R3 - R2} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 0 & 0 & -3 \end{array} \right]$$

Equation notation of triangular form:

$$\begin{array}{rclcl} x_1 & -2x_2 & +3x_3 & = & -1 \\ & 3x_2 & -6x_3 & = & 8 \\ & & 0 & = & -3 \end{array}$$

**Solution:**

$$\xrightarrow{R1 \leftrightarrow R2} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 5 & -7 & 9 & 0 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R3 - 5R1} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 3 & -6 & 5 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R3 - R2} \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 0 & 3 & -6 & 8 \\ 0 & 0 & 0 & -3 \end{array} \right]$$

Equation notation of triangular form:

$$\begin{array}{rclcl} x_1 & -2x_2 & +3x_3 & = & -1 \\ & 3x_2 & -6x_3 & = & 8 \\ & & 0 & = & -3 \end{array}$$

The original system is inconsistent!

### Example

For what values of  $h$  will the following system be consistent?

$$\begin{array}{rclcl} 3x_1 & - & 9x_2 & = & 4 \\ -2x_1 & + & 6x_2 & = & h \end{array}$$

**Solution:**

$$\left[ \begin{array}{cc|c} 3 & -9 & 4 \\ -2 & 6 & h \end{array} \right]$$



**Solution:**

$$R1 \xrightarrow{\rightarrow \frac{1}{3} R1} \left[ \begin{array}{cc|c} 3 & -9 & 4 \\ -2 & 6 & h \\ 1 & -3 & \frac{4}{3} \\ -2 & 6 & h \end{array} \right]$$

**Solution:**

$$\begin{array}{l} \xrightarrow{R1 \rightarrow \frac{1}{3}R1} \\ \xrightarrow{R2 \rightarrow R2 + 2R1} \end{array} \left[ \begin{array}{cc|c} 3 & -9 & 4 \\ -2 & 6 & h \\ 1 & -3 & \frac{4}{3} \\ -2 & 6 & h \\ 1 & -3 & \frac{4}{3} \\ 0 & 0 & h + \frac{8}{3} \end{array} \right]$$

**Solution:**

$$\begin{array}{l} \xrightarrow{R1 \rightarrow \frac{1}{3}R1} \\ \xrightarrow{R2 \rightarrow R2 + 2R1} \end{array} \left[ \begin{array}{cc|c} 3 & -9 & 4 \\ -2 & 6 & h \\ 1 & -3 & \frac{4}{3} \\ -2 & 6 & h \\ 1 & -3 & \frac{4}{3} \\ 0 & 0 & h + \frac{8}{3} \end{array} \right]$$

System is consistent if and only if  $h = -\frac{8}{3}$ .