

Math 415 - Lecture 36

Minima, maxima and saddle points, Constrained Optimization

Wednesday December 2nd 2015

Textbook reading: Chapter 6.1

Suggested practice exercises: Chapter 6.1, # 1, 16

Strang lecture: Lecture 27: Positive definite matrices and minima

1 Review

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.
- So, we can write

$$A = QDQ^T$$

where

$$D = \underbrace{\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \quad \text{and} \quad Q = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{q}_1 & \cdots & \mathbf{q}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}}$$

- A is called **positive definite** if $\mathbf{x}^T A \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then
 1. If all $\lambda_i > 0$, then A is positive definite,
 2. If all $\lambda_i < 0$, then $\mathbf{x}^T A \mathbf{x} < 0$ for all $\mathbf{x} \neq \mathbf{0}$
 3. If some $\lambda_i > 0$, some $\lambda_j < 0$, $\mathbf{x}^T A \mathbf{x}$ will have both positive and negative values.
- a function of the form $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ is called a **quadratic form**.

2 2nd derivative test

Definition 1. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable, the **Hessian** matrix of f

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(\mathbf{0}) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_2^2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n}(\mathbf{0}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_n \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_n^2}(\mathbf{0}) \end{bmatrix}$$

Idea. If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable and $\mathbf{0}$ is a critical point, then $f(\mathbf{x}) \approx f(\mathbf{0}) + \frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$.

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function, $\frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$!
- We understand $q(\mathbf{x}) = \frac{1}{2}\mathbf{x} \cdot H\mathbf{x} \implies$ we understand if $\mathbf{0}$ is a max, min or neither for f !
- Turns out: $q(\mathbf{x})$ is determined by eigenvectors and eigenvalues of H !

Theorem 1 (2nd derivative test). *If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$, then*

1. *If all eigenvalues of H are **positive**, then $\mathbf{0}$ is a local **min**.*
2. *If all eigenvalues of H are **negative**, then $\mathbf{0}$ is a local **max**.*
3. *If one eigenvalue of H is **positive** and one is **negative**, then $\mathbf{0}$ is **neither** a max nor a min.*
4. *Otherwise (e.g. all eigenvalues positive or zero), no information!*

Example 2. Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

(An example of such a function is $f(x, y) = \frac{1}{2}x^2 + 2xy + \frac{1}{2}y^2$).

Solution.





Figure 1: Graph of the function $f(x, y) = \frac{1}{2}x^2 + 2xy + \frac{1}{2}y^2$

Example 3. Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution.




Figure 2: Graph of the function $f(x, y) = \frac{1}{2}x^2 - xy + \frac{3}{2}y^2$

3 Constrained optimization

Problem: Given a quadratic form $q(\mathbf{x})$, find the maximum or minimum value $q(\mathbf{x})$ for \mathbf{x} in some specified set. Typically, the problem can be arranged such that \mathbf{x} varies over the set of vectors with $\mathbf{x}^T \mathbf{x} = 1$.

Example 4. Let $A = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 3 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution.



What if A is not diagonal?

Theorem 2. Let A be a symmetric matrix and let λ_m be the least eigenvalue and λ_M be the greatest eigenvalue of A . Then


$$\lambda_m = \min\{\mathbf{x}^T A \mathbf{x} : \mathbf{x}^T \mathbf{x} = 1\},$$

moreover if \mathbf{u}_m is a unit eigenvector corresponding to λ_m , then $\mathbf{u}_m^T A \mathbf{u}_m = \lambda_m$.
In addition,

$$\lambda_M = \max\{\mathbf{x}^T A \mathbf{x} : \mathbf{x}^T \mathbf{x} = 1\},$$

moreover if \mathbf{u}_M is a unit eigenvector corresponding to λ_M , then $\mathbf{u}_M^T A \mathbf{u}_M = \lambda_M$.

Proof.



Example 5. Let $A = \begin{bmatrix} 3 & 2 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 4 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution.