

Math 415 - Lecture 20

Fundamental Theorem of Linear algebra, orthogonal complement
of fundamental subspaces of a matrix

Monday October 12th 2015

Textbook reading: Chapter 3.1

Textbook reading: Chapter 3.1

Suggested practice exercises: Chapter 2.6, 5,6,7,36,37

Textbook reading: Chapter 3.1

Suggested practice exercises: Chapter 2.6, 5,6,7,36,37

Khan Academy video: Orthogonal complements

Textbook reading: Chapter 3.1

Suggested practice exercises: Chapter 2.6, 5,6,7,36,37

Khan Academy video: Orthogonal complements

Strang lecture: Lecture 14: Orthogonal vectors and subspaces

Review

Orthogonality and FTLA

- $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ are **orthogonal** iff
$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n = 0.$$
 - Non-zero orthogonal vectors are independent.
- If V is a subspace of \mathbb{R}^n then the *orthogonal complement* of V is

- $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ are **orthogonal** iff
$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \dots + v_n w_n = 0.$$
 - Non-zero orthogonal vectors are independent.
- If V is a subspace of \mathbb{R}^n then the *orthogonal complement* of V is

$$V^\perp = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{v} \cdot \mathbf{x} = 0, \text{ for all } \mathbf{v} \in V\}$$

- $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ are **orthogonal** iff
$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \dots + v_n w_n = 0.$$
 - Non-zero orthogonal vectors are independent.
- If V is a subspace of \mathbb{R}^n then the *orthogonal complement* of V is

$$V^\perp = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{v} \cdot \mathbf{x} = 0, \text{ for all } \mathbf{v} \in V\}$$

- If $W = V^\perp$ then $W^\perp = V$.

- $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ are **orthogonal** iff
$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \dots + v_n w_n = 0.$$
 - Non-zero orthogonal vectors are independent.
- If V is a subspace of \mathbb{R}^n then the *orthogonal complement* of V is

$$V^\perp = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{v} \cdot \mathbf{x} = 0, \text{ for all } \mathbf{v} \in V\}$$

- If $W = V^\perp$ then $W^\perp = V$.
- In other words $(V^\perp)^\perp = V$.

- $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ are **orthogonal** iff
$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \dots + v_n w_n = 0.$$
 - Non-zero orthogonal vectors are independent.
- If V is a subspace of \mathbb{R}^n then the *orthogonal complement* of V is

$$V^\perp = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{v} \cdot \mathbf{x} = 0, \text{ for all } \mathbf{v} \in V\}$$

- If $W = V^\perp$ then $W^\perp = V$.
- In other words $(V^\perp)^\perp = V$.
- $\dim(V) + \dim(V^\perp) = \dim(\mathbb{R}^n) = n$

Example

Let V be the horizontal x - y -plane in \mathbb{R}^3 and W the vertical y - z -plane.

Example

Let V be the horizontal x - y -plane in \mathbb{R}^3 and W the vertical y - z -plane.

- Is W the orthogonal complement of V ?

Example

Let V be the horizontal x - y -plane in \mathbb{R}^3 and W the vertical y - z -plane.

- Is W the orthogonal complement of V ?
- Is it true that W is orthogonal to V ?

Example

Let V be the horizontal x - y -plane in \mathbb{R}^3 and W the vertical y - z -plane.

- Is W the orthogonal complement of V ?
- Is it true that W is orthogonal to V ?
- What is the orthogonal complement of V ?

Example

Let V be the horizontal x - y -plane in \mathbb{R}^3 and W the vertical y - z -plane.

- Is W the orthogonal complement of V ?
- Is it true that W is orthogonal to V ?
- What is the orthogonal complement of V ?

Example

Given

$$\text{Nul} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} \right) =$$

Example

Given

$$\text{Nul} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} \right) =$$

Example

Given

$$\text{Nul} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} \right) = \text{Span} \left(\begin{bmatrix} 2 \\ -1 \end{bmatrix} \right),$$

get

Example

Given

$$\text{Nul} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} \right) = \text{Span} \left(\begin{bmatrix} 2 \\ -1 \end{bmatrix} \right),$$

get

Example

Given

$$\text{Nul} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} \right) = \text{Span} \left(\begin{bmatrix} 2 \\ -1 \end{bmatrix} \right),$$

get

$$\text{Col} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}^T \right) =$$

Example

Given

$$\text{Nul} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} \right) = \text{Span} \left(\begin{bmatrix} 2 \\ -1 \end{bmatrix} \right),$$

get

$$\text{Col} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}^T \right) = \text{span} \left(\begin{bmatrix} 1 \\ 2 \end{bmatrix} \right)$$

Example

Given

$$\text{Nul} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix} \right) = \text{Span} \left(\begin{bmatrix} 2 \\ -1 \end{bmatrix} \right),$$

get

$$\text{Col} \left(\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}^T \right) = \text{span} \left(\begin{bmatrix} 1 \\ 2 \end{bmatrix} \right)$$

Why?

Theorem (Fundamental Theorem of Linear Algebra)

Let A be a $m \times n$ -matrix. Then

- *$\text{Nul}(A)$ is the orthogonal complement of $\text{Col}(A^T)$ (in \mathbb{R}^n).*

Theorem (Fundamental Theorem of Linear Algebra)

Let A be a $m \times n$ -matrix. Then

- $\text{Nul}(A)$ is the orthogonal complement of $\text{Col}(A^T)$ (in \mathbb{R}^n).
Also, $\dim \text{Nul}(A) + \dim \text{Col}(A^T) = (n - r) + r = n$.

Theorem (Fundamental Theorem of Linear Algebra)

Let A be a $m \times n$ -matrix. Then

- $\text{Nul}(A)$ is the orthogonal complement of $\text{Col}(A^T)$ (in \mathbb{R}^n).
Also, $\dim \text{Nul}(A) + \dim \text{Col}(A^T) = (n - r) + r = n$.
- $\text{Col}(A)$ is the orthogonal complement of $\text{Nul}(A^T)$ (in \mathbb{R}^m).

Theorem (Fundamental Theorem of Linear Algebra)

Let A be a $m \times n$ -matrix. Then

- $\text{Nul}(A)$ is the orthogonal complement of $\text{Col}(A^T)$ (in \mathbb{R}^n).
Also, $\dim \text{Nul}(A) + \dim \text{Col}(A^T) = (n - r) + r = n$.
- $\text{Col}(A)$ is the orthogonal complement of $\text{Nul}(A^T)$ (in \mathbb{R}^m).

Why?

Theorem (Fundamental Theorem of Linear Algebra)

Let A be a $m \times n$ -matrix. Then

- $Nul(A)$ is the orthogonal complement of $Col(A^T)$ (in \mathbb{R}^n).
Also, $\dim Nul(A) + \dim Col(A^T) = (n - r) + r = n$.
- $Col(A)$ is the orthogonal complement of $Nul(A^T)$ (in \mathbb{R}^m).

Why? Suppose $\mathbf{x} \in Nul(A)$. That is,

Theorem (Fundamental Theorem of Linear Algebra)

Let A be a $m \times n$ -matrix. Then

- $Nul(A)$ is the orthogonal complement of $Col(A^T)$ (in \mathbb{R}^n).
Also, $\dim Nul(A) + \dim Col(A^T) = (n - r) + r = n$.
- $Col(A)$ is the orthogonal complement of $Nul(A^T)$ (in \mathbb{R}^m).

Why? Suppose $\mathbf{x} \in Nul(A)$. That is,

$$A\mathbf{x} = \mathbf{0}$$

What does this mean? (Think [row-column rule](#)).

Theorem (Fundamental Theorem of Linear Algebra)

Let A be a $m \times n$ -matrix. Then

- $Nul(A)$ is the orthogonal complement of $Col(A^T)$ (in \mathbb{R}^n).
Also, $\dim Nul(A) + \dim Col(A^T) = (n - r) + r = n$.
- $Col(A)$ is the orthogonal complement of $Nul(A^T)$ (in \mathbb{R}^m).

Why? Suppose $\mathbf{x} \in Nul(A)$. That is,

$$A\mathbf{x} = \mathbf{0}$$

What does this mean? (Think [row-column rule](#)).

- It means that the inner product of every row of A (transposed!) with \mathbf{x} is zero.

Theorem (Fundamental Theorem of Linear Algebra)

Let A be a $m \times n$ -matrix. Then

- $Nul(A)$ is the orthogonal complement of $Col(A^T)$ (in \mathbb{R}^n).
Also, $\dim Nul(A) + \dim Col(A^T) = (n - r) + r = n$.
- $Col(A)$ is the orthogonal complement of $Nul(A^T)$ (in \mathbb{R}^m).

Why? Suppose $\mathbf{x} \in Nul(A)$. That is,

$$A\mathbf{x} = \mathbf{0}$$

What does this mean? (Think **row-column rule**).

- It means that the inner product of every row of A (transposed!) with \mathbf{x} is zero. But that implies that \mathbf{x} is **orthogonal to the row space**.

FLTA in action

Example

Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

Example

Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

Solution

This means: Find the orthogonal complement of

Example

Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

Solution

This means: Find the orthogonal complement of $\text{Col} \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix} \right)$.

Example

Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

Solution

This means: Find the orthogonal complement of $\text{Col} \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix} \right)$.

Use the Fundamental Theorem: This is

Example

Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

Solution

This means: Find the orthogonal complement of $\text{Col} \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix} \right)$.

Use the Fundamental Theorem: This is

$$\text{Nul} \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix}^T \right) =$$

Example

Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

Solution

This means: Find the orthogonal complement of $\text{Col} \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix} \right)$.

Use the Fundamental Theorem: This is

$$\text{Nul} \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix}^T \right) = \text{Nul} \left(\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \right)$$

Compute Nul space: $Nul \left(\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \right) =$

Compute Nul space: $Nul \left(\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \right) = Nul \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} \right)$

Compute Nul space: $Nul \left(\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \right) = Nul \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} \right)$

Basis for Nul : $\left\{ \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \right\}$

Compute Nul space: $Nul \left(\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \right) = Nul \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} \right)$

Basis for Nul : $\left\{ \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \right\}$

Final answer: the set of vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ is

$$\text{span} \left\{ \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \right\}$$

Alternative solution.

Alternative solution. The FLTA is not magic! You can do this the **old-fashioned way**!

Alternative solution. The FLTA is not magic! You can do this the *old-fashioned way*!

Looking for all x so that

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot x = 0 \text{ and } \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \cdot x = 0$$

Alternative solution. The FLTA is not magic! You can do this the *old-fashioned way*!

Looking for all \mathbf{x} so that

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \mathbf{x} = 0 \text{ and } \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \cdot \mathbf{x} = 0$$

Matrix form:

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Alternative solution. The FLTA is not magic! You can do this the *old-fashioned way*!

Looking for all \mathbf{x} so that

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \mathbf{x} = 0 \text{ and } \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \cdot \mathbf{x} = 0$$

Matrix form:

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Get Null space:

$$\mathbf{x} \in \text{Nul} \left(\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \right)$$

Alternative solution. The FLTA is not magic! You can do this the **old-fashioned way**!

Looking for all \mathbf{x} so that

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \mathbf{x} = 0 \text{ and } \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \cdot \mathbf{x} = 0$$

Matrix form:

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Get Null space:

$$\mathbf{x} \in \text{Nul} \left(\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \right)$$

This is the **same null space** we obtained from the FTLA.

Example

$$\text{Let } V = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b = 2c \right\}.$$

Find a basis for the orthogonal complement of V .

Example

$$\text{Let } V = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b = 2c \right\}.$$

Find a basis for the orthogonal complement of V .

Solution

Write as Null space: $V = \text{Nul} \left(\begin{bmatrix} 1 & 1 & -2 \end{bmatrix} \right).$

Example

$$\text{Let } V = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b = 2c \right\}.$$

Find a basis for the orthogonal complement of V .

Solution

Write as Null space: $V = \text{Nul} \left(\begin{bmatrix} 1 & 1 & -2 \end{bmatrix} \right).$

By FTLA: the orthogonal complement is

Example

$$\text{Let } V = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b = 2c \right\}.$$

Find a basis for the orthogonal complement of V .

Solution

Write as Null space: $V = \text{Nul} \left(\begin{bmatrix} 1 & 1 & -2 \end{bmatrix} \right)$.

By FTLA: the orthogonal complement is $\text{Col} \left(\begin{bmatrix} 1 & 1 & -2 \end{bmatrix}^T \right)$.

Example

Let $V = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b = 2c \right\}.$

Find a basis for the orthogonal complement of V .

Solution

Write as Null space: $V = \text{Nul} \left(\begin{bmatrix} 1 & 1 & -2 \end{bmatrix} \right).$

By FTLA: the orthogonal complement is $\text{Col} \left(\begin{bmatrix} 1 & 1 & -2 \end{bmatrix}^T \right).$

Basis for the orthogonal complement: $\begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}$

Example

$$\text{Let } V = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b = 2c \right\}.$$

Find a basis for the orthogonal complement of V .

Solution

Write as Null space: $V = \text{Nul} \left(\begin{bmatrix} 1 & 1 & -2 \end{bmatrix} \right)$.

By FTLA: the orthogonal complement is $\text{Col} \left(\begin{bmatrix} 1 & 1 & -2 \end{bmatrix}^T \right)$.

Basis for the orthogonal complement: $\begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}$

Geometrically this makes sense: V is a plane with normal vector

$$\begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}$$

Alternative solution.

Notice that $a + b = 2c \iff \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = 0.$

Alternative solution.

Notice that $a + b = 2c \iff \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = 0.$

Interpret the above: V is actually defined as the orthogonal complement of

Alternative solution.

Notice that $a + b = 2c \iff \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = 0$.

Interpret the above: V is actually defined as the orthogonal complement of

$$\text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} \right\}.$$

Example

$$\text{Let } V = \left\{ \begin{bmatrix} 2a + b \\ -b \\ a + b \end{bmatrix} : a, b \in \mathbb{R} \right\}.$$

Find the orthogonal complement of V .

Example

$$\text{Let } V = \left\{ \begin{bmatrix} 2a + b \\ -b \\ a + b \end{bmatrix} : a, b \in \mathbb{R} \right\}.$$

Find the orthogonal complement of V .

Solution

Write as Column space: $V =$

Example

$$\text{Let } V = \left\{ \begin{bmatrix} 2a + b \\ -b \\ a + b \end{bmatrix} : a, b \in \mathbb{R} \right\}.$$

Find the orthogonal complement of V .

Solution

Write as Column space: $V = \text{span} \left(\begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right)$, so

Example

Let $V = \left\{ \begin{bmatrix} 2a + b \\ -b \\ a + b \end{bmatrix} : a, b \in \mathbb{R} \right\}.$

Find the orthogonal complement of V .

Solution

Write as Column space: $V = \text{span} \left(\begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right),$ so

$$V = \text{Col} \left(\begin{bmatrix} 2 & 1 \\ 0 & -1 \\ 1 & 1 \end{bmatrix} \right)$$

Example

Let $V = \left\{ \begin{bmatrix} 2a + b \\ -b \\ a + b \end{bmatrix} : a, b \in \mathbb{R} \right\}.$

Find the orthogonal complement of V .

Solution

Write as Column space: $V = \text{span} \left(\begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right),$ so

$$V = \text{Col} \left(\begin{bmatrix} 2 & 1 \\ 0 & -1 \\ 1 & 1 \end{bmatrix} \right)$$

By FTLA the orthogonal complement is

Example

Let $V = \left\{ \begin{bmatrix} 2a + b \\ -b \\ a + b \end{bmatrix} : a, b \in \mathbb{R} \right\}.$

Find the orthogonal complement of V .

Solution

Write as Column space: $V = \text{span} \left(\begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right),$ so

$$V = \text{Col} \left(\begin{bmatrix} 2 & 1 \\ 0 & -1 \\ 1 & 1 \end{bmatrix} \right)$$

By FTLA the orthogonal complement is $\text{Nul} \left(\begin{bmatrix} 2 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix} \right)$

Get RREF to compute Null space: $\begin{bmatrix} 2 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix} \rightarrow$

Get RREF to compute Null space: $\begin{bmatrix} 2 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & -1 & \frac{1}{2} \end{bmatrix}$

Get RREF to compute Null space: $\begin{bmatrix} 2 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & -1 & \frac{1}{2} \end{bmatrix}$

Basis for the Null space: $\left\{ \begin{bmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right\}$

Get RREF to compute Null space: $\begin{bmatrix} 2 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & -1 & \frac{1}{2} \end{bmatrix}$

Basis for the Null space: $\left\{ \begin{bmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right\}$

Get RREF to compute Null space: $\begin{bmatrix} 2 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & -1 & \frac{1}{2} \end{bmatrix}$

Basis for the Null space: $\left\{ \begin{bmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right\}$

So the orthogonal complement to V is: $\text{span} \left\{ \begin{bmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right\}$

Directions and Equations. Let V be a subspace of \mathbb{R}^n . Then there are *two* ways of describing V .

Directions and Equations. Let V be a subspace of \mathbb{R}^n . Then there are *two* ways of describing V .

By directions: If $V = \text{Col}(A)$ then you know that any vector \mathbf{v} in V is a linear combination of the columns of A ,

Directions and Equations. Let V be a subspace of \mathbb{R}^n . Then there are *two* ways of describing V .

By directions: If $V = \text{Col}(A)$ then you know that any vector \mathbf{v} in V is a linear combination of the columns of A , so you know in which directions \mathbf{v} can point.

Directions and Equations. Let V be a subspace of \mathbb{R}^n . Then there are *two* ways of describing V .

By directions: If $V = \text{Col}(A)$ then you know that any vector \mathbf{v} in V is a linear combination of the columns of A , so you know in which directions \mathbf{v} can point.

By equations: If $V = \text{Nul}(B)$ then you know that any \mathbf{v} in V satisfies the equations $\mathbf{R}_i^T \cdot \mathbf{v} = 0$, for all rows \mathbf{R}_i of B .

Directions and Equations. Let V be a subspace of \mathbb{R}^n . Then there are *two* ways of describing V .

By directions: If $V = \text{Col}(A)$ then you know that any vector \mathbf{v} in V is a linear combination of the columns of A , so you know in which directions \mathbf{v} can point.

By equations: If $V = \text{Nul}(B)$ then you know that any \mathbf{v} in V satisfies the equations $\mathbf{R}_i^T \cdot \mathbf{v} = 0$, for all rows \mathbf{R}_i of B .

Both descriptions are useful, and we will often switch between them, to answer any particular question we want to answer.

A new perspective on $A\mathbf{x} = \mathbf{b}$

To see if $A\mathbf{x} = \mathbf{b}$ has a solution, check that

Direct approach: $\mathbf{b} \in \text{Col}(A)$

To see if $A\mathbf{x} = \mathbf{b}$ has a solution, check that

Direct approach: $\mathbf{b} \in \text{Col}(A)$

Indirect approach: $\mathbf{b} \perp \text{Nul}(A^T)$

To see if $A\mathbf{x} = \mathbf{b}$ has a solution, check that

Direct approach: $\mathbf{b} \in \text{Col}(A)$

Indirect approach: $\mathbf{b} \perp \text{Nul}(A^T)$

The indirect approach means:

To see if $A\mathbf{x} = \mathbf{b}$ has a solution, check that

Direct approach: $\mathbf{b} \in \text{Col}(A)$

Indirect approach: $\mathbf{b} \perp \text{Nul}(A^T)$

The indirect approach means:

$$\text{if } \underbrace{\mathbf{y}^T A = \mathbf{0}}_{\mathbf{y} \in \text{Nul}(A^T)}, \text{ then } \underbrace{\mathbf{y}^T \mathbf{b} = 0}_{\mathbf{b} \perp \mathbf{y}}.$$

Example

Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix}$. For which \mathbf{b} does $A\mathbf{x} = \mathbf{b}$ have a solution?

Solution (old)

Solution (old)

Write augmented matrix, get Echelon form:

$$\left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 3 & 1 & b_2 \\ 0 & 5 & b_3 \end{array} \right] \longrightarrow$$

Solution (old)

Write augmented matrix, get Echelon form:

$$\left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 3 & 1 & b_2 \\ 0 & 5 & b_3 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 0 & -5 & -3b_1 + b_2 \\ 0 & 0 & -3b_1 + b_2 + b_3 \end{array} \right]$$

Solution (old)

Write augmented matrix, get Echelon form:

$$\left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 3 & 1 & b_2 \\ 0 & 5 & b_3 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 0 & -5 & -3b_1 + b_2 \\ 0 & 0 & -3b_1 + b_2 + b_3 \end{array} \right]$$

When is this consistent?

Solution (old)

Write augmented matrix, get Echelon form:

$$\left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 3 & 1 & b_2 \\ 0 & 5 & b_3 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 0 & -5 & -3b_1 + b_2 \\ 0 & 0 & -3b_1 + b_2 + b_3 \end{array} \right]$$

When is this consistent? Whenever $-3b_1 + b_2 + b_3 = 0$.

Solution (new)

Indirect approach says: $A\mathbf{x} = \mathbf{b}$ solvable $\iff \mathbf{b} \perp \text{Nul}(A^T)$.

Solution (new)

Indirect approach says: $A\mathbf{x} = \mathbf{b}$ solvable $\iff \mathbf{b} \perp \text{Nul}(A^T)$.

Find basis for $\text{Nul}(A^T)$:

$$\left[\begin{array}{ccc|c} 1 & 3 & 0 & 0 \\ 2 & 1 & 5 & 0 \end{array} \right] \longrightarrow$$

Solution (new)

Indirect approach says: $A\mathbf{x} = \mathbf{b}$ solvable $\iff \mathbf{b} \perp \text{Nul}(A^T)$.

Find basis for $\text{Nul}(A^T)$:

$$\left[\begin{array}{ccc|c} 1 & 3 & 0 & 0 \\ 2 & 1 & 5 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right]$$

Solution (new)

Indirect approach says: $A\mathbf{x} = \mathbf{b}$ solvable $\iff \mathbf{b} \perp \text{Nul}(A^T)$.

Find basis for $\text{Nul}(A^T)$:

$$\left[\begin{array}{ccc|c} 1 & 3 & 0 & 0 \\ 2 & 1 & 5 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right]$$

so $\text{Nul}(A^T)$ has basis $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$

Solution (new)

Indirect approach says: $A\mathbf{x} = \mathbf{b}$ solvable $\iff \mathbf{b} \perp \text{Nul}(A^T)$.

Find basis for $\text{Nul}(A^T)$:

$$\left[\begin{array}{ccc|c} 1 & 3 & 0 & 0 \\ 2 & 1 & 5 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right]$$

so $\text{Nul}(A^T)$ has basis $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$

Need $\mathbf{b} \perp \text{Nul}(A^T)$: $A\mathbf{x} = \mathbf{b}$ is solvable \iff

Solution (new)

Indirect approach says: $A\mathbf{x} = \mathbf{b}$ solvable $\iff \mathbf{b} \perp \text{Nul}(A^T)$.

Find basis for $\text{Nul}(A^T)$:

$$\left[\begin{array}{ccc|c} 1 & 3 & 0 & 0 \\ 2 & 1 & 5 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right]$$

so $\text{Nul}(A^T)$ has basis $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$

Need $\mathbf{b} \perp \text{Nul}(A^T)$: $A\mathbf{x} = \mathbf{b}$ is solvable $\iff \mathbf{b} \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix} = 0$

Solution (new)

Indirect approach says: $A\mathbf{x} = \mathbf{b}$ solvable $\iff \mathbf{b} \perp \text{Nul}(A^T)$.

Find basis for $\text{Nul}(A^T)$:

$$\left[\begin{array}{ccc|c} 1 & 3 & 0 & 0 \\ 2 & 1 & 5 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right]$$

so $\text{Nul}(A^T)$ has basis $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$

Need $\mathbf{b} \perp \text{Nul}(A^T)$: $A\mathbf{x} = \mathbf{b}$ is solvable $\iff \mathbf{b} \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix} = 0$

This is the same condition as before!

Motivation

How to find almost-solutions

Why do we care about orthogonality?

Why do we care about orthogonality? Not all linear systems have solutions.

Why do we care about orthogonality? Not all linear systems have solutions. In fact, for many applications, data needs to be fitted and there is **no hope** for a perfect match.

Why do we care about orthogonality? Not all linear systems have solutions. In fact, for many applications, data needs to be fitted and there is **no hope** for a perfect match. For example, $A\mathbf{x} = \mathbf{b}$ with

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \mathbf{x} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

has **no solution**:

Why do we care about orthogonality? Not all linear systems have solutions. In fact, for many applications, data needs to be fitted and there is **no hope** for a perfect match. For example, $A\mathbf{x} = \mathbf{b}$ with

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \mathbf{x} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

has **no solution**: $\begin{bmatrix} -1 \\ 2 \end{bmatrix}$ is not in $\text{Col}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$


Idea

Instead of giving up, we want the \mathbf{x} which makes $A\mathbf{x}$ and \mathbf{b} as close as possible.

Idea

Instead of giving up, we want the x which makes Ax and b as close as possible.


Such x is characterized by Ax being **orthogonal** to the error $b - Ax$.




Application: Directed graphs

Set up


- Graphs appear in **network analysis** (e.g. internet) or **circuit analysis**.



- Graphs appear in **network analysis** (e.g. internet) or **circuit analysis**.
- Arrow indicates direction of flow



- Graphs appear in **network analysis** (e.g. internet) or **circuit analysis**.
- Arrow indicates direction of flow
- No edges from a node to itself



Definition

Let G be a graph with m edges and n nodes.


Definition

Let G be a graph with m edges and n nodes. The **edge-node incidence matrix** of G is the $m \times n$ matrix A with

$$A_{i,j} = \begin{cases} -1, & \text{if edge } i \text{ leaves node } j \\ +1, & \text{if edge } i \text{ enters node } j \\ 0, & \text{otherwise} \end{cases}$$


Example

Give the edge-node incidence matrix of our graph.



Example

Give the edge-node incidence matrix of our graph.




Solution

$$\begin{bmatrix} -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 1 \end{bmatrix}$$

- Each column represents a node

Example

Give the edge-node incidence matrix of our graph.



Solution

$$\begin{bmatrix} -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 1 \end{bmatrix}$$

- Each column represents a node
- Each row represents an edge

We are going to use linear algebra to study networks!