

# Math 415 - Lecture 9

## Vector spaces and subspaces

Monday September 14th 2015

Textbook: Chapter 2.1.

Textbook: Chapter 2.1.

Suggested practice exercises: Chapter 2.1: 1, 2, 10, 11, 17, 18.

Textbook: Chapter 2.1.

Suggested practice exercises: Chapter 2.1: 1, 2, 10, 11, 17, 18.

Khan Academy video: Linear Subspaces

# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix.

# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix. What about  $3 \times 3, \dots, n \times n$ ?

# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix. What about  $3 \times 3, \dots, n \times n$ ? We use:

## Theorem

*An  $n \times n$  matrix  $A$  is invertible if and only if  $A$  is row equivalent to  $I_n$ ,*

# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix. What about  $3 \times 3, \dots, n \times n$ ? We use:

## Theorem

*An  $n \times n$  matrix  $A$  is invertible if and only if  $A$  is row equivalent to  $I_n$ , and in this case, any sequence of elementary row operations that reduces  $A$  to  $I_n$  will also transform  $I_n$  to  $A^{-1}$ .*



# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix. What about  $3 \times 3, \dots, n \times n$ ? We use:

## Theorem

*An  $n \times n$  matrix  $A$  is invertible if and only if  $A$  is row equivalent to  $I_n$ , and in this case, any sequence of elementary row operations that reduces  $A$  to  $I_n$  will also transform  $I_n$  to  $A^{-1}$ .*

Note that this tells us a lot about  $Ax = b$  if  $A$  is invertible.

# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix. What about  $3 \times 3, \dots, n \times n$ ? We use:

## Theorem

*An  $n \times n$  matrix  $A$  is invertible if and only if  $A$  is row equivalent to  $I_n$ , and in this case, any sequence of elementary row operations that reduces  $A$  to  $I_n$  will also transform  $I_n$  to  $A^{-1}$ .*

Note that this tells us a lot about  $Ax = b$  if  $A$  is invertible.

- $Ax = b$  has how many pivots in  $A$ ?

# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix. What about  $3 \times 3, \dots, n \times n$ ? We use:

## Theorem

*An  $n \times n$  matrix  $A$  is invertible if and only if  $A$  is row equivalent to  $I_n$ , and in this case, any sequence of elementary row operations that reduces  $A$  to  $I_n$  will also transform  $I_n$  to  $A^{-1}$ .*

Note that this tells us a lot about  $Ax = b$  if  $A$  is invertible.

- $Ax = b$  has how many pivots in  $A$ ?
- How many free variables?

# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix. What about  $3 \times 3, \dots, n \times n$ ? We use:

## Theorem

*An  $n \times n$  matrix  $A$  is invertible if and only if  $A$  is row equivalent to  $I_n$ , and in this case, any sequence of elementary row operations that reduces  $A$  to  $I_n$  will also transform  $I_n$  to  $A^{-1}$ .*

Note that this tells us a lot about  $Ax = b$  if  $A$  is invertible.

- $Ax = b$  has how many pivots in  $A$ ?
- How many free variables?
- Can  $Ax = b$  be inconsistent?

# Inverses

We know how to find the inverse of a  $2 \times 2$  matrix. What about  $3 \times 3, \dots, n \times n$ ? We use:

## Theorem

*An  $n \times n$  matrix  $A$  is invertible if and only if  $A$  is row equivalent to  $I_n$ , and in this case, any sequence of elementary row operations that reduces  $A$  to  $I_n$  will also transform  $I_n$  to  $A^{-1}$ .*

Note that this tells us a lot about  $Ax = b$  if  $A$  is invertible.

- $Ax = b$  has how many pivots in  $A$ ?
- How many free variables?
- Can  $Ax = b$  be inconsistent?

Here is the algorithm to find the inverse of a matrix  $A$ , called the  
**Gauss-Jordan Method**

- Place  $A$  and  $I$  side-by-side to form an augmented matrix  $[A \mid I]$ .

Here is the algorithm to find the inverse of a matrix  $A$ , called the **Gauss-Jordan Method**

- Place  $A$  and  $I$  side-by-side to form an augmented matrix  $[A \mid I]$ . This is an  $n \times 2n$  matrix (**Big Augmented Matrix**), instead of  $n \times (n + 1)$ , for the usual augmented matrix.

Here is the algorithm to find the inverse of a matrix  $A$ , called the **Gauss-Jordan Method**

- Place  $A$  and  $I$  side-by-side to form an augmented matrix  $[A \mid I]$ .
- Then perform row operations on this matrix (which will produce identical operations on  $A$  and  $I$ ).



Here is the algorithm to find the inverse of a matrix  $A$ , called the **Gauss-Jordan Method**

- Place  $A$  and  $I$  side-by-side to form an augmented matrix  $[A \mid I]$ .
- Then perform row operations on this matrix (which will produce identical operations on  $A$  and  $I$ ).
- So by the Theorem:

$$[A \mid I] \text{ will row reduce to } [I \mid A^{-1}]$$

or  $A$  is not invertible.

### Example

Find the inverse of  $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ , if it exists.

### Example

Find the inverse of  $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ , if it exists.

### Example

Find the inverse of  $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ , if it exists.

*Solution:*

$$\begin{aligned} [A \mid I] &= \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \sim \dots \sim \\ &\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right] \end{aligned}$$

### Example

Find the inverse of  $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ , if it exists.

*Solution:*

$$\begin{aligned} [A \mid I] &= \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \sim \dots \sim \\ &\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right] \end{aligned}$$

So

$$A^{-1} =$$

### Example

Find the inverse of  $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ , if it exists.

*Solution:*

$$\begin{aligned} [A \mid I] &= \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \sim \dots \sim \\ &\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right] \end{aligned}$$

So

$$A^{-1} = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 \\ \frac{3}{2} & 1 & 0 \end{bmatrix}$$

Example (Let's do the previous example step by step.)

$$[A \mid I] =$$

Example (Let's do the previous example step by step.)

$$[A \mid I] =$$



Example (Let's do the previous example step by step.)

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

Example (Let's do the previous example step by step.)

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R1 \rightarrow \frac{1}{2}R1} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

Example (Let's do the previous example step by step.)

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R1 \rightarrow \frac{1}{2}R1} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \rightarrow R2 + 3R1} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

Example (Let's do the previous example step by step.)

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R1 \rightarrow \frac{1}{2}R1} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \rightarrow R2 + 3R1} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \leftrightarrow R3} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right]$$

Example (Let's do the previous example step by step.)

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R1 \rightarrow \frac{1}{2}R1} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \rightarrow R2 + 3R1} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \leftrightarrow R3} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right]$$

Check at home that  $AA^{-1} = I_3$ .

## Remark

Why does this algorithm work?

## Remark

Why does this algorithm work?

- At each step, we get

$$[ A \mid I ] \rightsquigarrow [ E_1 A \mid E_1 ] \rightsquigarrow [ E_2 E_1 A \mid E_2 E_1 ] \rightsquigarrow \dots$$

## Remark

Why does this algorithm work?

- At each step, we get

$$[ A \mid I ] \rightsquigarrow [ E_1 A \mid E_1 ] \rightsquigarrow [ E_2 E_1 A \mid E_2 E_1 ] \rightsquigarrow \dots$$

- So each step is of the form

$$[ FA \mid F ], \quad F = E_r \dots E_3 E_2 E_1$$



## Remark

Why does this algorithm work?

- At each step, we get

$$[ A \mid I ] \rightsquigarrow [ E_1 A \mid E_1 ] \rightsquigarrow [ E_2 E_1 A \mid E_2 E_1 ] \rightsquigarrow \dots$$

- So each step is of the form

$$[ FA \mid F ], \quad F = E_r \dots E_3 E_2 E_1$$

- If we succeed in row reducing  $A$  to  $I$ , the final step is

$$[ FA \mid F ] = [ I \mid F ]$$

## Remark

Why does this algorithm work?

- At each step, we get

$$[ A \mid I ] \rightsquigarrow [ E_1 A \mid E_1 ] \rightsquigarrow [ E_2 E_1 A \mid E_2 E_1 ] \rightsquigarrow \dots$$

- So each step is of the form

$$[ FA \mid F ], \quad F = E_r \dots E_3 E_2 E_1$$

- If we succeed in row reducing  $A$  to  $I$ , the final step is

$$[ FA \mid F ] = [ I \mid F ]$$

- So  $FA = I$ , which means that

## Remark

Why does this algorithm work?

- At each step, we get

$$[ A \mid I ] \rightsquigarrow [ E_1 A \mid E_1 ] \rightsquigarrow [ E_2 E_1 A \mid E_2 E_1 ] \rightsquigarrow \dots$$

- So each step is of the form

$$[ FA \mid F ], \quad F = E_r \dots E_3 E_2 E_1$$

- If we succeed in row reducing  $A$  to  $I$ , the final step is

$$[ FA \mid F ] = [ I \mid F ]$$

- So  $FA = I$ , which means that

## Remark

Why does this algorithm work?

- At each step, we get

$$[ A \mid I ] \rightsquigarrow [ E_1 A \mid E_1 ] \rightsquigarrow [ E_2 E_1 A \mid E_2 E_1 ] \rightsquigarrow \dots$$

- So each step is of the form

$$[ FA \mid F ], \quad F = E_r \dots E_3 E_2 E_1$$

- If we succeed in row reducing  $A$  to  $I$ , the final step is

$$[ FA \mid F ] = [ I \mid F ]$$

- So  $FA = I$ , which means that  $A^{-1} = F$ .

# One more example

## Example

Use the Gauss Jordan method to compute the inverse of

$$\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ -1 & 0 & 1 \end{bmatrix}$$

# One more example

Solution

$$[A \mid I] =$$

# One more example

Solution

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ -1 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

# One more example

## Solution

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ -1 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R1 + R3} \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 0 & -1 & 1 & 1 & 0 & 1 \end{array} \right]$$



# One more example

## Solution

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ -1 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R1 + R3} \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 0 & -1 & 1 & 1 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R2 + R3} \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{array} \right]$$

# One more example

## Solution

$$[A \mid I] = \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ -1 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R1 + R3} \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 0 & -1 & 1 & 1 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R3 \rightarrow R2 + R3} \left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{array} \right]$$

Failure: the reduced row echelon form of  $A$  will not be  $I$ , so  $A$  has no inverse!

**Practice Problems.** Find the inverse of  $A$ :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

**Practice Problems.** Find the inverse of  $A$ :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}.$  Hint: What is  $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ ?

**Practice Problems.** Find the inverse of  $A$ :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}.$  Hint: What is  $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ ?

- $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$

**Practice Problems.** Find the inverse of  $A$ :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}.$  Hint: What is  $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ ?

- $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 2 & 3 & 0 \\ 4 & 5 & 6 & 0 \\ 6 & 7 & 8 & 0 \\ 9 & 0 & 1 & 0 \end{bmatrix}.$

**Practice Problems.** Find the inverse of  $A$ :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}.$  Hint: What is  $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ ?

- $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 2 & 3 & 0 \\ 4 & 5 & 6 & 0 \\ 6 & 7 & 8 & 0 \\ 9 & 0 & 1 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}.$

# Vector Spaces and Subspaces



# Vector Spaces and Subspaces

- The most important property of column vectors in  $\mathbb{R}^n$  is that you can take *linear combinations* of them.

# Vector Spaces and Subspaces

- The most important property of column vectors in  $\mathbb{R}^n$  is that you can take *linear combinations* of them.
- There are many mathematical objects  $X, Y, \dots$  for which a linear combination  $cX + dY$  make sense, and have the usual properties of linear combination in  $\mathbb{R}^n$

# Vector Spaces and Subspaces

- The most important property of column vectors in  $\mathbb{R}^n$  is that you can take *linear combinations* of them.
- There are many mathematical objects  $X, Y, \dots$  for which a linear combination  $cX + dY$  make sense, and have the usual properties of linear combination in  $\mathbb{R}^n$
- We are going to define a *vector space* in general as a collection of objects for which linear combinations make sense. The objects of such a set are called vectors.

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense.

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below.

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all  $u$ ,  $v$ , and  $w$  in  $V$  and for all scalars  $c$  and  $d$ .

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all  $u$ ,  $v$ , and  $w$  in  $V$  and for all scalars  $c$  and  $d$ .

1.  $\mathbf{u} + \mathbf{v}$  is in  $V$ .

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all  $u$ ,  $v$ , and  $w$  in  $V$  and for all scalars  $c$  and  $d$ .

1.  $\mathbf{u} + \mathbf{v}$  is in  $V$ . ( $V$  is “closed under addition”.)



## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all  $u$ ,  $v$ , and  $w$  in  $V$  and for all scalars  $c$  and  $d$ .

1.  $\mathbf{u} + \mathbf{v}$  is in  $V$ . ( $V$  is “closed under addition”.)
2.  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all  $u$ ,  $v$ , and  $w$  in  $V$  and for all scalars  $c$  and  $d$ .

1.  $\mathbf{u} + \mathbf{v}$  is in  $V$ . ( $V$  is “closed under addition”.)
2.  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ .

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all  $u$ ,  $v$ , and  $w$  in  $V$  and for all scalars  $c$  and  $d$ .

1.  $\mathbf{u} + \mathbf{v}$  is in  $V$ . ( $V$  is “closed under addition”.)
2.  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ .
4. There is a vector (called the zero vector)  $\mathbf{0}$  in  $V$  such that  $\mathbf{u} + \mathbf{0} = \mathbf{u}$ .

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all  $u$ ,  $v$ , and  $w$  in  $V$  and for all scalars  $c$  and  $d$ .

1.  $\mathbf{u} + \mathbf{v}$  is in  $V$ . ( $V$  is “closed under addition”.)
2.  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ .
4. There is a vector (called the zero vector)  $\mathbf{0}$  in  $V$  such that  $\mathbf{u} + \mathbf{0} = \mathbf{u}$ .
5. For each  $\mathbf{u}$  in  $V$ , there is a vector  $-\mathbf{u}$  in  $V$  satisfying  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .

## Definition

A **vector space** is a non-empty set  $V$  of objects, called *vectors*, for which linear combinations make sense. More precisely: on  $V$  there are defined two operations, called *addition* and *multiplication* by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all  $u$ ,  $v$ , and  $w$  in  $V$  and for all scalars  $c$  and  $d$ .

1.  $\mathbf{u} + \mathbf{v}$  is in  $V$ . ( $V$  is “closed under addition”.)
2.  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ .
4. There is a vector (called the zero vector)  $\mathbf{0}$  in  $V$  such that  $\mathbf{u} + \mathbf{0} = \mathbf{u}$ .
5. For each  $\mathbf{u}$  in  $V$ , there is a vector  $-\mathbf{u}$  in  $V$  satisfying  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .

### Definition Continued

6.  $c\mathbf{u}$  is in  $V$ .

### Definition Continued

6.  $c\mathbf{u}$  is in  $V$ . ( $V$  is “closed under scalar multiplication”.)

### Definition Continued

6.  $c\mathbf{u}$  is in  $V$ . ( $V$  is “closed under scalar multiplication”.)
7.  $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .



### Definition Continued

6.  $c\mathbf{u}$  is in  $V$ . ( $V$  is “closed under scalar multiplication”.)
7.  $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8.  $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ .

### Definition Continued

6.  $c\mathbf{u}$  is in  $V$ . ( $V$  is “closed under scalar multiplication”.)
7.  $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8.  $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ .
9.  $(cd)\mathbf{u} = c(d\mathbf{u})$ .

### Definition Continued

6.  $c\mathbf{u}$  is in  $V$ . ( $V$  is “closed under scalar multiplication”.)
7.  $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8.  $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ .
9.  $(cd)\mathbf{u} = c(d\mathbf{u})$ .
10.  $1\mathbf{u} = \mathbf{u}$ .

### Definition Continued

6.  $c\mathbf{u}$  is in  $V$ . ( $V$  is “closed under scalar multiplication”.)
7.  $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8.  $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ .
9.  $(cd)\mathbf{u} = c(d\mathbf{u})$ .
10.  $1\mathbf{u} = \mathbf{u}$ .

## Vector Space Examples

# Vector Space Examples

## Example

Let  $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ . This is a vector space.

# Vector Space Examples

## Example

Let  $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ . This is a vector space.

# Vector Space Examples

## Example

Let  $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ . This is a vector space.

We need to say what the two operations are.



# Vector Space Examples

## Example

Let  $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ . This is a vector space.

We need to say what the two operations are.

Addition:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} e & f \\ g & h \end{bmatrix} =$$

# Vector Space Examples

## Example

Let  $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ . This is a vector space.

We need to say what the two operations are.

Addition:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} a+e & b+f \\ c+g & d+h \end{bmatrix}.$$

# Vector Space Examples

## Example

Let  $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ . This is a vector space.

We need to say what the two operations are.

Addition:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} a+e & b+f \\ c+g & d+h \end{bmatrix}.$$

Scalar Multiplication:

$$e \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} =$$

# Vector Space Examples

## Example

Let  $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ . This is a vector space.

We need to say what the two operations are.

Addition:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} a+e & b+f \\ c+g & d+h \end{bmatrix}.$$

Scalar Multiplication:

$$e \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} ea & eb \\ ec & ed \end{bmatrix}.$$

Next we need to say what the zero vector is.

Next we need to say what the zero vector is.

Question: What is the matrix  $\mathbf{0}$  such that  $\mathbf{0} + A = A$  for any  $(2 \times 2)$  matrix  $A$ ?

Next we need to say what the zero vector is.

Question: What is the matrix  $\mathbf{0}$  such that  $\mathbf{0} + A = A$  for any  $(2 \times 2)$  matrix  $A$ ?

Answer: We see that the  $\mathbf{0}$  vector is  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ .

Next we need to say what the zero vector is.

Question: What is the matrix  $\mathbf{0}$  such that  $\mathbf{0} + A = A$  for any  $(2 \times 2)$  matrix  $A$ ?

Answer: We see that the  $\mathbf{0}$  vector is  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ .

Then we need to check all the 10 axioms. They follow from the corresponding properties of ordinary numbers.



## Remarks

## Remarks

- We can take instead of matrices of size  $2 \times 2$  matrices of any shape: you can check that the set  $M_{m \times n}$  of  $m \times n$  matrices is also a vector space, in the same way as we indicated above.

## Remarks

- We can take instead of matrices of size  $2 \times 2$  matrices of any shape: you can check that the set  $M_{m \times n}$  of  $m \times n$  matrices is also a vector space, in the same way as we indicated above.
- Confusing: in the vector space  $M_{2 \times 2}$  the vectors are in fact  $2 \times 2$  matrices!

## Remarks

- We can take instead of matrices of size  $2 \times 2$  matrices of any shape: you can check that the set  $M_{m \times n}$  of  $m \times n$  matrices is also a vector space, in the same way as we indicated above.
- Confusing: in the vector space  $M_{2 \times 2}$  the vectors are in fact  $2 \times 2$  matrices!
- In the definition of the vector space  $M_{2 \times 2}$  the multiplication of matrices plays no role; matrix multiplication will show up when we study the connections *between* vector spaces.

## Remarks

- We can take instead of matrices of size  $2 \times 2$  matrices of any shape: you can check that the set  $M_{m \times n}$  of  $m \times n$  matrices is also a vector space, in the same way as we indicated above.
- Confusing: in the vector space  $M_{2 \times 2}$  the vectors are in fact  $2 \times 2$  matrices!
- In the definition of the vector space  $M_{2 \times 2}$  the multiplication of matrices plays no role; matrix multiplication will show up when we study the connections *between* vector spaces.
- a “vector”  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$  behaves very much like a column vector  $\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$ .

## Remarks

- We can take instead of matrices of size  $2 \times 2$  matrices of any shape: you can check that the set  $M_{m \times n}$  of  $m \times n$  matrices is also a vector space, in the same way as we indicated above.
- Confusing: in the vector space  $M_{2 \times 2}$  the vectors are in fact  $2 \times 2$  matrices!
- In the definition of the vector space  $M_{2 \times 2}$  the multiplication of matrices plays no role; matrix multiplication will show up when we study the connections *between* vector spaces.
- a “vector”  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$  behaves very much like a column vector  $\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$ .

## Remarks

- We can take instead of matrices of size  $2 \times 2$  matrices of any shape: you can check that the set  $M_{m \times n}$  of  $m \times n$  matrices is also a vector space, in the same way as we indicated above.
- Confusing: in the vector space  $M_{2 \times 2}$  the vectors are in fact  $2 \times 2$  matrices!
- In the definition of the vector space  $M_{2 \times 2}$  the multiplication of matrices plays no role; matrix multiplication will show up when we study the connections *between* vector spaces.
- a “vector”  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$  behaves very much like a column vector  $\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$ . A fancy person would say that the vector spaces  $M_{2 \times 2}$  and  $\mathbb{R}^4$  are *isomorphic*.

# Vector Space Examples

## Example

Let  $n \geq 0$  be an integer and let

$\mathbf{P}_n$  = the set of all polynomials of degree at most  $n$ .



# Vector Space Examples

## Example

Let  $n \geq 0$  be an integer and let

$\mathbf{P}_n$  = the set of all polynomials of degree at most  $n$ .

This is a vector space.

Members of  $\mathbf{P}_n$  have the form

$$\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$$

where  $a_0, a_1, \dots, a_n$  are real numbers and  $t$  is a variable.

# Vector Space Examples

## Example

Let  $n \geq 0$  be an integer and let

$\mathbf{P}_n$  = the set of all polynomials of degree at most  $n$ .

This is a vector space.

Members of  $\mathbf{P}_n$  have the form

$$\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$$

where  $a_0, a_1, \dots, a_n$  are real numbers and  $t$  is a variable. We will just verify 3 out of the 10 axioms here.

# Vector Space Examples

Let  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$  and  $\mathbf{q}(t) = b_0 + b_1t + \cdots + b_nt^n$  and let  $c$  be a scalar.

# Vector Space Examples

Let  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$  and  $\mathbf{q}(t) = b_0 + b_1t + \cdots + b_nt^n$  and let  $c$  be a scalar.

The polynomial  $\mathbf{p} + \mathbf{q}$  is defined as follows:

$$(\mathbf{p} + \mathbf{q})(t) = \mathbf{p}(t) + \mathbf{q}(t).$$

# Vector Space Examples

Let  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$  and  $\mathbf{q}(t) = b_0 + b_1t + \cdots + b_nt^n$  and let  $c$  be a scalar.

The polynomial  $\mathbf{p} + \mathbf{q}$  is defined as follows:

$$(\mathbf{p} + \mathbf{q})(t) = \mathbf{p}(t) + \mathbf{q}(t).$$

Therefore,

$$\begin{aligned}(\mathbf{p} + \mathbf{q})(t) &= \mathbf{p}(t) + \mathbf{q}(t) \\ &= (a_0 + b_0) + (a_1 + b_1)t + \cdots + (a_n + b_n)t^n.\end{aligned}$$

which is also a polynomial of degree at most  $n$ .

# Vector Space Examples

Let  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$  and  $\mathbf{q}(t) = b_0 + b_1t + \cdots + b_nt^n$  and let  $c$  be a scalar.

The polynomial  $\mathbf{p} + \mathbf{q}$  is defined as follows:

$$(\mathbf{p} + \mathbf{q})(t) = \mathbf{p}(t) + \mathbf{q}(t).$$

Therefore,

$$\begin{aligned}(\mathbf{p} + \mathbf{q})(t) &= \mathbf{p}(t) + \mathbf{q}(t) \\ &= (a_0 + b_0) + (a_1 + b_1)t + \cdots + (a_n + b_n)t^n.\end{aligned}$$

which is also a polynomial of degree at most  $n$ . So  $\mathbf{p} + \mathbf{q}$  is in  $\mathbf{P}_n$  (i.e.  $\mathbf{P}_n$  is closed under addition). This verifies Axiom 1.

# Vector Space Examples

Next we need to find a zero vector.

# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?



# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?

**Answer:** Take  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$  (zero vector in  $\mathbf{P}_n$ ) Then

# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?

**Answer:** Take  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$  (zero vector in  $\mathbf{P}_n$ ) Then

$$\begin{aligned}(\mathbf{p} + \mathbf{0})(t) &= (a_0 + a_1t + \cdots + a_nt^n) + (0 + 0t + \cdots + 0t^n) \\&= (a_0 + 0) + (a_1 + 0)t + \cdots + (a_n + 0)t^n \\&= a_0 + a_1t + \cdots + a_nt^n\end{aligned}$$

and so  $\mathbf{p} + \mathbf{0} = \mathbf{p}$ . This verifies Axiom 4.

# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?

**Answer:** Take  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$  (zero vector in  $\mathbf{P}_n$ ) Then

$$\begin{aligned}(\mathbf{p} + \mathbf{0})(t) &= (a_0 + a_1t + \cdots + a_nt^n) + (0 + 0t + \cdots + 0t^n) \\&= (a_0 + 0) + (a_1 + 0)t + \cdots + (a_n + 0)t^n \\&= a_0 + a_1t + \cdots + a_nt^n\end{aligned}$$

and so  $\mathbf{p} + \mathbf{0} = \mathbf{p}$ . This verifies Axiom 4.

Next we define scalar multiplication.

# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?

**Answer:** Take  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$  (zero vector in  $\mathbf{P}_n$ ) Then

$$\begin{aligned}(\mathbf{p} + \mathbf{0})(t) &= (a_0 + a_1t + \cdots + a_nt^n) + (0 + 0t + \cdots + 0t^n) \\&= (a_0 + 0) + (a_1 + 0)t + \cdots + (a_n + 0)t^n \\&= a_0 + a_1t + \cdots + a_nt^n\end{aligned}$$

and so  $\mathbf{p} + \mathbf{0} = \mathbf{p}$ . This verifies Axiom 4.

Next we define scalar multiplication. Remember  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$ .

# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?

**Answer:** Take  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$  (zero vector in  $\mathbf{P}_n$ ) Then

$$\begin{aligned}(\mathbf{p} + \mathbf{0})(t) &= (a_0 + a_1t + \cdots + a_nt^n) + (0 + 0t + \cdots + 0t^n) \\&= (a_0 + 0) + (a_1 + 0)t + \cdots + (a_n + 0)t^n \\&= a_0 + a_1t + \cdots + a_nt^n\end{aligned}$$

and so  $\mathbf{p} + \mathbf{0} = \mathbf{p}$ . This verifies Axiom 4.

Next we define scalar multiplication. Remember  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$ . We define

$$(c\mathbf{p})(t) =$$

# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?

**Answer:** Take  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$  (zero vector in  $\mathbf{P}_n$ ) Then

$$\begin{aligned}(\mathbf{p} + \mathbf{0})(t) &= (a_0 + a_1t + \cdots + a_nt^n) + (0 + 0t + \cdots + 0t^n) \\&= (a_0 + 0) + (a_1 + 0)t + \cdots + (a_n + 0)t^n \\&= a_0 + a_1t + \cdots + a_nt^n\end{aligned}$$

and so  $\mathbf{p} + \mathbf{0} = \mathbf{p}$ . This verifies Axiom 4.

Next we define scalar multiplication. Remember  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$ . We define

$$(c\mathbf{p})(t) = c\mathbf{p}(t) =$$

# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?

**Answer:** Take  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$  (zero vector in  $\mathbf{P}_n$ ) Then

$$\begin{aligned}(\mathbf{p} + \mathbf{0})(t) &= (a_0 + a_1t + \cdots + a_nt^n) + (0 + 0t + \cdots + 0t^n) \\&= (a_0 + 0) + (a_1 + 0)t + \cdots + (a_n + 0)t^n \\&= a_0 + a_1t + \cdots + a_nt^n\end{aligned}$$

and so  $\mathbf{p} + \mathbf{0} = \mathbf{p}$ . This verifies Axiom 4.

Next we define scalar multiplication. Remember  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$ . We define

$$(c\mathbf{p})(t) = c\mathbf{p}(t) = (ca_0) + (ca_1)t + \cdots + (ca_n)t^n$$

which is in  $\mathbf{P}_n$ . so that Axiom 6 holds.

# Vector Space Examples

Next we need to find a zero vector.

**Question:** What this is the polynomial  $\mathbf{0}(t)$  such that  $\mathbf{0}(t) + p(t) = p(t)$ ?

**Answer:** Take  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$  (zero vector in  $\mathbf{P}_n$ ) Then

$$\begin{aligned}(\mathbf{p} + \mathbf{0})(t) &= (a_0 + a_1t + \cdots + a_nt^n) + (0 + 0t + \cdots + 0t^n) \\&= (a_0 + 0) + (a_1 + 0)t + \cdots + (a_n + 0)t^n \\&= a_0 + a_1t + \cdots + a_nt^n\end{aligned}$$

and so  $\mathbf{p} + \mathbf{0} = \mathbf{p}$ . This verifies Axiom 4.

Next we define scalar multiplication. Remember  $\mathbf{p}(t) = a_0 + a_1t + \cdots + a_nt^n$ . We define

$$(c\mathbf{p})(t) = c\mathbf{p}(t) = (ca_0) + (ca_1)t + \cdots + (ca_n)t^n$$

which is in  $\mathbf{P}_n$ . so that Axiom 6 holds.

The other 7 axioms also hold, so  $\mathbf{P}_n$  is a vector space.



## Subspaces

# Subspaces

New vector spaces may be formed from subsets of other vector spaces.

# Subspaces

New vector spaces may be formed from subsets of other vector spaces. These are called **subspaces**.

# Subspaces

New vector spaces may be formed from subsets of other vector spaces. These are called **subspaces**.

## Definition

A *subspace* of a vector space  $V$  is a subset  $H$  of  $V$  that satisfies 3 properties:

# Subspaces

New vector spaces may be formed from subsets of other vector spaces. These are called **subspaces**.

## Definition

A *subspace* of a vector space  $V$  is a subset  $H$  of  $V$  that satisfies 3 properties:

- The zero vector (of  $V$ ) belongs to  $H$ .

# Subspaces

New vector spaces may be formed from subsets of other vector spaces. These are called **subspaces**.

## Definition

A *subspace* of a vector space  $V$  is a subset  $H$  of  $V$  that satisfies 3 properties:

- The zero vector (of  $V$ ) belongs to  $H$ .
- If  $\mathbf{u}, \mathbf{v}$  both belong to  $H$  also the sum  $\mathbf{u} + \mathbf{v}$  belongs to  $H$ . ( $H$  is *closed* under vector addition).

# Subspaces

New vector spaces may be formed from subsets of other vector spaces. These are called **subspaces**.

## Definition

A *subspace* of a vector space  $V$  is a subset  $H$  of  $V$  that satisfies 3 properties:

- The zero vector (of  $V$ ) belongs to  $H$ .
- If  $\mathbf{u}, \mathbf{v}$  both belong to  $H$  also the sum  $\mathbf{u} + \mathbf{v}$  belongs to  $H$ . ( $H$  is *closed* under vector addition).
- If  $\mathbf{u}$  is in  $H$  and  $c$  is any scalar also  $c\mathbf{u}$  belongs to  $H$ . ( $H$  is closed under scalar multiplication.)

# Subspaces

New vector spaces may be formed from subsets of other vector spaces. These are called **subspaces**.

## Definition

A *subspace* of a vector space  $V$  is a subset  $H$  of  $V$  that satisfies 3 properties:

- The zero vector (of  $V$ ) belongs to  $H$ .
- If  $\mathbf{u}, \mathbf{v}$  both belong to  $H$  also the sum  $\mathbf{u} + \mathbf{v}$  belongs to  $H$ . ( $H$  is *closed* under vector addition).
- If  $\mathbf{u}$  is in  $H$  and  $c$  is any scalar also  $c\mathbf{u}$  belongs to  $H$ . ( $H$  is closed under scalar multiplication.)



# Subspaces

New vector spaces may be formed from subsets of other vector spaces. These are called **subspaces**.

## Definition

A *subspace* of a vector space  $V$  is a subset  $H$  of  $V$  that satisfies 3 properties:

- The zero vector (of  $V$ ) belongs to  $H$ .
- If  $\mathbf{u}, \mathbf{v}$  both belong to  $H$  also the sum  $\mathbf{u} + \mathbf{v}$  belongs to  $H$ . ( $H$  is *closed* under vector addition).
- If  $\mathbf{u}$  is in  $H$  and  $c$  is any scalar also  $c\mathbf{u}$  belongs to  $H$ . ( $H$  is closed under scalar multiplication.)

Note that if the subset  $H$  satisfies these three properties, then  $H$  itself is a vector space.

# Subspaces

## Example

$Z = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$  is a subspace of  $\mathbb{R}^2$ . Why?

# Subspaces

## Example

$Z = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$  is a subspace of  $\mathbb{R}^2$ . Why?

### Check:

- $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  is in  $Z$ .
- $\begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0+0 \\ 0+0 \end{bmatrix}$  is in  $Z$ .
- $c \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} c0 \\ c0 \end{bmatrix}$  is in  $Z$ .

# Subspaces

## Example

$Z = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$  is a subspace of  $\mathbb{R}^2$ . Why?

### Check:

- $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  is in  $Z$ .
- $\begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0+0 \\ 0+0 \end{bmatrix}$  is in  $Z$ .
- $c \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} c0 \\ c0 \end{bmatrix}$  is in  $Z$ .

$Z$  is called the zero subspace of  $\mathbb{R}^2$ .

# Subspaces

## Example

$Z = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$  is a subspace of  $\mathbb{R}^2$ . Why?

### Check:

- $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  is in  $Z$ .
- $\begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0+0 \\ 0+0 \end{bmatrix}$  is in  $Z$ .
- $c \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} c0 \\ c0 \end{bmatrix}$  is in  $Z$ .

$Z$  is called the zero subspace of  $\mathbb{R}^2$ . Every vectorspace has a zero subspace consisting just of the zero vector.

# Subspaces

## Example

$H = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$  is a subspace of  $\mathbb{R}^2$ . Why?

# Subspaces

## Example

$H = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$  is a subspace of  $\mathbb{R}^2$ . Why?

### Check:

- $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  is in  $H$ .
- $\begin{bmatrix} a \\ a \end{bmatrix} + \begin{bmatrix} b \\ b \end{bmatrix} = \begin{bmatrix} a+b \\ a+b \end{bmatrix}$  is in  $H$ .
- $c \begin{bmatrix} a \\ a \end{bmatrix} = \begin{bmatrix} ca \\ ca \end{bmatrix}$  is in  $H$ .

# Subspaces

## Example

Let  $H = \left\{ \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ . Show that  $H$  is a subspace of  $\mathbb{R}^3$ .



# Subspaces

## Example

Let  $H = \left\{ \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ . Show that  $H$  is a subspace of  $\mathbb{R}^3$ .

Verify properties 1, 2, and 3 of the definition of a subspace.

- The zero vector of  $\mathbb{R}^3$  is in  $H$ .

$$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \in H, \quad (a = b = 0)$$

# Subspaces

- Adding two vectors in  $H$  always produces another vector whose second entry is 0 and therefore the sum of two vectors in  $H$  is also in  $H$ . ( $H$  is closed under addition.)

$$\begin{bmatrix} a \\ 0 \\ b \end{bmatrix} + \begin{bmatrix} c \\ 0 \\ d \end{bmatrix} = \begin{bmatrix} a + c \\ 0 \\ b + d \end{bmatrix}.$$

# Subspaces

- Adding two vectors in  $H$  always produces another vector whose second entry is 0 and therefore the sum of two vectors in  $H$  is also in  $H$ . ( $H$  is closed under addition.)

$$\begin{bmatrix} a \\ 0 \\ b \end{bmatrix} + \begin{bmatrix} c \\ 0 \\ d \end{bmatrix} = \begin{bmatrix} a + c \\ 0 \\ b + d \end{bmatrix}.$$

- Multiplying a vector in  $H$  by a scalar produces another vector in  $H$ . ( $H$  is closed under scalar multiplication.)

$$c \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} = \begin{bmatrix} ca \\ 0 \\ cb \end{bmatrix}.$$

# Subspaces

Since those three properties hold,  $H$  is a subspace of  $\mathbb{R}^3$ .

## Remark

Vectors  $(a, 0, b)$  look and act like the points  $(a, b)$  in  $\mathbb{R}^2$ .  
But they are **not** the same!

# Subspaces

## Example

Is  $H = \left\{ \begin{bmatrix} x \\ x+1 \end{bmatrix} : x \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ?

(i.e. does  $H$  satisfy the properties of a subspace?)

# Subspaces

## Example

Is  $H = \left\{ \begin{bmatrix} x \\ x+1 \end{bmatrix} : x \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ?

(i.e. does  $H$  satisfy the properties of a subspace?)

$H$  does not contain the zero vector (property 1).

$$\begin{bmatrix} x \\ x+1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

cannot be true for any value of  $x$ .

Therefore,  $H$  is **not** a subspace!

### Example

Is  $H = \left\{ \begin{bmatrix} x \\ x+1 \end{bmatrix} : x \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ?

(i.e. does  $H$  satisfy the properties of a subspace?)

### Example

Is  $H = \left\{ \begin{bmatrix} x \\ x+1 \end{bmatrix} : x \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ?

(i.e. does  $H$  satisfy the properties of a subspace?)

Another way to show that  $H$  is not a subspace of  $\mathbb{R}^2$  is to check whether  $H$  is closed under addition (property 2).



### Example

Is  $H = \left\{ \begin{bmatrix} x \\ x+1 \end{bmatrix} : x \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ?


(i.e. does  $H$  satisfy the properties of a subspace?)

Another way to show that  $H$  is not a subspace of  $\mathbb{R}^2$  is to check whether  $H$  is closed under addition (property 2).

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \end{bmatrix} \in H$$

but

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} \notin H.$$



## Problem

*Find as many subspaces in  $\mathbb{R}^2$  as you can.*

## Problem

*Find as many subspaces in  $\mathbb{R}^2$  as you can.*


## Problem


*Find as many subspaces in  $\mathbb{R}^2$  as you can.*

Think of this at home.

## Example

Is one of the following a subspace of  $\mathbb{R}^2$ ?





## Example

Is this a subspace of  $\mathbb{R}^3$ ?

