

# Math 415 - Lecture 6

## Elementary Matrices, LU Decomposition

Friday September 4th 2015

Textbook: Chapter 1.4, 1.5

Textbook: Chapter 1.4, 1.5

Suggested Practice Exercise: Chapter 1.4 Exercise 22, 27, Chapter  
1.5: 4, 5, 11, 23, 29

**Textbook:** Chapter 1.4, 1.5

**Suggested Practice Exercise:** Chapter 1.4 Exercise 22, 27, Chapter 1.5: 4, 5, 11, 23, 29

**Khan Academy Video:** Matrix multiplication (part I), Matrix multiplication (part II), Defined and undefined matrix operations

# Review of Matrix Multiplication

- **Matrix multiplication is linear combination:**  $A\mathbf{x}$  is a linear combination of the columns of  $A$  with weights given by the entries of  $\mathbf{x}$ .

# Review of Matrix Multiplication

- **Matrix multiplication is linear combination:**  $A\mathbf{x}$  is a linear combination of the columns of  $A$  with weights given by the entries of  $\mathbf{x}$ .

## Example

$$A\mathbf{x} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & -1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} =$$

# Review of Matrix Multiplication

- **Matrix multiplication is linear combination:**  $A\mathbf{x}$  is a linear combination of the columns of  $A$  with weights given by the entries of  $\mathbf{x}$ .

## Example

$$A\mathbf{x} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & -1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 4 \end{bmatrix} + 0 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + (-1) \begin{bmatrix} 3 \\ 0 \end{bmatrix}$$

# Review of Matrix Multiplication

- **Matrix multiplication is linear combination:**  $Ax$  is a linear combination of the columns of  $A$  with weights given by the entries of  $x$ .

## Example

$$Ax = \begin{bmatrix} 1 & 2 & 3 \\ 4 & -1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 4 \end{bmatrix} + 0 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + (-1) \begin{bmatrix} 3 \\ 0 \end{bmatrix}$$

- **Linear Combination is Linear System**

## Example

$$x_1 \begin{bmatrix} 1 \\ 4 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 3 \\ 0 \end{bmatrix} = \begin{bmatrix} -2 \\ 4 \end{bmatrix} \Leftrightarrow$$



# Review of Matrix Multiplication

- **Matrix multiplication is linear combination:**  $Ax$  is a linear combination of the columns of  $A$  with weights given by the entries of  $x$ .

## Example

$$Ax = \begin{bmatrix} 1 & 2 & 3 \\ 4 & -1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 4 \end{bmatrix} + 0 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + (-1) \begin{bmatrix} 3 \\ 0 \end{bmatrix}$$

- **Linear Combination is Linear System**

## Example

$$x_1 \begin{bmatrix} 1 \\ 4 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 3 \\ 0 \end{bmatrix} = \begin{bmatrix} -2 \\ 4 \end{bmatrix} \Leftrightarrow \begin{cases} x_1 + 2x_2 + 3x_3 = -2 \\ 4x_1 + (-1)x_2 + 0x_3 = 4 \end{cases}$$

# Review of Matrix Multiplication, Cont

- $A\mathbf{x} = \mathbf{b}$  is the matrix form of the linear system with augmented matrix  $\left[ \begin{array}{c|c} A & \mathbf{b} \end{array} \right]$ .

# Review of Matrix Multiplication, Cont

- $A\mathbf{x} = \mathbf{b}$  is the matrix form of the linear system with augmented matrix  $\left[ \begin{array}{c|c} A & \mathbf{b} \end{array} \right]$ .

## Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & -1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2 \\ 4 \end{bmatrix} \leftrightarrow$$

# Review of Matrix Multiplication, Cont

- $A\mathbf{x} = \mathbf{b}$  is the matrix form of the linear system with augmented matrix  $\left[ \begin{array}{c|c} A & \mathbf{b} \end{array} \right]$ .

## Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & -1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2 \\ 4 \end{bmatrix} \leftrightarrow \begin{array}{l} x_1 + 2x_2 + 3x_3 = -2 \\ 4x_1 + (-1)x_2 + 0x_3 = 4 \end{array}$$

# Review of Matrix Multiplication, Cont

- $A\mathbf{x} = \mathbf{b}$  is the matrix form of the linear system with augmented matrix  $\left[ \begin{array}{ccc|c} A & & & \mathbf{b} \end{array} \right]$ .

## Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & -1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2 \\ 4 \end{bmatrix} \leftrightarrow \begin{array}{l} x_1 + 2x_2 + 3x_3 = -2 \\ 4x_1 + (-1)x_2 + 0x_3 = 4 \end{array}$$

$$\leftrightarrow \left[ \begin{array}{ccc|c} 1 & 2 & 3 & -2 \\ 4 & -1 & 0 & 4 \end{array} \right]$$

- Each column of  $AB$  is a linear combination of the columns of  $A$  with weights given by the corresponding column of  $B$ :

$$AB = A \begin{bmatrix} \mathbf{b}_1 & \dots & \mathbf{b}_p \end{bmatrix} =$$

- Each column of  $AB$  is a linear combination of the columns of  $A$  with weights given by the corresponding column of  $B$ :

$$AB = A \begin{bmatrix} \mathbf{b}_1 & \dots & \mathbf{b}_p \end{bmatrix} =$$

- Each column of  $AB$  is a linear combination of the columns of  $A$  with weights given by the corresponding column of  $B$ :

$$AB = A \begin{bmatrix} \mathbf{b}_1 & \dots & \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} A\mathbf{b}_1 & \dots & A\mathbf{b}_p \end{bmatrix}$$



- Each column of  $AB$  is a linear combination of the columns of  $A$  with weights given by the corresponding column of  $B$ :

$$AB = A \begin{bmatrix} \mathbf{b}_1 & \dots & \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} A\mathbf{b}_1 & \dots & A\mathbf{b}_p \end{bmatrix}$$

### Example

If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 4 \\ 0 & 2 \end{bmatrix}$ , then

$$AB = \left[ A \begin{bmatrix} 3 \\ 0 \end{bmatrix} \quad A \begin{bmatrix} 4 \\ 2 \end{bmatrix} \right] =$$

- Each column of  $AB$  is a linear combination of the columns of  $A$  with weights given by the corresponding column of  $B$ :

$$AB = A \begin{bmatrix} \mathbf{b}_1 & \dots & \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} A\mathbf{b}_1 & \dots & A\mathbf{b}_p \end{bmatrix}$$

### Example

If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 4 \\ 0 & 2 \end{bmatrix}$ , then

$$AB = \left[ A \begin{bmatrix} 3 \\ 0 \end{bmatrix} \quad A \begin{bmatrix} 4 \\ 2 \end{bmatrix} \right] =$$

- Each column of  $AB$  is a linear combination of the columns of  $A$  with weights given by the corresponding column of  $B$ :

$$AB = A \begin{bmatrix} \mathbf{b}_1 & \dots & \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} A\mathbf{b}_1 & \dots & A\mathbf{b}_p \end{bmatrix}$$

### Example

If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 4 \\ 0 & 2 \end{bmatrix}$ , then

$$AB = \left[ A \begin{bmatrix} 3 \\ 0 \end{bmatrix} \quad A \begin{bmatrix} 4 \\ 2 \end{bmatrix} \right] = \left[ 3 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 0 \begin{bmatrix} 2 \\ 1 \end{bmatrix} \quad 4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right]$$

- Each column of  $AB$  is a linear combination of the columns of  $A$  with weights given by the corresponding column of  $B$ :

$$AB = A \begin{bmatrix} \mathbf{b}_1 & \dots & \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} A\mathbf{b}_1 & \dots & A\mathbf{b}_p \end{bmatrix}$$

### Example

If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 4 \\ 0 & 2 \end{bmatrix}$ , then

$$\begin{aligned} AB &= \left[ A \begin{bmatrix} 3 \\ 0 \end{bmatrix} \quad A \begin{bmatrix} 4 \\ 2 \end{bmatrix} \right] = \left[ 3 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 0 \begin{bmatrix} 2 \\ 1 \end{bmatrix} \quad 4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right] \\ &= \begin{bmatrix} 3 & 8 \\ 6 & 10 \end{bmatrix} \end{aligned}$$

- Row-column rule: The  $ij$ -th entry of  $AB$  is  $a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$ .

- Row-column rule: The  $ij$ -th entry of  $AB$  is  $a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$ .

- Row-column rule: The  $ij$ -th entry of  $AB$  is  $a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$ .

### Example

If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 4 \\ 0 & 2 \end{bmatrix}$ , then the 22 entry of  $AB$  is

$$AB_{22} =$$

- Row-column rule: The  $ij$ -th entry of  $AB$  is  $a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$ .

### Example

If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 4 \\ 0 & 2 \end{bmatrix}$ , then the 22 entry of  $AB$  is

$$AB_{22} =$$



- Row-column rule: The  $ij$ -th entry of  $AB$  is  $a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$ .

### Example

If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 4 \\ 0 & 2 \end{bmatrix}$ , then the 22 entry of  $AB$  is

$$AB_{22} = \begin{bmatrix} 2 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ 2 \end{bmatrix} =$$

- Row-column rule: The  $ij$ -th entry of  $AB$  is  $a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$ .

### Example

If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 4 \\ 0 & 2 \end{bmatrix}$ , then the 22 entry of  $AB$  is

$$AB_{22} = \begin{bmatrix} 2 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ 2 \end{bmatrix} = 2 \times 4 + 1 \times 2 = 10$$

- Matrix multiplication is not commutative: usually,  $AB \neq BA$ .

# Powers of $A$

We write:  $A^k = A \cdots A$ ,  $k$ -times.

# Powers of $A$

We write:  $A^k = A \cdots A$ ,  $k$ -times.

For which matrices  $A$  does this make sense? If  $A$  is  $m \times n$  what can  $m$ ,  $n$  be?

# Powers of $A$

We write:  $A^k = A \cdots A$ ,  $k$ -times.

For which matrices  $A$  does this make sense? If  $A$  is  $m \times n$  what can  $m, n$  be?

## Example

$$\begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix}^3 = \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix}$$

# Powers of $A$

We write:  $A^k = A \cdots A$ ,  $k$ -times.

For which matrices  $A$  does this make sense? If  $A$  is  $m \times n$  what can  $m$ ,  $n$  be?

## Example

$$\begin{aligned} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix}^3 &= \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 9 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \end{aligned}$$

# Powers of $A$

We write:  $A^k = A \cdots A$ ,  $k$ -times.

For which matrices  $A$  does this make sense? If  $A$  is  $m \times n$  what can  $m, n$  be?

## Example

$$\begin{aligned}
 \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix}^3 &= \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \\
 &= \begin{bmatrix} 1 & 0 \\ 9 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \\
 &= \begin{bmatrix} 1 & 0 \\ 21 & 8 \end{bmatrix}
 \end{aligned}$$

# Powers of $A$

We write:  $A^k = A \cdots A$ ,  $k$ -times.

For which matrices  $A$  does this make sense? If  $A$  is  $m \times n$  what can  $m$ ,  $n$  be?

## Example

$$\begin{aligned} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix}^3 &= \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 9 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 21 & 8 \end{bmatrix} \end{aligned}$$

Calculating high powers of large matrix is hard. We will later learn a clever way to do this efficiently.



## Definition

If  $A$  is  $m \times n$ , the **transpose** of  $A$  is the  $n \times m$  matrix, denoted by  $A^T$ , whose columns are formed from the corresponding rows of  $A$ .

## Definition

If  $A$  is  $m \times n$ , the **transpose** of  $A$  is the  $n \times m$  matrix, denoted by  $A^T$ , whose columns are formed from the corresponding rows of  $A$ . In terms of matrix elements  $(A^T)_{ij} = A_{ji}$ .

## Definition

If  $A$  is  $m \times n$ , the **transpose** of  $A$  is the  $n \times m$  matrix, denoted by  $A^T$ , whose columns are formed from the corresponding rows of  $A$ . In terms of matrix elements  $(A^T)_{ij} = A_{ji}$ .

## Example

Let  $A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 6 & 7 & 8 & 9 & 8 \\ 7 & 6 & 5 & 4 & 3 \end{bmatrix}$ .

## Definition

If  $A$  is  $m \times n$ , the **transpose** of  $A$  is the  $n \times m$  matrix, denoted by  $A^T$ , whose columns are formed from the corresponding rows of  $A$ . In terms of matrix elements  $(A^T)_{ij} = A_{ji}$ .

## Example

$$\text{Let } A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 6 & 7 & 8 & 9 & 8 \\ 7 & 6 & 5 & 4 & 3 \end{bmatrix}.$$

$$\text{Then } A^T = \begin{bmatrix} 1 & 6 & 7 \\ 2 & 7 & 6 \\ 3 & 8 & 5 \\ 4 & 9 & 4 \\ 5 & 8 & 3 \end{bmatrix}$$

### Example

Let  $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix}$ . Compute  $AB$ ,  $(AB)^T$ ,  $A^T B^T$  and  $B^T A^T$ .

## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix}$$

## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 1 & 10 \end{bmatrix}$$

## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 1 & 10 \end{bmatrix}$$

$$(AB)^T =$$



## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 1 & 10 \end{bmatrix}$$

$$(AB)^T = \begin{bmatrix} 1 & 1 \\ 4 & 10 \end{bmatrix}$$

## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 1 & 10 \end{bmatrix}$$

$$(AB)^T = \begin{bmatrix} 1 & 1 \\ 4 & 10 \end{bmatrix}$$

$$A^T B^T = \begin{bmatrix} 1 & 3 \\ 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -2 \\ 2 & 1 & 4 \end{bmatrix}$$

## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 1 & 10 \end{bmatrix}$$

$$(AB)^T = \begin{bmatrix} 1 & 1 \\ 4 & 10 \end{bmatrix}$$

$$A^T B^T = \begin{bmatrix} 1 & 3 \\ 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -2 \\ 2 & 1 & 4 \end{bmatrix} = \begin{bmatrix} 7 & 3 & 10 \\ 2 & 0 & -4 \\ 2 & 1 & 4 \end{bmatrix}$$

## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 1 & 10 \end{bmatrix}$$

$$(AB)^T = \begin{bmatrix} 1 & 1 \\ 4 & 10 \end{bmatrix}$$

$$A^T B^T = \begin{bmatrix} 1 & 3 \\ 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -2 \\ 2 & 1 & 4 \end{bmatrix} = \begin{bmatrix} 7 & 3 & 10 \\ 2 & 0 & -4 \\ 2 & 1 & 4 \end{bmatrix}$$

$$B^T A^T =$$

## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 1 & 10 \end{bmatrix}$$

$$(AB)^T = \begin{bmatrix} 1 & 1 \\ 4 & 10 \end{bmatrix}$$

$$A^T B^T = \begin{bmatrix} 1 & 3 \\ 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -2 \\ 2 & 1 & 4 \end{bmatrix} = \begin{bmatrix} 7 & 3 & 10 \\ 2 & 0 & -4 \\ 2 & 1 & 4 \end{bmatrix}$$

$$B^T A^T = \begin{bmatrix} 1 & 0 & -2 \\ 2 & 1 & 4 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$$

## Solution

$$AB = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 1 & 10 \end{bmatrix}$$

$$(AB)^T = \begin{bmatrix} 1 & 1 \\ 4 & 10 \end{bmatrix}$$

$$A^T B^T = \begin{bmatrix} 1 & 3 \\ 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -2 \\ 2 & 1 & 4 \end{bmatrix} = \begin{bmatrix} 7 & 3 & 10 \\ 2 & 0 & -4 \\ 2 & 1 & 4 \end{bmatrix}$$

$$B^T A^T = \begin{bmatrix} 1 & 0 & -2 \\ 2 & 1 & 4 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 2 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 4 & 10 \end{bmatrix}$$



## Conclusion

The transpose of a product is the product of transposes **IN OPPOSITE ORDER**:

$$(AB)^T = B^T A^T$$



## Definition

$A$  is **symmetric** if  $A = A^T$ .

## Example

Which of these is symmetric?

$$\begin{bmatrix} 2 & 3 \\ 3 & 1 \end{bmatrix}^T =$$

$$\begin{bmatrix} 0 & 3 \\ 2 & 0 \end{bmatrix}^T =$$

## Theorem

*Let  $A$  and  $B$  denote matrices whose sizes are appropriate for the following sums and products.*

## Theorem

*Let  $A$  and  $B$  denote matrices whose sizes are appropriate for the following sums and products.*

(a)  $(A^T)^T = A,$

## Theorem

*Let  $A$  and  $B$  denote matrices whose sizes are appropriate for the following sums and products.*

(a)  $(A^T)^T = A,$

(b)  $(A + B)^T = A^T + B^T$

## Theorem

*Let  $A$  and  $B$  denote matrices whose sizes are appropriate for the following sums and products.*

- (a)  $(A^T)^T = A,$
- (b)  $(A + B)^T = A^T + B^T$
- (c) *For any scalar  $r$ ,  $(rA)^T = rA^T$*

## Theorem

*Let  $A$  and  $B$  denote matrices whose sizes are appropriate for the following sums and products.*

- (a)  $(A^T)^T = A,$
- (b)  $(A + B)^T = A^T + B^T$
- (c) *For any scalar  $r$ ,  $(rA)^T = rA^T$*
- (d)  $(AB)^T = B^T A^T$

### Example

Prove that  $(ABC)^T =$

### Example

Prove that  $(ABC)^T = C^T B^T A^T$ .

### Solution

By part d of the  
Theorem,  $(ABC)^T =$



### Example

Prove that  $(ABC)^T = C^T B^T A^T$ .

### Solution

By part d of the  
Theorem,  $(ABC)^T = (A(BC))^T =$

### Example

Prove that  $(ABC)^T = C^T B^T A^T$ .

### Solution

By part d of the  
Theorem,  $(ABC)^T = (A(BC))^T = (BC)^T A^T =$

### Example

Prove that  $(ABC)^T = C^T B^T A^T$ .

### Solution

By part d of the Theorem,  $(ABC)^T = (A(BC))^T = (BC)^T A^T = C^T B^T A^T$ .

## Definition

The  $n \times n$  **identity matrix**  $I_n$  has all entries 0, except on the main diagonal where the entries are 1. For example

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

### Definition

The  $n \times n$  **identity matrix**  $I_n$  has all entries 0, except on the main diagonal where the entries are 1. For example

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

### Definition

An **elementary matrix** is one that is obtained by performing a single elementary row operation on an identity matrix.

A **permutation matrix** is one that is obtained by performing row exchanges on an identity matrix.

### Example

$$\text{Let } E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix}.$$

$E_1$ ,  $E_2$ , and  $E_3$  are elementary matrices. Why? Are there any permutation matrices?

## Solution

Observe the following products and describe how these products can be obtained by elementary row operations on  $A$ .

$$E_1 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} =$$

## Solution

Observe the following products and describe how these products can be obtained by elementary row operations on  $A$ .

$$E_1 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ 2d & 2e & 2f \\ g & h & i \end{bmatrix}$$



## Solution

Observe the following products and describe how these products can be obtained by elementary row operations on  $A$ .

$$E_1 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ 2d & 2e & 2f \\ g & h & i \end{bmatrix}$$

$$E_2 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} =$$

## Solution

Observe the following products and describe how these products can be obtained by elementary row operations on  $A$ .

$$E_1 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ 2d & 2e & 2f \\ g & h & i \end{bmatrix}$$

$$E_2 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ g & h & i \\ d & e & f \end{bmatrix}$$

## Solution

Observe the following products and describe how these products can be obtained by elementary row operations on  $A$ .

$$E_1A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ 2d & 2e & 2f \\ g & h & i \end{bmatrix}$$

$$E_2A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ g & h & i \\ d & e & f \end{bmatrix}$$

$$E_3A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g+3a & h+3b & i+3c \end{bmatrix}$$

## Theorem

*If an elementary row operation is performed on an  $m \times n$ - matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$ -matrix  $E$  is created by performing the same row operations on  $I_m$ .*

## Theorem

*If an elementary row operation is performed on an  $m \times n$ - matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$ -matrix  $E$  is created by performing the same row operations on  $I_m$ .*

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} =$$

## Theorem

*If an elementary row operation is performed on an  $m \times n$ - matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$ -matrix  $E$  is created by performing the same row operations on  $I_m$ .*

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 \\ \\ \end{bmatrix}$$

## Theorem

*If an elementary row operation is performed on an  $m \times n$ - matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$ -matrix  $E$  is created by performing the same row operations on  $I_m$ .*

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

## Theorem

*If an elementary row operation is performed on an  $m \times n$ - matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$ -matrix  $E$  is created by performing the same row operations on  $I_m$ .*

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



## Theorem

*If an elementary row operation is performed on an  $m \times n$ - matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$ -matrix  $E$  is created by performing the same row operations on  $I_m$ .*

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & & \\ 0 & & \end{bmatrix}$$

## Theorem

*If an elementary row operation is performed on an  $m \times n$ - matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$ -matrix  $E$  is created by performing the same row operations on  $I_m$ .*

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

## Theorem

*If an elementary row operation is performed on an  $m \times n$ -matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$ -matrix  $E$  is created by performing the same row operations on  $I_m$ .*

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

We write this as

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix}$$

More on inverses soon.

## Remark

Elementary matrices are *invertible* because row operations are *reversible*. To determine the inverse of an elementary matrix  $E$ , determine the elementary row operation needed to transform  $E$  back into  $I$  and apply this operation to  $I$  to find the inverse.

## Remark

Elementary matrices are *invertible* because row operations are *reversible*. To determine the inverse of an elementary matrix  $E$ , determine the elementary row operation needed to transform  $E$  back into  $I$  and apply this operation to  $I$  to find the inverse.

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} =$$

## Remark

Elementary matrices are *invertible* because row operations are *reversible*. To determine the inverse of an elementary matrix  $E$ , determine the elementary row operation needed to transform  $E$  back into  $I$  and apply this operation to  $I$  to find the inverse.

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

## Remark

Elementary matrices are *invertible* because row operations are *reversible*. To determine the inverse of an elementary matrix  $E$ , determine the elementary row operation needed to transform  $E$  back into  $I$  and apply this operation to  $I$  to find the inverse.

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} =$$

## Remark

Elementary matrices are *invertible* because row operations are *reversible*. To determine the inverse of an elementary matrix  $E$ , determine the elementary row operation needed to transform  $E$  back into  $I$  and apply this operation to  $I$  to find the inverse.

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



## Remark

Elementary matrices are *invertible* because row operations are *reversible*. To determine the inverse of an elementary matrix  $E$ , determine the elementary row operation needed to transform  $E$  back into  $I$  and apply this operation to  $I$  to find the inverse.

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}^{-1} =$$

## Remark

Elementary matrices are *invertible* because row operations are *reversible*. To determine the inverse of an elementary matrix  $E$ , determine the elementary row operation needed to transform  $E$  back into  $I$  and apply this operation to  $I$  to find the inverse.

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

## Example

Compute the following products using the row or column interpretation of matrix multiplication. (Don't just use the row-column rule.)

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 2 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} =$$

## Example

Compute the following products using the row or column interpretation of matrix multiplication. (Don't just use the row-column rule.)

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 2 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 2 \\ 0 & 1 & 1 \end{bmatrix}$$

## Example

Compute the following products using the row or column interpretation of matrix multiplication. (Don't just use the row-column rule.)

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 2 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 2 \\ 0 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 2 \\ 0 & 2 & 1 \end{bmatrix} =$$

## Example

Compute the following products using the row or column interpretation of matrix multiplication. (Don't just use the row-column rule.)

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 2 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 2 \\ 0 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 2 \\ 0 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 2 \\ -2 & 1 & -1 \end{bmatrix}$$