

Preparation problems for the discussion sections on October 27th and 29th

1. Let $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \right\}$ be a subspace of \mathbb{R}^4 .

(i) Find the closest point to $\begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}$ on the subspace W .

(ii) Find the projection matrix P corresponding to the projection onto W .

(iii) Use the projection matrix P to find the projection of $\begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}$ onto the subspace W .

Solution. (i)

$$\begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}_W = \frac{\begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \frac{2}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ \frac{1}{2} \\ \frac{1}{2} \end{bmatrix}$$

(ii) We have:

$$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}_W = \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{3}{4} \\ \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \end{bmatrix},$$

$$\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}_W = \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{-1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} -\frac{1}{4} \\ \frac{3}{4} \\ \frac{1}{4} \\ \frac{1}{4} \end{bmatrix},$$

$$\begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}_W = \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \end{bmatrix},$$

$$\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}_W = \frac{\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \end{bmatrix},$$

Hence, the projection matrix is:

$$P = \begin{bmatrix} \frac{3}{4} & -\frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ -\frac{1}{4} & \frac{3}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

(iii) Using P , we have:

$$\begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}_W = \begin{bmatrix} \frac{3}{4} & -\frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ -\frac{1}{4} & \frac{3}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ \frac{1}{2} \\ \frac{1}{2} \end{bmatrix}$$

□

2. Let $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\}$ and $V = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} \right\}$ be subspaces of \mathbb{R}^3 .

(i) Find the projection matrices, P and Q , corresponding to the projections onto W and V , respectively.

(ii) Check that $PQ = QP$. Can you interpret PQ as a projection matrix?

Solution. (i)

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}_W = \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{5}{6} \\ \frac{1}{6} \\ \frac{1}{3} \end{bmatrix},$$

$$\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}_W = \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} -\frac{1}{6} \\ \frac{5}{6} \\ \frac{1}{3} \end{bmatrix},$$

$$\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}_W = \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \\ \frac{1}{3} \end{bmatrix},$$

Hence, the projection matrix corresponding to the orthogonal projection onto W is:

$$P = \begin{bmatrix} \frac{5}{6} & -\frac{1}{6} & \frac{1}{3} \\ -\frac{1}{6} & \frac{5}{6} & \frac{1}{3} \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \end{bmatrix}$$

Also,

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}_V = \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{1}{6} \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 0 \end{bmatrix},$$

$$\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}_V = \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{1}{6} \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 0 \end{bmatrix},$$

$$\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}_V = \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - \frac{1}{3} \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix},$$

Hence, the projection matrix corresponding to the orthogonal projection onto V is:

$$P = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

(ii) $PQ = QP$ is the matrix corresponding to the orthogonal projection onto $W \cap V = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$. [Note:

since $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}^T \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = 0$ if you compute orthogonal projection onto W and then onto V
the answer will be same as computing orthogonal projection onto V and then onto W]
□

3. Let $A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \\ 0 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$.

- Does \mathbf{b} belong to the column space of A ? Can you solve $A\mathbf{x} = \mathbf{b}$?
- What do you expect the projection of \mathbf{b} onto $W = \text{Col}(A)$ to be?
- Find the projection $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{Col}(A)$, and then solve $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$. (The vector $\hat{\mathbf{x}}$ is called the least square solution of $A\mathbf{x} = \mathbf{b}$.)
- Solve the equation $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$. Compare with your result of the previous part! (This equation is called the normal equation of $A\mathbf{x} = \mathbf{b}$.)
- Answer these questions for A as above but with $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ (and then $\mathbf{b} = \begin{bmatrix} 0 \\ 0 \\ 4 \end{bmatrix}$).

Solution. **a.** No, \mathbf{b} does not belong to the column space of A , because it is not a linear combination of $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$. Hence there is no solution to $A\mathbf{x} = \mathbf{b}$.

b. W is the span of $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$. It can easily be seen that W is the set of vectors in \mathbb{R}^3

whose third entry is 0. Hence $\begin{bmatrix} 4 \\ 5 \\ 0 \end{bmatrix}$ is in W . Note

$$\begin{bmatrix} 0 \\ 0 \\ 6 \end{bmatrix} = \mathbf{b} - \begin{bmatrix} 4 \\ 5 \\ 0 \end{bmatrix}$$

is orthogonal to W . Hence $\begin{bmatrix} 4 \\ 5 \\ 0 \end{bmatrix}$ should be the orthogonal projection of \mathbf{b} onto W .

c. Step 1: Find the orthogonal projection of \mathbf{b} onto W .

$$\hat{\mathbf{b}} = \frac{\begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + \frac{\begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \cdot \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 4.5 \\ 4.5 \\ 0 \end{bmatrix} + \begin{bmatrix} -.5 \\ .5 \\ 0 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ 0 \end{bmatrix}.$$

Step 2: Solve $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$.

$$\left[\begin{array}{cc|c} 1 & -1 & 4 \\ 1 & 1 & 5 \\ 0 & 0 & 0 \end{array} \right] \xrightarrow{R2 \rightarrow R2 - R1} \left[\begin{array}{cc|c} 1 & -1 & 4 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{array} \right] \xrightarrow{R1 \rightarrow R1 + .5R2} \left[\begin{array}{cc|c} 1 & 0 & 4.5 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{array} \right].$$

Hence,

$$\hat{\mathbf{x}} = \begin{bmatrix} 4.5 \\ .5 \end{bmatrix}.$$

d. We first calculate $A^T A$ and $A^T \mathbf{b}$:

$$A^T A = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix},$$

$$A^T \mathbf{b} = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 9 \\ 1 \end{bmatrix}.$$

Now we have to solve

$$\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 9 \\ 1 \end{bmatrix}.$$

It is easy to check that then

$$\hat{\mathbf{x}} = \begin{bmatrix} 4.5 \\ .5 \end{bmatrix}.$$

e. I leave the details to you, but here are the solutions. For $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$:

$$\hat{\mathbf{b}} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Note that in this case $\hat{\mathbf{b}} = \mathbf{b}$ and $\hat{\mathbf{x}}$ is a solution (not just a least square solution) of $A\mathbf{x} = \mathbf{b}$.

For $\mathbf{b} = \begin{bmatrix} 0 \\ 0 \\ 4 \end{bmatrix}$:

$$\hat{\mathbf{b}} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \hat{\mathbf{x}} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

□

4. Let $A = \begin{bmatrix} 0 & 1 \\ -2 & 2 \\ 2 & 2 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$. Find the least squares solution $\hat{\mathbf{x}}$ of $A\mathbf{x} = \mathbf{b}$.

Solution. We first calculate $A^T A$ and $A^T \mathbf{b}$:

$$A^T A = \begin{bmatrix} 0 & -2 & 2 \\ 1 & 2 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -2 & 2 \\ 2 & 2 \end{bmatrix} = \begin{bmatrix} 8 & 0 \\ 0 & 9 \end{bmatrix},$$

$$A^T \mathbf{b} = \begin{bmatrix} 0 & -2 & 2 \\ 1 & 2 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -2 \\ 3 \end{bmatrix}.$$

Now we have to solve

$$\begin{bmatrix} 8 & 0 \\ 0 & 9 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} -2 \\ 3 \end{bmatrix}.$$

It is easy to check that then

$$\hat{\mathbf{x}} = \begin{bmatrix} -\frac{1}{4} \\ \frac{1}{3} \end{bmatrix}.$$

□

5. A scientist tries to find the relation between the mysterious quantities x and y . She measures the following values:

x	1	2	3	4
y	2	5	9	17

- Suppose that y is a linear function of x of the form $a + bx$. Setup the system of equations to find the coefficients a and b .
- Find the best estimate for the coefficients a and b .
- Same question if we suppose that y is a quadratic function of the $a + bx + cx^2$.

Solution. (i) We set up the equation as follows:

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \\ 9 \\ 17 \end{bmatrix}.$$

(ii) We calculate

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}^T \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 4 & 10 \\ 10 & 30 \end{bmatrix}$$

and

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}^T \begin{bmatrix} 2 \\ 5 \\ 9 \\ 17 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ 5 \\ 9 \\ 17 \end{bmatrix} = \begin{bmatrix} 33 \\ 107 \end{bmatrix}$$

Now we solve

$$\left[\begin{array}{cc|c} 4 & 10 & 33 \\ 10 & 30 & 107 \end{array} \right] \xrightarrow{R2 \rightarrow R2 - 2.5R1} \left[\begin{array}{cc|c} 4 & 10 & 33 \\ 0 & 5 & 24.5 \end{array} \right] \xrightarrow{R1 \rightarrow R1 - 2R2} \left[\begin{array}{cc|c} 4 & 0 & -16 \\ 0 & 5 & 24.5 \end{array} \right].$$

Hence $a = -4$ and $b = 4.9$.

(iii) We set up the equation as follows:

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \\ 1 & 4 & 16 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \\ 9 \\ 17 \end{bmatrix}.$$

We calculate

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \\ 1 & 4 & 16 \end{bmatrix}^T \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \\ 1 & 4 & 16 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \\ 1 & 4 & 9 & 16 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \\ 1 & 4 & 16 \end{bmatrix} = \begin{bmatrix} 4 & 10 & 30 \\ 10 & 30 & 100 \\ 30 & 100 & 354 \end{bmatrix}$$

and

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \\ 1 & 4 & 16 \end{bmatrix}^T \begin{bmatrix} 2 \\ 5 \\ 9 \\ 17 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \\ 1 & 4 & 9 & 16 \end{bmatrix} \begin{bmatrix} 2 \\ 5 \\ 9 \\ 17 \end{bmatrix} = \begin{bmatrix} 33 \\ 107 \\ 375 \end{bmatrix}.$$

One can row reduce

$$\left[\begin{array}{ccc|c} 4 & 10 & 30 & 33 \\ 10 & 30 & 100 & 107 \\ 30 & 100 & 354 & 375 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 0 & 2.25 \\ 0 & 1 & 0 & -1.35 \\ 0 & 0 & 1 & 1.25 \end{array} \right].$$

So $a = 2.25$, $b = -1.35$ and $c = 1.25$. □

6. The system of the equations $A\mathbf{x} = \mathbf{b}$ with

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} 5 \\ 0 \\ 5 \\ 10 \end{bmatrix},$$

is not consistent.

(i) Find the least squares solution $\hat{\mathbf{x}}$ for the equation $A\mathbf{x} = \mathbf{b}$.

(ii) Determine the least squares line for the data points $(-1, 5), (0, 0), (1, 5), (2, 10)$.

Solution. (i) We first calculate $A^T A$ and $A^T \mathbf{b}$:

$$A^T A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ -1 & 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 4 & 2 \\ 2 & 6 \end{bmatrix},$$

$$A^T \mathbf{b} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ -1 & 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 5 \\ 0 \\ 5 \\ 10 \end{bmatrix} = \begin{bmatrix} 20 \\ 20 \end{bmatrix}.$$

Now we have to solve

$$\begin{bmatrix} 4 & 2 \\ 2 & 6 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 20 \\ 20 \end{bmatrix}.$$

We have

$$\left[\begin{array}{cc|c} 4 & 2 & 20 \\ 2 & 6 & 20 \end{array} \right] \xrightarrow{R2 \rightarrow R2 - 1/2 R1} \left[\begin{array}{cc|c} 4 & 2 & 20 \\ 0 & 5 & 10 \end{array} \right].$$

Hence,

$$\hat{\mathbf{x}} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}.$$

(ii) Denoting the least squares line as $y = ax + b$, we have to find $\begin{bmatrix} b \\ a \end{bmatrix}$ so that $\begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} b \\ a \end{bmatrix}$

is the closest possible value to $\begin{bmatrix} 5 \\ 0 \\ 5 \\ 10 \end{bmatrix}$. From the first part, $\begin{bmatrix} b \\ a \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$. Hence, the least squares line is $y = ax + b = 2x + 4$. □