

1. Six samples of each of four types of cereal grain grown in a certain region were analyzed to determine thiamin content, resulting in the following data ($\mu\text{g/g}$):

Wheat	5.2	4.5	6.0	6.1	6.7	5.7
Barley	6.5	8.0	6.1	7.5	5.9	5.6
Maize	5.8	4.7	6.4	4.9	6.0	5.2
Oats	8.3	6.1	7.8	7.0	5.6	7.2

	n_i	\bar{x}_i	s_i	s_i^2
Wheat	6	5.7	0.7668	0.588
Barley	6	6.6	0.9508	0.904
Maize	6	5.5	0.6693	0.448
Oats	6	7.0	1.0139	1.028

Source	SS	DF	MS	F
Between	9.24	3	3.08	4.151
Within	14.84	20	0.742	
Total	24.08	23		

$F_{0.05}(3, 20) = 3.10$
 $t_{0.025}(20) = 2.086$

A $100 \times (1 - \gamma)$ -percent confidence interval the difference $\mu_i - \mu_j$, $i \neq j$, is given by

$$\bar{Y}_i - \bar{Y}_j \pm t_{\gamma/2}(N - J \text{ d.f.}) \cdot s_{pooled} \cdot \sqrt{\frac{1}{n_i} + \frac{1}{n_j}}$$

where $s_{pooled} = \sqrt{\text{MSW}}$.

- a) Construct a 95% confidence interval for the difference between the average thiamin content for Oats and Maize.

$$(7.0 - 5.5) \pm 2.086 \cdot \sqrt{0.742} \cdot \sqrt{\frac{1}{6} + \frac{1}{6}} \quad \mathbf{1.5 \pm 1.0374}$$

Tukey's pairwise comparison:

With $100 \times (1 - \gamma)$ -percent confidence *all* pairwise differences $\mu_i - \mu_j$ are bracketed by the bounds

$$(\bar{Y}_i - \bar{Y}_j) \pm \frac{q_{\gamma, J, N-J}}{\sqrt{2}} \cdot s_{pooled} \cdot \sqrt{\frac{1}{n_i} + \frac{1}{n_j}}$$

where $s_{pooled} = \sqrt{MSW}$,

$q_{\gamma, J, N-J}$ = values from Studentized Range table.

- b) Use a 95% confidence level and Tukey's pairwise comparison procedure to compare the average thiamin content for Oats with the average thiamin content for Maize.

$$q_{0.05, 4, 20} = 3.96.$$

$$(7.0 - 5.5) \pm \frac{3.96}{\sqrt{2}} \cdot \sqrt{0.742} \cdot \sqrt{\frac{1}{6} + \frac{1}{6}} \quad \mathbf{1.5 \pm 1.3926}$$

```
qtukey(0.95,4,20)
```

```
## [1] 3.958293
```

```
fit <- lm(Thiamin ~ Grain, data = Cereal)
TukeyHSD(aov(fit))
```

```
## Tukey multiple comparisons of means
## 95% family-wise confidence level
##
## Fit: aov(formula = fit)
##
## $Grain
##          diff          lwr          upr          p adj
## Maize-Barley -1.1 -2.4919842 0.29198423 0.1541415
## Oats-Barley   0.4 -0.9919842 1.79198423 0.8516188
## Wheat-Barley -0.9 -2.2919842 0.49198423 0.2980981
## Oats-Maize     1.5  0.1080158 2.89198423 0.0318296
## Wheat-Maize    0.2 -1.1919842 1.59198423 0.9773911
## Wheat-Oats    -1.3 -2.6919842 0.09198423 0.0724920
```

TukeyHSD – Tukey's Honest Significant Difference

Contrast in the means $\mu_1, \mu_2, \dots, \mu_J$

$$c_1 \mu_1 + c_2 \mu_2 + \dots + c_J \mu_J = \sum_{j=1}^J c_j \mu_j \quad \text{where } \sum_{j=1}^J c_j = 0$$

Scheffé's multiple comparison:

With $100 \times (1 - \gamma)$ -percent confidence *all* contrasts in the J population means of

the form $\sum_{j=1}^J c_j \mu_j$ are bracketed by the bounds

$$\sum_{j=1}^J c_j \bar{Y}_j \pm \sqrt{F_{\gamma}(J-1, N-J) \cdot s_{pooled}^2 \cdot (J-1) \cdot \sum_{j=1}^J \frac{c_j^2}{n_j}}$$

where $s_{pooled} = \sqrt{MSW}$.

c) Use a 95% confidence level and Scheffé's multiple comparison procedure to compare ...

(i) the average thiamin content for Oats with the average thiamin content for Maize;

$$c_W = 0, \quad c_B = 0, \quad c_M = -1, \quad c_O = 1.$$

$$(7.0 - 5.5) \pm \sqrt{3.10} \cdot \sqrt{0.742} \cdot \sqrt{3 \cdot \left(\frac{1}{6} + \frac{1}{6}\right)} \quad \mathbf{1.5 \pm 1.51664}$$

(ii) the average thiamin content for Oats and Barley with the average thiamin content for Maize;

$$c_W = 0, \quad c_B = \frac{1}{2}, \quad c_M = -1, \quad c_O = \frac{1}{2}.$$

$$\left(\frac{6.6 + 7.0}{2} - 5.5\right) \pm \sqrt{3.10} \cdot \sqrt{0.742} \cdot \sqrt{3 \cdot \left(\frac{1}{24} + \frac{1}{6} + \frac{1}{24}\right)} \quad \mathbf{1.3 \pm 1.31345}$$

- (iii) the average thiamin content for Oats and Barley with the average thiamin content for Maize and Wheat;

$$c_W = -1/2, \quad c_B = 1/2, \quad c_M = -1/2, \quad c_O = 1/2.$$

$$\left(\frac{6.6+7.0}{2} - \frac{5.7+5.5}{2} \right) \pm \sqrt{3.10} \cdot \sqrt{0.742} \cdot \sqrt{3 \cdot \left(\frac{1}{24} + \frac{1}{24} + \frac{1}{24} + \frac{1}{24} \right)} \quad \mathbf{1.2 \pm 1.0724}$$

=====

Bonferroni method:

To make m confidence intervals with *simultaneous* confidence level of at least

$(1 - \gamma) \times 100\%$, use $(1 - \frac{\gamma}{m}) \times 100\%$ for the individual confidence level.

We have 6 pairwise differences:

$$\mu_1 - \mu_2 \quad \mu_1 - \mu_3 \quad \mu_1 - \mu_4 \quad \mu_2 - \mu_3 \quad \mu_2 - \mu_4 \quad \mu_3 - \mu_4$$

To make 6 confidence intervals with *simultaneous* confidence level of at least 95%,

$$\frac{\gamma}{m} = \frac{0.05}{6}, \quad \frac{\gamma/m}{2} = \frac{0.05}{12}.$$

```
qt(1-0.05/12,20)
```

```
## [1] 2.927119
```

$$\text{Margin of error:} \quad \pm 2.927119 \cdot \sqrt{0.742} \cdot \sqrt{\frac{1}{6} + \frac{1}{6}} \quad \mathbf{\pm 1.4557}$$

Maize-Barley	-1.1	±	1.4557
Oats-Barley	0.4	±	1.4557
Wheat-Barley	-0.9	±	1.4557
Oats-Maize	1.5	±	1.4557
Wheat-Maize	0.2	±	1.4557
Wheat-Oats	-1.3	±	1.4557