

Math 415 - Lecture 26

Orthogonal Matrices and QR Decomposition

Monday October 26th 2015

Textbook reading: Chapter 3.4

Textbook reading: Chapter 3.4

Suggested practice exercises: 3.4: 13, 16, 17, 18. 13,

Textbook reading: Chapter 3.4

Suggested practice exercises: 3.4: 13, 16, 17, 18. 13,

Khan Academy video: Gram-Schmidt Example

Textbook reading: Chapter 3.4

Suggested practice exercises: 3.4: 13, 16, 17, 18. 13,

Khan Academy video: Gram-Schmidt Example

Strang lecture: Orthogonal Matrices and Gram-Schmidt Process.

Review

- Vectors $\mathbf{q}_1, \dots, \mathbf{q}_n$ are orthonormal if

$$\mathbf{q}_i^T \mathbf{q}_j = \begin{cases} 0, & \text{if } i \neq j, \\ 1, & \text{if } i = j. \end{cases}$$

- Vectors $\mathbf{q}_1, \dots, \mathbf{q}_n$ are orthonormal if

$$\mathbf{q}_i^T \mathbf{q}_j = \begin{cases} 0, & \text{if } i \neq j, \\ 1, & \text{if } i = j. \end{cases}$$

- **Gram-Schmidt** orthonormalization:
input: basis $\mathbf{a}_1, \dots, \mathbf{a}_n$ for V .

- Vectors $\mathbf{q}_1, \dots, \mathbf{q}_n$ are orthonormal if

$$\mathbf{q}_i^T \mathbf{q}_j = \begin{cases} 0, & \text{if } i \neq j, \\ 1, & \text{if } i = j. \end{cases}$$

- **Gram-Schmidt** orthonormalization:
input: basis $\mathbf{a}_1, \dots, \mathbf{a}_n$ for V .

- Vectors $\mathbf{q}_1, \dots, \mathbf{q}_n$ are orthonormal if

$$\mathbf{q}_i^T \mathbf{q}_j = \begin{cases} 0, & \text{if } i \neq j, \\ 1, & \text{if } i = j. \end{cases}$$

- **Gram-Schmidt** orthonormalization:

input: basis $\mathbf{a}_1, \dots, \mathbf{a}_n$ for V .

output: orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$ for V .

$$\mathbf{b}_1 = \mathbf{a}_1, \quad \mathbf{q}_1 = \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}$$

- Vectors $\mathbf{q}_1, \dots, \mathbf{q}_n$ are orthonormal if

$$\mathbf{q}_i^T \mathbf{q}_j = \begin{cases} 0, & \text{if } i \neq j, \\ 1, & \text{if } i = j. \end{cases}$$

- **Gram-Schmidt** orthonormalization:

input: basis $\mathbf{a}_1, \dots, \mathbf{a}_n$ for V .

output: orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$ for V .

$$\begin{aligned} \mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \end{aligned}$$

- Vectors $\mathbf{q}_1, \dots, \mathbf{q}_n$ are orthonormal if


$$\mathbf{q}_i^T \mathbf{q}_j = \begin{cases} 0, & \text{if } i \neq j, \\ 1, & \text{if } i = j. \end{cases}$$

- **Gram-Schmidt** orthonormalization:

input: basis $\mathbf{a}_1, \dots, \mathbf{a}_n$ for V .

output: orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$ for V .

$$\begin{aligned} \mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ & \dots & \dots \end{aligned}$$



Fact

if A is any matrix $A^T A$ is the matrix of dot products of the columns of A :

Fact

if A is any matrix $A^T A$ is the matrix of dot products of the columns of A :

Fact

if A is any matrix $A^T A$ is the matrix of dot products of the columns of A :

Write $A = [\mathbf{a}_1, \dots, \mathbf{a}_n]$ then

Fact

if A is any matrix $A^T A$ is the matrix of dot products of the columns of A :

Write $A = [\mathbf{a}_1, \dots, \mathbf{a}_n]$ then

$$A^T A = \begin{bmatrix} \mathbf{a}_1 \cdot \mathbf{a}_1 & \mathbf{a}_1 \cdot \mathbf{a}_2 & \mathbf{a}_1 \cdot \mathbf{a}_3 & \dots \\ \mathbf{a}_2 \cdot \mathbf{a}_1 & \mathbf{a}_2 \cdot \mathbf{a}_2 & \mathbf{a}_2 \cdot \mathbf{a}_3 & \dots \\ \mathbf{a}_3 \cdot \mathbf{a}_1 & \mathbf{a}_3 \cdot \mathbf{a}_2 & \mathbf{a}_3 \cdot \mathbf{a}_3 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Theorem

The columns of Q are orthonormal $\iff Q^T Q = I$

Theorem

The columns of Q are orthonormal $\iff Q^T Q = I$

Definition

An **orthogonal matrix** is a square matrix Q with orthonormal columns.

The QR decomposition

In linear algebra “everything” is a matrix factorization.

In linear algebra “everything” is a matrix factorization.

- Gaussian elimination in terms of matrices: $A = LU$

In linear algebra “everything” is a matrix factorization.

- Gaussian elimination in terms of matrices: $A = LU$
- Gram-Schmidt in terms of matrices $A = QR$

In linear algebra “everything” is a matrix factorization.

- Gaussian elimination in terms of matrices: $A = LU$
- Gram-Schmidt in terms of matrices $A = QR$

Theorem (QR decomposition)

Let A be an $m \times n$ matrix of rank n . There is is a orthogonal matrix $m \times n$ -matrix Q and an upper triangular $n \times n$ invertible matrix R such that

$$A = QR.$$

In linear algebra “everything” is a matrix factorization.

- Gaussian elimination in terms of matrices: $A = LU$
- Gram-Schmidt in terms of matrices $A = QR$

Theorem (QR decomposition)

Let A be an $m \times n$ matrix of rank n . There is is a orthogonal matrix $m \times n$ -matrix Q and an upper triangular $n \times n$ invertible matrix R such that

$$A = QR.$$

Idea. Gram-Schmidt on the columns of A to get columns of Q .

Recipe

In general, to obtain $A = QR$:

Recipe

In general, to obtain $A = QR$:

- Gram-Schmidt on (columns of) A , to get (columns of) Q .

Recipe

In general, to obtain $A = QR$:

- Gram-Schmidt on (columns of) A , to get (columns of) Q .
- Then $R = Q^T A$.

Recipe

In general, to obtain $A = QR$:

- Gram-Schmidt on (columns of) A , to get (columns of) Q .
- Then $R = Q^T A$.

Recipe

In general, to obtain $A = QR$:

- Gram-Schmidt on (columns of) A , to get (columns of) Q .
- Then $R = Q^T A$.

The resulting R is indeed upper triangular, and we get:

$$\begin{bmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{bmatrix} = \begin{bmatrix} | & | & \cdots \\ \mathbf{q}_1 & \mathbf{q}_2 & \cdots \\ | & | & \end{bmatrix} \begin{bmatrix} \mathbf{q}_1^T \mathbf{a}_1 & \mathbf{q}_1^T \mathbf{a}_2 & \mathbf{q}_1^T \mathbf{a}_3 & \cdots \\ & \mathbf{q}_2^T \mathbf{a}_2 & \mathbf{q}_2^T \mathbf{a}_3 & \\ & & \mathbf{q}_3^T \mathbf{a}_3 & \\ & & & \ddots \end{bmatrix}$$

Recipe

In general, to obtain $A = QR$:

- Gram-Schmidt on (columns of) A , to get (columns of) Q .
- Then $R = Q^T A$.

The resulting R is indeed upper triangular, and we get:

$$\begin{bmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{bmatrix} = \begin{bmatrix} | & | & \cdots \\ \mathbf{q}_1 & \mathbf{q}_2 & \cdots \\ | & | & \end{bmatrix} \begin{bmatrix} \mathbf{q}_1^T \mathbf{a}_1 & \mathbf{q}_1^T \mathbf{a}_2 & \mathbf{q}_1^T \mathbf{a}_3 & \cdots \\ & \mathbf{q}_2^T \mathbf{a}_2 & \mathbf{q}_2^T \mathbf{a}_3 & \\ & & \mathbf{q}_3^T \mathbf{a}_3 & \\ & & & \ddots \end{bmatrix}$$

It should be noted that, no extra work is needed for computing R :

Recipe

In general, to obtain $A = QR$:

- Gram-Schmidt on (columns of) A , to get (columns of) Q .
- Then $R = Q^T A$.

The resulting R is indeed upper triangular, and we get:

$$\begin{bmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{bmatrix} = \begin{bmatrix} | & | & \cdots \\ \mathbf{q}_1 & \mathbf{q}_2 & \cdots \\ | & | & \end{bmatrix} \begin{bmatrix} \mathbf{q}_1^T \mathbf{a}_1 & \mathbf{q}_1^T \mathbf{a}_2 & \mathbf{q}_1^T \mathbf{a}_3 & \cdots \\ & \mathbf{q}_2^T \mathbf{a}_2 & \mathbf{q}_2^T \mathbf{a}_3 & \\ & & \mathbf{q}_3^T \mathbf{a}_3 & \\ & & & \ddots \end{bmatrix}$$

It should be noted that, no extra work is needed for computing R : all the inner products in R have been computed during Gram-Schmidt.

Recipe

In general, to obtain $A = QR$:

- Gram-Schmidt on (columns of) A , to get (columns of) Q .
- Then $R = Q^T A$.

The resulting R is indeed upper triangular, and we get:

$$\begin{bmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{bmatrix} = \begin{bmatrix} | & | & \cdots \\ \mathbf{q}_1 & \mathbf{q}_2 & \cdots \\ | & | & \end{bmatrix} \begin{bmatrix} \mathbf{q}_1^T \mathbf{a}_1 & \mathbf{q}_1^T \mathbf{a}_2 & \mathbf{q}_1^T \mathbf{a}_3 & \cdots \\ & \mathbf{q}_2^T \mathbf{a}_2 & \mathbf{q}_2^T \mathbf{a}_3 & \\ & & \mathbf{q}_3^T \mathbf{a}_3 & \\ & & & \ddots \end{bmatrix}$$

It should be noted that, no extra work is needed for computing R : all the inner products in R have been computed during Gram-Schmidt.

(Just the LU decomposition encodes the steps of Gaussian elimination, the QR decomposition encodes the steps of Gram-Schmidt.)

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix}$.

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A :

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A :

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \mathbf{q}_1$$

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A :

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \mathbf{q}_1$$

$$\begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix} - \langle \begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix}, \mathbf{q}_1 \rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}, \quad \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \mathbf{q}_2$$

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A :

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \mathbf{q}_1$$

$$\begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix} - \langle \begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix}, \mathbf{q}_1 \rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}, \quad \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \mathbf{q}_2$$

$$\begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} - \langle \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \mathbf{q}_2 \rangle \mathbf{q}_2 = \begin{bmatrix} 0 \\ 5 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \mathbf{q}_3$$

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Note Q is a permutation matrix and so orthogonal.

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Note Q is a permutation matrix and so orthogonal. Why?

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Note Q is a permutation

matrix and so orthogonal. Why? Q has orthonormal columns so $Q^T Q = I$!

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Note Q is a permutation

matrix and so orthogonal. Why? Q has orthonormal columns so $Q^T Q = I$! To find R in $A = QR$, note that $Q^T A = Q^T QR = R$.

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Note Q is a permutation

matrix and so orthogonal. Why? Q has orthonormal columns so $Q^T Q = I$! To find R in $A = QR$, note that $Q^T A = Q^T QR = R$.

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Note Q is a permutation

matrix and so orthogonal. Why? Q has orthonormal columns so $Q^T Q = I$! To find R in $A = QR$, note that $Q^T A = Q^T QR = R$.

$$R = Q^T A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 5 \end{bmatrix}$$

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Note Q is a permutation

matrix and so orthogonal. Why? Q has orthonormal columns so $Q^T Q = I$! To find R in $A = QR$, note that $Q^T A = Q^T QR = R$.

$$R = Q^T A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 5 \end{bmatrix}$$

Note R is upper triangular.

Solution (continued)

Hence: $Q = [\mathbf{q}_1 \ \mathbf{q}_2 \ \mathbf{q}_3] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Note Q is a permutation

matrix and so orthogonal. Why? Q has orthonormal columns so $Q^T Q = I$! To find R in $A = QR$, note that $Q^T A = Q^T QR = R$.

$$R = Q^T A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 5 \end{bmatrix}$$

Note R is upper triangular.

Summarizing, we have

$$\begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 5 \end{bmatrix}$$

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 & 2 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$.

Applications of $A = QR$

Using QR to solve systems of equations

QR decomposition can be used to solve systems of linear equations.

$$A\mathbf{x} = \mathbf{b}$$

QR decomposition can be used to solve systems of linear equations.

$$A\mathbf{x} = \mathbf{b} \iff QR\mathbf{x} = \mathbf{b}$$

QR decomposition can be used to solve systems of linear equations.

$$\begin{aligned} A\mathbf{x} = \mathbf{b} &\iff QR\mathbf{x} = \mathbf{b} \\ &\iff R\mathbf{x} = Q^T\mathbf{b} \end{aligned}$$

QR decomposition can be used to solve systems of linear equations.

$$\begin{aligned} A\mathbf{x} = \mathbf{b} &\iff QR\mathbf{x} = \mathbf{b} \\ &\iff R\mathbf{x} = Q^T\mathbf{b} \end{aligned}$$

$R\mathbf{x} = Q^T\mathbf{b}$ is triangular, so solve it by back substitution.

QR decomposition can be used to solve systems of linear equations.

$$\begin{aligned} A\mathbf{x} = \mathbf{b} &\iff QR\mathbf{x} = \mathbf{b} \\ &\iff R\mathbf{x} = Q^T\mathbf{b} \end{aligned}$$

$R\mathbf{x} = Q^T\mathbf{b}$ is triangular, so solve it by back substitution.

QR is a little slower than LU, but makes up in numerical stability.

QR decomposition can be used to solve systems of linear equations.

$$\begin{aligned} A\mathbf{x} = \mathbf{b} &\iff QR\mathbf{x} = \mathbf{b} \\ &\iff R\mathbf{x} = Q^T\mathbf{b} \end{aligned}$$

$R\mathbf{x} = Q^T\mathbf{b}$ is triangular, so solve it by back substitution.

QR is a little slower than LU, but makes up in numerical stability.

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Proof.

$$A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$$

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Proof.

$$A^T A \hat{\mathbf{x}} = A^T \mathbf{b} \iff \underbrace{(QR)^T QR}_{=R^T Q^T QR} \hat{\mathbf{x}} = (QR)^T \mathbf{b}$$

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Proof.

$$\begin{aligned} A^T A \hat{\mathbf{x}} &= A^T \mathbf{b} \iff \underbrace{(QR)^T QR}_{=R^T Q^T QR} \hat{\mathbf{x}} = (QR)^T \mathbf{b} \\ &\iff R^T R \hat{\mathbf{x}} = R^T Q^T \mathbf{b} \end{aligned}$$

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Proof.

$$\begin{aligned} A^T A \hat{\mathbf{x}} &= A^T \mathbf{b} \iff \underbrace{(QR)^T QR}_{=R^T Q^T QR} \hat{\mathbf{x}} = (QR)^T \mathbf{b} \\ &\iff R^T R \hat{\mathbf{x}} = R^T Q^T \mathbf{b} \\ &\iff R \hat{\mathbf{x}} = Q^T \mathbf{b} \end{aligned}$$

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Proof.

$$\begin{aligned} A^T A \hat{\mathbf{x}} &= A^T \mathbf{b} && \iff \underbrace{(QR)^T QR}_{=R^T Q^T QR} \hat{\mathbf{x}} = (QR)^T \mathbf{b} \\ &&& \iff R^T R \hat{\mathbf{x}} = R^T Q^T \mathbf{b} \\ &&& \iff R \hat{\mathbf{x}} = Q^T \mathbf{b} \end{aligned}$$

Again, this is triangular, solved by back substitution.

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Proof.

$$\begin{aligned} A^T A \hat{\mathbf{x}} &= A^T \mathbf{b} \iff \underbrace{(QR)^T QR}_{=R^T Q^T QR} \hat{\mathbf{x}} = (QR)^T \mathbf{b} \\ &\iff R^T R \hat{\mathbf{x}} = R^T Q^T \mathbf{b} \\ &\iff R \hat{\mathbf{x}} = Q^T \mathbf{b} \end{aligned}$$

Again, this is triangular, solved by back substitution.

$\hat{\mathbf{x}}$ is a least square solution of $A\mathbf{x} = \mathbf{b}$

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Proof.

$$\begin{aligned}
 A^T A \hat{\mathbf{x}} &= A^T \mathbf{b} \iff \underbrace{(QR)^T QR}_{=R^T Q^T QR} \hat{\mathbf{x}} = (QR)^T \mathbf{b} \\
 &\iff R^T R \hat{\mathbf{x}} = R^T Q^T \mathbf{b} \\
 &\iff R \hat{\mathbf{x}} = Q^T \mathbf{b}
 \end{aligned}$$

Again, this is triangular, solved by back substitution.

$\hat{\mathbf{x}}$ is a least square solution of $A\mathbf{x} = \mathbf{b}$

$$\iff R\hat{\mathbf{x}} = Q^T \mathbf{b} \text{ (where } A = QR\text{)}$$

Theorem

Let A be matrix with linear independent columns. Suppose $A\mathbf{x} = \mathbf{b}$ has no solution.

Then the solution of $R\mathbf{x} = Q^T\mathbf{b}$ is the least square solution of $A\mathbf{x} = \mathbf{b}$.

Proof.

$$\begin{aligned} A^T A \hat{\mathbf{x}} &= A^T \mathbf{b} \iff \underbrace{(QR)^T QR}_{=R^T Q^T QR} \hat{\mathbf{x}} = (QR)^T \mathbf{b} \\ &\iff R^T R \hat{\mathbf{x}} = R^T Q^T \mathbf{b} \\ &\iff R \hat{\mathbf{x}} = Q^T \mathbf{b} \end{aligned}$$

Again, this is triangular, solved by back substitution.

$\hat{\mathbf{x}}$ is a least square solution of $A\mathbf{x} = \mathbf{b}$

$$\iff R \hat{\mathbf{x}} = Q^T \mathbf{b} \text{ (where } A = QR)$$

Example

Let $A = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 2 & 1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$. Find the least square solution of $A\mathbf{x} = \mathbf{b}$ using QR-decomposition.

Solution

Let us first apply Gram-Schmidt to the columns of A .

Example

Let $A = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 2 & 1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$. Find the least square solution of $A\mathbf{x} = \mathbf{b}$ using QR-decomposition.

Solution

Let us first apply Gram-Schmidt to the columns of A . (We are going to work first with unnormalized vectors, and normalize at the end. Check that this also works!)

Example

Let $A = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 2 & 1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$. Find the least square solution of $A\mathbf{x} = \mathbf{b}$ using QR-decomposition.

Solution

Let us first apply Gram-Schmidt to the columns of A . (We are going to work first with unnormalized vectors, and normalize at the

end. Check that this also works!) We have $\mathbf{b}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$ and

$$\mathbf{b}_2 = \mathbf{a}_2 - \frac{\langle \mathbf{a}_2, \mathbf{b}_1 \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} \mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - \frac{9}{9} \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}.$$

Solution (continued)

Normalizing we get

$$Q = \begin{bmatrix} 1/3 & 0 \\ 2/3 & 1/\sqrt{2} \\ 2/3 & -1/\sqrt{2} \end{bmatrix}.$$

We have $A = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 2 & 1 \end{bmatrix}$, and $Q = \begin{bmatrix} 1/3 & 0 \\ 2/3 & 1/\sqrt{2} \\ 2/3 & -1/\sqrt{2} \end{bmatrix}$.

Solution (continued)

Normalizing we get

$$Q = \begin{bmatrix} 1/3 & 0 \\ 2/3 & 1/\sqrt{2} \\ 2/3 & -1/\sqrt{2} \end{bmatrix}.$$

We have $A = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 2 & 1 \end{bmatrix}$, and $Q = \begin{bmatrix} 1/3 & 0 \\ 2/3 & 1/\sqrt{2} \\ 2/3 & -1/\sqrt{2} \end{bmatrix}$. Then

$$R = \begin{bmatrix} \mathbf{q}_1 \cdot \mathbf{a}_1 & \mathbf{q}_1 \cdot \mathbf{a}_2 \\ 0 & \mathbf{q}_2 \cdot \mathbf{a}_2 \end{bmatrix} =$$

Solution (continued)

Normalizing we get

$$Q = \begin{bmatrix} 1/3 & 0 \\ 2/3 & 1/\sqrt{2} \\ 2/3 & -1/\sqrt{2} \end{bmatrix}.$$

We have $A = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 2 & 1 \end{bmatrix}$, and $Q = \begin{bmatrix} 1/3 & 0 \\ 2/3 & 1/\sqrt{2} \\ 2/3 & -1/\sqrt{2} \end{bmatrix}$. Then

$$R = \begin{bmatrix} \mathbf{q}_1 \cdot \mathbf{a}_1 & \mathbf{q}_1 \cdot \mathbf{a}_2 \\ 0 & \mathbf{q}_2 \cdot \mathbf{a}_2 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 0 & \sqrt{2} \end{bmatrix}.$$

Solution (continued)

Normalizing we get

$$Q = \begin{bmatrix} 1/3 & 0 \\ 2/3 & 1/\sqrt{2} \\ 2/3 & -1/\sqrt{2} \end{bmatrix}.$$

We have $A = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 2 & 1 \end{bmatrix}$, and $Q = \begin{bmatrix} 1/3 & 0 \\ 2/3 & 1/\sqrt{2} \\ 2/3 & -1/\sqrt{2} \end{bmatrix}$. Then

$$R = \begin{bmatrix} \mathbf{q}_1 \cdot \mathbf{a}_1 & \mathbf{q}_1 \cdot \mathbf{a}_2 \\ 0 & \mathbf{q}_2 \cdot \mathbf{a}_2 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 0 & \sqrt{2} \end{bmatrix}.$$

Now $A\mathbf{x} = \mathbf{b}$ is not consistent.

Solution

So we do least squares, but in this case ($A = QR$) we know the normal equations are

$$R\hat{\mathbf{x}} = Q^T \mathbf{b},$$

Solution

So we do least squares, but in this case ($A = QR$) we know the normal equations are

$$R\hat{\mathbf{x}} = Q^T \mathbf{b},$$

Solution

So we do least squares, but in this case ($A = QR$) we know the normal equations are

$$R\hat{\mathbf{x}} = Q^T \mathbf{b}, \quad \begin{bmatrix} 3 & 3 \\ 0 & \sqrt{2} \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 1/3 & 2/3 & 2/3 \\ 0 & 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} =$$

Solution

So we do least squares, but in this case ($A = QR$) we know the normal equations are

$$R\hat{\mathbf{x}} = Q^T \mathbf{b}, \quad \begin{bmatrix} 3 & 3 \\ 0 & \sqrt{2} \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 1/3 & 2/3 & 2/3 \\ 0 & 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1/3 \\ 0 \end{bmatrix}$$

Solution

So we do least squares, but in this case ($A = QR$) we know the normal equations are

$$R\hat{\mathbf{x}} = Q^T \mathbf{b}, \quad \begin{bmatrix} 3 & 3 \\ 0 & \sqrt{2} \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 1/3 & 2/3 & 2/3 \\ 0 & 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1/3 \\ 0 \end{bmatrix}$$

$$\text{So } \hat{\mathbf{x}} = \begin{bmatrix} 1/9 \\ 0 \end{bmatrix}, \text{ and } \hat{\mathbf{b}} = A\hat{\mathbf{x}} = 1/9 \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}.$$