

Math 415 - Lecture 37

Singular Value Decomposition

Friday December 4th 2015

Textbook reading: Chapter 6.3

Textbook reading: Chapter 6.3

Suggested practice exercises: Chapter 6.3, # 1, 2, 3, 5, 8, 9, 15

Textbook reading: Chapter 6.3

Suggested practice exercises: Chapter 6.3, # 1, 2, 3, 5, 8, 9, 15

Strang lecture: Lecture 29: Singular Value Decomposition

Review

- Spectral theorem: If A is an $n \times n$ symmetric matrix, then it has an orthonormal basis of eigenvectors $\mathbf{v}_1 \dots \mathbf{v}_n$, and all eigenvalues are real.

- Spectral theorem: If A is an $n \times n$ symmetric matrix, then it has an orthonormal basis of eigenvectors $\mathbf{v}_1 \dots \mathbf{v}_n$, and all eigenvalues are real.
- We can write

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{v}_1 & \dots & \mathbf{v}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}} \underbrace{\begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ & & \ddots \\ 0 & 0 & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}$$

- Spectral theorem: If A is an $n \times n$ symmetric matrix, then it has an orthonormal basis of eigenvectors $\mathbf{v}_1 \dots \mathbf{v}_n$, and all eigenvalues are real.
- We can write

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{v}_1 & \dots & \mathbf{v}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}} \underbrace{\begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ & & \ddots \\ 0 & 0 & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}$$

- Today: There is a similar decomposition for any $m \times n$ matrix A .

- Spectral theorem: If A is an $n \times n$ symmetric matrix, then it has an orthonormal basis of eigenvectors $\mathbf{v}_1 \dots \mathbf{v}_n$, and all eigenvalues are real.
- We can write

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{v}_1 & \dots & \mathbf{v}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}} \underbrace{\begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ & & \ddots \\ 0 & 0 & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}$$

- Today: There is a similar decomposition for any $m \times n$ matrix A .
 - Doesn't even have to be square!

- Spectral theorem: If A is an $n \times n$ symmetric matrix, then it has an orthonormal basis of eigenvectors $\mathbf{v}_1 \dots \mathbf{v}_n$, and all eigenvalues are real.
- We can write

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{v}_1 & \dots & \mathbf{v}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}} \underbrace{\begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ & & \ddots \\ 0 & 0 & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}$$

- Today: There is a similar decomposition for any $m \times n$ matrix A .
 - Doesn't even have to be square!
 - The price we pay: different bases on the left and right sides.

Singular Value Decomposition

Goals

Starting with an $m \times n$ matrix A we want to

- Describe the *geometry* of the corresponding map $\mathbb{R}^n \rightarrow \mathbb{R}^m$,

Starting with an $m \times n$ matrix A we want to

- Describe the *geometry* of the corresponding map $\mathbb{R}^n \rightarrow \mathbb{R}^m$,
- Find a way to *approximate* A by simpler matrices, that are easier/cheaper to calculate with.

Starting with an $m \times n$ matrix A we want to

- Describe the *geometry* of the corresponding map $\mathbb{R}^n \rightarrow \mathbb{R}^m$,
- Find a way to *approximate* A by simpler matrices, that are easier/cheaper to calculate with.

Starting with an $m \times n$ matrix A we want to

- Describe the *geometry* of the corresponding map $\mathbb{R}^n \rightarrow \mathbb{R}^m$,
- Find a way to *approximate* A by simpler matrices, that are easier/cheaper to calculate with.

How?

Starting with an $m \times n$ matrix A we want to

- Describe the *geometry* of the corresponding map $\mathbb{R}^n \rightarrow \mathbb{R}^m$,
- Find a way to *approximate* A by simpler matrices, that are easier/cheaper to calculate with.

How?

Remember: for each A we get 4 subspaces

- Input space \mathbb{R}^n contains row space $\text{Col}(A^T)$ and Null space $\text{Nul}(A)$. Dimensions are r and $n - r$.

Starting with an $m \times n$ matrix A we want to

- Describe the *geometry* of the corresponding map $\mathbb{R}^n \rightarrow \mathbb{R}^m$,
- Find a way to *approximate* A by simpler matrices, that are easier/cheaper to calculate with.

How?

Remember: for each A we get 4 subspaces

- Input space \mathbb{R}^n contains row space $\text{Col}(A^T)$ and Null space $\text{Nul}(A)$. Dimensions are r and $n - r$.
- Output space \mathbb{R}^m contains columns space $\text{Col}(A)$ and left null space $\text{Nul}(A^T)$. Dimensions are r and $m - r$.

Idea

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$.

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$. Then think of what happens when we apply A to each of the basis vectors:

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$. Then think of what happens when we apply A to each of the basis vectors:

- $A\mathbf{v}_{r+1} = 0 = \dots = A\mathbf{v}_n$, since each vector belongs to $\text{Nul}(A)$.

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$. Then think of what happens when we apply A to each of the basis vectors:

- $A\mathbf{v}_{r+1} = 0 = \dots = A\mathbf{v}_n$, since each vector belongs to $\text{Nul}(A)$.

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$. Then think of what happens when we apply A to each of the basis vectors:

- $A\mathbf{v}_{r+1} = 0 = \dots = A\mathbf{v}_n$, since each vector belongs to $\text{Nul}(A)$.
- The other vectors $A\mathbf{v}_i$, $i = 1, 2, \dots, r$, will all be nonzero, in fact will give a basis of $\text{Col}(A)$!

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$. Then think of what happens when we apply A to each of the basis vectors:

- $A\mathbf{v}_{r+1} = 0 = \dots = A\mathbf{v}_n$, since each vector belongs to $\text{Nul}(A)$.
- The other vectors $A\mathbf{v}_i$, $i = 1, 2, \dots, r$, will all be nonzero, in fact will give a basis of $\text{Col}(A)$!
- Rescale these basis vectors to get unit vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r$.

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$. Then think of what happens when we apply A to each of the basis vectors:

- $A\mathbf{v}_{r+1} = 0 = \dots = A\mathbf{v}_n$, since each vector belongs to $\text{Nul}(A)$.
- The other vectors $A\mathbf{v}_i$, $i = 1, 2, \dots, r$, will all be nonzero, in fact will give a basis of $\text{Col}(A)$!
- Rescale these basis vectors to get unit vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r$.

Idea:

Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .

Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$. Then think of what happens when we apply A to each of the basis vectors:


- $A\mathbf{v}_{r+1} = 0 = \dots = A\mathbf{v}_n$, since each vector belongs to $\text{Nul}(A)$.
- The other vectors $A\mathbf{v}_i$, $i = 1, 2, \dots, r$, will all be nonzero, in fact will give a basis of $\text{Col}(A)$!
- Rescale these basis vectors to get unit vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r$. By a miracle they turn out to be orthogonal, if we choose the $\mathbf{v}_1, \mathbf{v}_2, \dots$ in the right way.
- We get $A\mathbf{v}_i = \sigma_i \mathbf{u}_i$ for $i = 1, 2, \dots, r$. The stretch factors $\sigma_i > 0$, $i = 1, 2, \dots, r$ are called the *Singular Values* of A


Idea:


Find an orthonormal basis in each of the input space subspaces, and watch what happens to these basis vectors if we multiply by A .


Choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the row space $\text{Col}(A^T)$, and a basis $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ of the null space $\text{Nul}(A)$. Then think of what happens when we apply A to each of the basis vectors:


- $A\mathbf{v}_{r+1} = 0 = \dots = A\mathbf{v}_n$, since each vector belongs to $\text{Nul}(A)$.
- The other vectors $A\mathbf{v}_i$, $i = 1, 2, \dots, r$, will all be nonzero, in fact will give a basis of $\text{Col}(A)$!
- Rescale these basis vectors to get unit vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r$. By a miracle they turn out to be orthogonal, if we choose the $\mathbf{v}_1, \mathbf{v}_2, \dots$ in the right way.
- We get $A\mathbf{v}_i = \sigma_i \mathbf{u}_i$ for $i = 1, 2, \dots, r$. The stretch factors $\sigma_i > 0$, $i = 1, 2, \dots, r$ are called the *Singular Values* of A
- Extend the \mathbf{u}_i basis of $\text{Col}(A)$ to a basis $\mathbf{u}_{r+1}, \dots, \mathbf{u}_m$ of the output space.




















What is SVD?

What is SVD?

Motto

In Linear Algebra everything is a matrix factorization.

What is SVD?

Motto

In Linear Algebra everything is a matrix factorization.

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_{\Sigma} \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_{\Sigma} \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m .

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_{\Sigma} \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m .

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m . U is $m \times m$

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_{\Sigma} \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m . U is $m \times m$
- Rows of V are an orthonormal basis for \mathbb{R}^n .

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_{\Sigma} \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m . U is $m \times m$
- Rows of V are an orthonormal basis for \mathbb{R}^n .

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m . U is $m \times m$
- Rows of V are an orthonormal basis for \mathbb{R}^n . V is $n \times n$

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m . U is $m \times m$
- Rows of V are an orthonormal basis for \mathbb{R}^n . V is $n \times n$
- Σ is rectangular $m \times n$ and diagonal,

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m . U is $m \times m$
- Rows of V are an orthonormal basis for \mathbb{R}^n . V is $n \times n$
- Σ is rectangular $m \times n$ and diagonal,

Motto

In Linear Algebra everything is a matrix factorization.

The complicated story with orthonormal basis and singular values for A gives a factorization, called **Singular Value Decomposition**:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_{\Sigma} \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

- $A = U\Sigma V^T$
- U, V are orthogonal
- Columns of U are an orthonormal basis for \mathbb{R}^m . U is $m \times m$
- Rows of V are an orthonormal basis for \mathbb{R}^n . V is $n \times n$
- Σ is rectangular $m \times n$ and diagonal, the r non zero diagonal entries are called **singular values**, they are positive

How to Compute SVD

Here is a recipe for computing SVD:

Here is a recipe for computing SVD:

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

How to Compute SVD

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

Make V : • Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$.

How to Compute SVD

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

Make V : • Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$.

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

Make V :

- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$.

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V ,

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V ,

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and
 $\sigma_{r+1} = \dots = \sigma_n = 0$.

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and
 $\sigma_{r+1} = \dots = \sigma_n = 0$.

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and $\sigma_{r+1} = \dots = \sigma_n = 0$. Put these into diagonal of **rectangular** $m \times n$ matrix Σ .

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and $\sigma_{r+1} = \dots = \sigma_n = 0$. Put these into diagonal of **rectangular** $m \times n$ matrix Σ .

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and
 $\sigma_{r+1} = \dots = \sigma_n = 0$. Put these into diagonal of
rectangular $m \times n$ matrix Σ .

Make U : • Set $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1, \dots, \mathbf{u}_r = \frac{1}{\sigma_r} A \mathbf{v}_r$.

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and
 $\sigma_{r+1} = \dots = \sigma_n = 0$. Put these into diagonal of
rectangular $m \times n$ matrix Σ .

- Make U :
- Set $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1, \dots, \mathbf{u}_r = \frac{1}{\sigma_r} A \mathbf{v}_r$.
 - Extend $\mathbf{u}_1, \dots, \mathbf{u}_r$ to an orthonormal basis
 $\mathbf{u}_1, \dots, \mathbf{u}_m$ for \mathbb{R}^m .

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and $\sigma_{r+1} = \dots = \sigma_n = 0$. Put these into diagonal of **rectangular** $m \times n$ matrix Σ .

- Make U :
- Set $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1, \dots, \mathbf{u}_r = \frac{1}{\sigma_r} A \mathbf{v}_r$.
 - Extend $\mathbf{u}_1, \dots, \mathbf{u}_r$ to an orthonormal basis $\mathbf{u}_1, \dots, \mathbf{u}_m$ for \mathbb{R}^m .
 - Put $\mathbf{u}_1, \dots, \mathbf{u}_m$ into matrix U .

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues
 $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and $\sigma_{r+1} = \dots = \sigma_n = 0$. Put these into diagonal of **rectangular** $m \times n$ matrix Σ .

- Make U :
- Set $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1, \dots, \mathbf{u}_r = \frac{1}{\sigma_r} A \mathbf{v}_r$.
 - Extend $\mathbf{u}_1, \dots, \mathbf{u}_r$ to an orthonormal basis $\mathbf{u}_1, \dots, \mathbf{u}_m$ for \mathbb{R}^m .
 - Put $\mathbf{u}_1, \dots, \mathbf{u}_m$ into matrix U .

Here is a recipe for computing SVD:

Compute $A^T A$. This is a symmetric matrix!! (Why?)

- Make V :
- Find orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$. (Why can we do this?)
 - Of $\mathbf{v}_1, \dots, \mathbf{v}_n$, make the first ones, $\mathbf{v}_1, \dots, \mathbf{v}_r$, be ones with **nonzero** eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$.
 - **Magic:** $\lambda_1, \dots, \lambda_r$ always positive!
 - Put $\mathbf{v}_1, \dots, \mathbf{v}_n$ into matrix V , those with non-zero eigenvalues come first!

Make Σ : Set $\sigma_i = \sqrt{\lambda_i}$ for $i = 1 \dots r$ and $\sigma_{r+1} = \dots = \sigma_n = 0$. Put these into diagonal of **rectangular** $m \times n$ matrix Σ .

- Make U :
- Set $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1, \dots, \mathbf{u}_r = \frac{1}{\sigma_r} A \mathbf{v}_r$.
 - Extend $\mathbf{u}_1, \dots, \mathbf{u}_r$ to an orthonormal basis $\mathbf{u}_1, \dots, \mathbf{u}_m$ for \mathbb{R}^m .
 - Put $\mathbf{u}_1, \dots, \mathbf{u}_m$ into matrix U .

Now you have $A = U \Sigma V^T$!

Example

Compute the SVD of

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

Solution

Example

Compute the SVD of

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

Solution

Compute $A^T A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.

Make V : Basis of eigenvectors for $A^T A$: $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and

$$\mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Example

Compute the SVD of

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

Solution

Compute $A^T A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.

Make V : Basis of eigenvectors for $A^T A$: $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and

$$\mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$V = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Solution (continued)

Make Σ : Eigenvalues are 1 and 1.

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} =$

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} = 1$.

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} = 1$.

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 =$

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} = 1$.

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} = 1$.

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$ and $\mathbf{u}_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$.

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} = 1$.

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$ and $\mathbf{u}_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$.

$$U = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

We get

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} = 1$.

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$ and $\mathbf{u}_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$.

$$U = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

We get

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Note: $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ has no real eigenvalues.

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} = 1$.

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$ and $\mathbf{u}_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$.

$$U = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

We get

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Note: $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ has no real eigenvalues. It's **not diagonalizable** with real matrices!

Solution (continued)

Make Σ : Eigenvalues are 1 and 1. So, $\sigma_1 = \sigma_2 = \sqrt{1} = 1$.

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$ and $\mathbf{u}_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$.

$$U = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

We get

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Note: $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ has no real eigenvalues. It's **not diagonalizable** with real matrices! But, it **has an SVD!** Wolfram Alpha

Example

Compute the SVD of

$$A = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$$

Solution

Example

Compute the SVD of

$$A = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$$

Solution

Compute $A^T A = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix}$.

Make V : Basis of eigenvectors for $A^T A$:

$$\mathbf{v}_1 = \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{bmatrix}$$

with eigenvalues $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$.

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$.

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$.

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1$

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$.

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix}$

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$.

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$.

Make U : $\mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$

$$\mathbf{u}_2 = \frac{1}{\sigma_2} A \mathbf{v}_2$$

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$.

$$\text{Make } U: \mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$
$$\mathbf{u}_2 = \frac{1}{\sigma_2} A \mathbf{v}_2 = \frac{1}{1} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$.

$$\text{Make } U: \mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$\mathbf{u}_2 = \frac{1}{\sigma_2} A \mathbf{v}_2 = \frac{1}{1} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

Solution (continued)

Make Σ : Eigenvalues were $\lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$, so
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$.

$$\text{Make } U: \mathbf{u}_1 = \frac{1}{\sigma_1} A \mathbf{v}_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$\mathbf{u}_2 = \frac{1}{\sigma_2} A \mathbf{v}_2 = \frac{1}{1} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

Final result:

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}$$

Notice how A behaves in the basis $\{\mathbf{u}_1, \mathbf{u}_2\}$ and $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$:

Notice how A behaves in the basis $\{\mathbf{u}_1, \mathbf{u}_2\}$ and $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$:

$$A\mathbf{v}_1 = \sigma_1\mathbf{u}_1:$$

Notice how A behaves in the basis $\{\mathbf{u}_1, \mathbf{u}_2\}$ and $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$:

$$A\mathbf{v}_1 = \sigma_1\mathbf{u}_1:$$

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \sqrt{3} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

Notice how A behaves in the basis $\{\mathbf{u}_1, \mathbf{u}_2\}$ and $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$:

$$A\mathbf{v}_1 = \sigma_1\mathbf{u}_1:$$

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \sqrt{3} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$A\mathbf{v}_2 = \sigma_2\mathbf{u}_2:$$

Notice how A behaves in the basis $\{\mathbf{u}_1, \mathbf{u}_2\}$ and $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$:

$$A\mathbf{v}_1 = \sigma_1\mathbf{u}_1:$$

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \sqrt{3} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$A\mathbf{v}_2 = \sigma_2\mathbf{u}_2:$$

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \end{bmatrix} = 1 \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

Notice how A behaves in the basis $\{\mathbf{u}_1, \mathbf{u}_2\}$ and $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$:

$$A\mathbf{v}_1 = \sigma_1\mathbf{u}_1:$$

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} \\ -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} = \sqrt{3} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$A\mathbf{v}_2 = \sigma_2\mathbf{u}_2:$$

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \end{bmatrix} = 1 \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$A\mathbf{v}_3 = 0.$$

A matrix might not be diagonalizable:

- If A is rectangular, it does not even have eigenvalues.

A matrix might not be diagonalizable:

- If A is rectangular, it does not even have eigenvalues.

A matrix might not be diagonalizable:

- If A is rectangular, it does not even have eigenvalues.

But A will always have an SVD!

A matrix might not be diagonalizable:

- If A is rectangular, it does not even have eigenvalues.

But A will always have an SVD! This comes at a cost:

- The SVD is not unique.

A matrix might not be diagonalizable:

- If A is rectangular, it does not even have eigenvalues.

But A will always have an SVD! This comes at a cost:

- The SVD is not unique.
- The singular values σ_i are not eigenvalues.

A matrix might not be diagonalizable:

- If A is rectangular, it does not even have eigenvalues.

But A will always have an SVD! This comes at a cost:

- The SVD is not unique.
- The singular values σ_i are not eigenvalues.

A matrix might not be diagonalizable:

- If A is rectangular, it does not even have eigenvalues.

But A will always have an SVD! This comes at a cost:

- The SVD is not unique.
- The singular values σ_i are not eigenvalues.

Note the difference: for $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ the eigenvalues are $\lambda = i, -i$ but the singular values are $\sigma = 1, 1$.

Approximation

- * To calculate matrix product AB we can use the **ROW** times **COLUMN** method: the ij component is the product $R_i B_j$, where R_i is row i of A and B_j is the j th column of B .

- * To calculate matrix product AB we can use the **ROW** times **COLUMN** method: the ij component is the product $R_i B_j$, where R_i is row i of A and B_j is the j th column of B .
- * What about **COLUMN** times **ROW**?

- * To calculate matrix product AB we can use the **ROW** times **COLUMN** method: the ij component is the product $R_i B_j$, where R_i is row i of A and B_j is the j th column of B .
- * What about **COLUMN** times **ROW**?

- * To calculate matrix product AB we can use the **ROW** times **COLUMN** method: the ij component is the product $R_i B_j$, where R_i is row i of A and B_j is the j th column of B .
- * What about **COLUMN** times **ROW**?
- *

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} a+3b & 2a+4b \\ c+3d & 2c+4d \end{bmatrix} = \\ = \begin{bmatrix} a \\ c \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix} + \begin{bmatrix} b \\ d \end{bmatrix} \begin{bmatrix} 3 & 4 \end{bmatrix}$$

- * To calculate matrix product AB we can use the **ROW** times **COLUMN** method: the ij component is the product $R_i B_j$, where R_i is row i of A and B_j is the j th column of B .
- * What about **COLUMN** times **ROW**?

*

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} a+3b & 2a+4b \\ c+3d & 2c+4d \end{bmatrix} = \\ = \begin{bmatrix} a \\ c \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix} + \begin{bmatrix} b \\ d \end{bmatrix} \begin{bmatrix} 3 & 4 \end{bmatrix}$$

- * This works for any matrix multiplication: AB is a sum of **COLUMN** times **ROW** matrices.

It turns out we can write A as a sum:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

$$= \sigma_1 \mathbf{u}_1 \mathbf{v}_1^T + \sigma_2 \mathbf{u}_2 \mathbf{v}_2^T + \dots + \sigma_r \mathbf{u}_r \mathbf{v}_r^T$$

It turns out we can write A as a sum:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

$$= \sigma_1 \mathbf{u}_1 \mathbf{v}_1^T + \sigma_2 \mathbf{u}_2 \mathbf{v}_2^T + \dots + \sigma_r \mathbf{u}_r \mathbf{v}_r^T$$

(Sanity check: An $m \times 1$ column vector times a $1 \times n$ row vector is an $m \times n$ matrix.)

Idea

We can get a good approximation to A by taking the entries of the sum with the largest singular values!

It turns out we can write A as a sum:

$$A = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{u}_1 & \dots & \mathbf{u}_m \\ | & & | \end{bmatrix}}_U \underbrace{\begin{bmatrix} \sigma_1 & 0 & \\ 0 & \sigma_2 & \\ & & \ddots \end{bmatrix}}_\Sigma \underbrace{\begin{bmatrix} - & \mathbf{v}_1^T & - \\ & \vdots & \\ - & \mathbf{v}_n^T & - \end{bmatrix}}_{V^T}$$

$$= \sigma_1 \mathbf{u}_1 \mathbf{v}_1^T + \sigma_2 \mathbf{u}_2 \mathbf{v}_2^T + \dots + \sigma_r \mathbf{u}_r \mathbf{v}_r^T$$

(Sanity check: An $m \times 1$ column vector times a $1 \times n$ row vector is an $m \times n$ matrix.)

Idea

We can get a good approximation to A by taking the entries of the sum with the largest singular values! We'll see this when we talk about image compression later.

Example

If \mathbf{u}, \mathbf{v} are non-zero, then the matrix $\mathbf{u}\mathbf{v}^T$ has rank 1. Why?

Example

Use

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}$$

to write $\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$ as a sum of rank 1 matrices.

Solution

SVD and the Four Fundamental Subspaces

The SVD of A gives orthonormal bases for all four fundamental subspaces of A .

Given $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ and $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$,

- $\text{Col}(A^T) = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$

The SVD of A gives orthonormal bases for all four fundamental subspaces of A .

Given $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ and $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$,

- $\text{Col}(A^T) = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$
- $\text{Nul}(A) = \text{Span}\{\mathbf{v}_{r+1}, \dots, \mathbf{v}_n\}$

The SVD of A gives orthonormal bases for all four fundamental subspaces of A .

Given $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ and $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$,

- $\text{Col}(A^T) = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$
- $\text{Nul}(A) = \text{Span}\{\mathbf{v}_{r+1}, \dots, \mathbf{v}_n\}$
- $\text{Col}(A) = \text{Span}\{\mathbf{u}_1, \dots, \mathbf{u}_r\}$

The SVD of A gives orthonormal bases for all four fundamental subspaces of A .

Given $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ and $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$,

- $\text{Col}(A^T) = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$
- $\text{Nul}(A) = \text{Span}\{\mathbf{v}_{r+1}, \dots, \mathbf{v}_n\}$
- $\text{Col}(A) = \text{Span}\{\mathbf{u}_1, \dots, \mathbf{u}_r\}$
- $\text{Nul}(A^T) = \text{Span}\{\mathbf{u}_{r+1}, \dots, \mathbf{u}_m\}$

Practice Questions

Example

Suppose A is an invertible square matrix. Find a singular value decomposition of A^{-1} .

Example

If A is a square matrix, then $|\det(A)|$ is the product of the singular values of A . Why?

Example

Find the singular value decomposition of $A = \begin{bmatrix} 3 & 2 & 2 \\ 2 & 3 & -2 \end{bmatrix}$.