

MATH 415 – Lecture 37

Review for Exam 3

Thursday 30 July 2015

Big topics for midterm 3

Big topics for midterm 3

- Orthogonal projection

Big topics for midterm 3

- Orthogonal projection
- Least Squares

Big topics for midterm 3

- Orthogonal projection
- Least Squares
- Gram-Schmidt

Big topics for midterm 3

- Orthogonal projection
- Least Squares
- Gram-Schmidt
- Determinants

Big topics for midterm 3

- Orthogonal projection
- Least Squares
- Gram-Schmidt
- Determinants
- Eigenvalues and eigenvectors

Big topics for midterm 3

- Orthogonal projection
- Least Squares
- Gram-Schmidt
- Determinants
- Eigenvalues and eigenvectors
- Diagonalization

Orthogonal Projection

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j =$$

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j =$$

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- Suppose that V is a subspace of W and \mathbf{x} is in W , then the **orthogonal projection** of \mathbf{x} onto V is given by

$$\hat{\mathbf{x}} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j =$$

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- Suppose that V is a subspace of W and \mathbf{x} is in W , then the **orthogonal projection** of \mathbf{x} onto V is given by

$$\hat{\mathbf{x}} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j =$$

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- Suppose that V is a subspace of W and \mathbf{x} is in W , then the **orthogonal projection** of \mathbf{x} onto V is given by

$$\hat{\mathbf{x}} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- Suppose that V is a subspace of W and \mathbf{x} is in W , then the **orthogonal projection** of \mathbf{x} onto V is given by

$$\hat{\mathbf{x}} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- The basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ has to be orthogonal for this formula!!

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- Suppose that V is a subspace of W and \mathbf{x} is in W , then the **orthogonal projection** of \mathbf{x} onto V is given by

$$\hat{\mathbf{x}} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- The basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ has to be orthogonal for this formula!!
- This decomposes $\mathbf{x} = \underbrace{\hat{\mathbf{x}}}_{\text{in } V} + \underbrace{\mathbf{x}^\perp}_{\text{in } V^\perp}$, where

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- Suppose that V is a subspace of W and \mathbf{x} is in W , then the **orthogonal projection** of \mathbf{x} onto V is given by

$$\hat{\mathbf{x}} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- The basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ has to be orthogonal for this formula!!
- This decomposes $\mathbf{x} = \underbrace{\hat{\mathbf{x}}}_{\text{in } V} + \underbrace{\mathbf{x}^\perp}_{\text{in } V^\perp}$, where

- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- Suppose that V is a subspace of W and \mathbf{x} is in W , then the **orthogonal projection** of \mathbf{x} onto V is given by

$$\hat{\mathbf{x}} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- The basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ has to be orthogonal for this formula!!
- This decomposes $\mathbf{x} = \underbrace{\hat{\mathbf{x}}}_{\text{in } V} + \underbrace{\mathbf{x}^\perp}_{\text{in } V^\perp}$, where the error \mathbf{x}^\perp is orthogonal to V .


- If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **orthogonal basis** for V , and \mathbf{x} is in V , then

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- Suppose that V is a subspace of W and \mathbf{x} is in W , then the **orthogonal projection** of \mathbf{x} onto V is given by

$$\hat{\mathbf{x}} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n \quad \text{with } c_j = \frac{\langle \mathbf{x}, \mathbf{v}_j \rangle}{\langle \mathbf{v}_j, \mathbf{v}_j \rangle}.$$

- The basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ has to be orthogonal for this formula!!
- This decomposes $\mathbf{x} = \underbrace{\hat{\mathbf{x}}}_{\text{in } V} + \underbrace{\mathbf{x}^\perp}_{\text{in } V^\perp}$, where the error \mathbf{x}^\perp is orthogonal to V . (This decomposition is unique.)



Example

What is the orthogonal projection of $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ onto $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Example

What is the orthogonal projection of $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ onto $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Solution

The projection is $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$.

Example

What is the orthogonal projection of $\begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}$ onto

$$\text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\}?$$

Example

What is the orthogonal projection of $\begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}$ onto

$$\text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\}?$$

Solution (First try:)

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Example

What is the orthogonal projection of $\begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}$ onto

$$\text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\}?$$

Solution (First try:)

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Example

What is the orthogonal projection of $\begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}$ onto

$$\text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\}?$$

Solution (First try:)

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Is the projection $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$?

Example

What is the orthogonal projection of $\begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}$ onto

$$\text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\}?$$

Solution (First try:)

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Is the projection $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$? No!

Example

What is the orthogonal projection of $\begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}$ onto

$$\text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\} ?$$

Solution (First try:)

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Is the projection $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$? No! Wrong approach!!

Example

What is the orthogonal projection of $\begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}$ onto

$$\text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\}?$$

Solution (First try:)

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Is the projection $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$? No! Wrong approach!! (This is because the basis is not orthogonal.)

Solution (Corrected:)

Form orthogonal basis first:

Solution (Corrected:)

Form orthogonal basis first:

Solution (Corrected:)

Form orthogonal basis first:

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Solution (Corrected:)

Form orthogonal basis first: $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ (for instance, using Gram-Schmidt)

Solution (Corrected:)

Form orthogonal basis first: $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ (for

instance, using Gram-Schmidt)

Then compute:

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} +$$

Solution (Corrected:)

Form orthogonal basis first: $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ (for instance, using Gram-Schmidt)

Then compute:

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\rangle} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} =$$

Solution (Corrected:)

Form orthogonal basis first: $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ (for instance, using Gram-Schmidt)

Then compute:

$$\frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\rangle} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\left\langle \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\rangle} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + (-2) \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Answer: $\begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix}$.

Example

What is the projection matrix corresponding to orthogonal projection onto $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Example

What is the projection matrix corresponding to orthogonal projection onto $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Solution

The projection matrix is $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$.

Example

What is the projection matrix corresponding to orthogonal projection onto $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Solution

The projection matrix is $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$.

Example

What is the projection matrix corresponding to orthogonal projection onto $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Solution

The projection matrix is $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$.

What would Gram-Schmidt do?

Example

What is the projection matrix corresponding to orthogonal projection onto $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Solution

The projection matrix is $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$.

What would Gram-Schmidt do? $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$.

Example

What is the orthogonal projection of $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ onto $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Example

What is the orthogonal projection of $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ onto $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$?

Solution

The projection is $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$

- The space of all nice functions with period 2π has the natural inner product $\langle f, g \rangle = \int_0^{2\pi} f(x)g(x)dx$.

- The space of all nice functions with period 2π has the natural inner product $\langle f, g \rangle = \int_0^{2\pi} f(x)g(x)dx$.

- The space of all nice functions with period 2π has the natural inner product $\langle f, g \rangle = \int_0^{2\pi} f(x)g(x)dx$.
[in \mathbb{R}^n : $\langle \mathbf{x}, \mathbf{y} \rangle = x_1y_1 + \dots + x_ny_n$]

- The space of all nice functions with period 2π has the natural inner product $\langle f, g \rangle = \int_0^{2\pi} f(x)g(x)dx$.
[in \mathbb{R}^n : $\langle \mathbf{x}, \mathbf{y} \rangle = x_1y_1 + \dots + x_ny_n$]
- The functions

$$1, \cos(x), \sin(x), \cos(2x), \sin(2x), \dots$$

are orthogonal basis for this space.

- The space of all nice functions with period 2π has the natural inner product $\langle f, g \rangle = \int_0^{2\pi} f(x)g(x)dx$.
[in \mathbb{R}^n : $\langle \mathbf{x}, \mathbf{y} \rangle = x_1y_1 + \dots + x_ny_n$]
- The functions

$$1, \cos(x), \sin(x), \cos(2x), \sin(2x), \dots$$

are orthogonal basis for this space.

- Expanding a function $f(x)$ in this basis produces its **Fourier series**

$$f(x) = a_0 + a_1 \cos(x) + b_1 \sin(x) + a_2 \cos(2x) + b_2 \sin(2x) + \dots$$

Example

How do we compute the norm (length) of $\cos(2x)$?

Example

How do we compute the norm (length) of $\cos(2x)$?

Solution

The length of any vector v is always $\sqrt{\langle v, v \rangle}$. So,

Example

How do we compute the norm (length) of $\cos(2x)$?

Solution

The length of any vector v is always $\sqrt{\langle v, v \rangle}$. So,

Example

How do we compute the norm (length) of $\cos(2x)$?

Solution

The length of any vector v is always $\sqrt{\langle v, v \rangle}$. So,

$$||\cos(2x)||^2 = \langle \cos(2x), \cos(2x) \rangle =$$

Example

How do we compute the norm (length) of $\cos(2x)$?

Solution

The length of any vector v is always $\sqrt{\langle v, v \rangle}$. So,

$$||\cos(2x)||^2 = \langle \cos(2x), \cos(2x) \rangle = \int_0^{2\pi} \cos(2x)^2 dx = \pi$$

Example

How do we compute the norm (length) of $\cos(2x)$?

Solution

The length of any vector v is always $\sqrt{\langle v, v \rangle}$. So,

$$||\cos(2x)||^2 = \langle \cos(2x), \cos(2x) \rangle = \int_0^{2\pi} \cos(2x)^2 dx = \pi$$

You don't need to know how to compute this integral.

Example

How do we compute the norm (length) of $\cos(2x)$?

Solution

The length of any vector v is always $\sqrt{\langle v, v \rangle}$. So,

$$||\cos(2x)||^2 = \langle \cos(2x), \cos(2x) \rangle = \int_0^{2\pi} \cos(2x)^2 dx = \pi$$

You don't need to know how to compute this integral. But, you should be able to find the norm (length) of a function!

Example

How do we compute the norm (length) of $\cos(2x)$?

Solution

The length of any vector v is always $\sqrt{\langle v, v \rangle}$. So,

$$||\cos(2x)||^2 = \langle \cos(2x), \cos(2x) \rangle = \int_0^{2\pi} \cos(2x)^2 dx = \pi$$

You don't need to know how to compute this integral. But, you should be able to find the norm (length) of a function! Lengths of other functions work the same way!

Example

If $f(x) = a_0 + a_1 \cos(x) + b_1 \sin(x) + a_2 \cos(2x) + b_2 \sin(2x) + \dots$
how can we compute b_2 ?

Example

If $f(x) = a_0 + a_1 \cos(x) + b_1 \sin(x) + a_2 \cos(2x) + b_2 \sin(2x) + \dots$
how can we compute b_2 ?

Solution

$b_2 \sin(2x)$ is the *orthogonal projection* of f onto the span of $\sin(2x)$.

Example

If $f(x) = a_0 + a_1 \cos(x) + b_1 \sin(x) + a_2 \cos(2x) + b_2 \sin(2x) + \dots$
how can we compute b_2 ?

Solution

$b_2 \sin(2x)$ is the *orthogonal projection* of f onto the span of $\sin(2x)$.

Example

If $f(x) = a_0 + a_1 \cos(x) + b_1 \sin(x) + a_2 \cos(2x) + b_2 \sin(2x) + \dots$
how can we compute b_2 ?

Solution

$b_2 \sin(2x)$ is the *orthogonal projection* of f onto the span of $\sin(2x)$. Hence:

$$b_2 = \frac{\langle f(x), \sin(2x) \rangle}{\langle \sin(2x), \sin(2x) \rangle} =$$

Example

If $f(x) = a_0 + a_1 \cos(x) + b_1 \sin(x) + a_2 \cos(2x) + b_2 \sin(2x) + \dots$
how can we compute b_2 ?

Solution

$b_2 \sin(2x)$ is the *orthogonal projection* of f onto the span of $\sin(2x)$. Hence:

$$b_2 = \frac{\langle f(x), \sin(2x) \rangle}{\langle \sin(2x), \sin(2x) \rangle} = \frac{\int_0^{2\pi} f(x) \sin(2x) dx}{\int_0^{2\pi} \sin^2(2x) dx}$$

Example

If $f(x) = a_0 + a_1 \cos(x) + b_1 \sin(x) + a_2 \cos(2x) + b_2 \sin(2x) + \dots$
how can we compute b_2 ?

Solution

$b_2 \sin(2x)$ is the *orthogonal projection* of f onto the span of $\sin(2x)$. Hence:

$$b_2 = \frac{\langle f(x), \sin(2x) \rangle}{\langle \sin(2x), \sin(2x) \rangle} = \frac{\int_0^{2\pi} f(x) \sin(2x) dx}{\int_0^{2\pi} \sin^2(2x) dx}$$

Projections onto the span of an other function work the same way!

Example

If $f(x) = a_0 + a_1 \cos(x) + b_1 \sin(x) + a_2 \cos(2x) + b_2 \sin(2x) + \dots$
how can we compute b_2 ?

Solution

$b_2 \sin(2x)$ is the *orthogonal projection* of f onto the span of $\sin(2x)$. Hence:

$$b_2 = \frac{\langle f(x), \sin(2x) \rangle}{\langle \sin(2x), \sin(2x) \rangle} = \frac{\int_0^{2\pi} f(x) \sin(2x) dx}{\int_0^{2\pi} \sin^2(2x) dx}$$

Projections onto the span of an other function work the same way!
Again, you should be able to project functions onto the span of a function!

Least Squares

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\iff A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

The projection of \mathbf{b} on the column space of A is then just $A\hat{\mathbf{x}}$.

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

The projection of \mathbf{b} on the column space of A is then just $A\hat{\mathbf{x}}$.

Example

Find the least squares line for the data points
 $(2, 1), (5, 2), (7, 3), (8, 3)$.

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

The projection of \mathbf{b} on the column space of A is then just $A\hat{\mathbf{x}}$.

Example

Find the least squares line for the data points
 $(2, 1), (5, 2), (7, 3), (8, 3)$.

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

The projection of \mathbf{b} on the column space of A is then just $A\hat{\mathbf{x}}$.

Example

Find the least squares line for the data points
 $(2, 1), (5, 2), (7, 3), (8, 3)$.

Solution

Looking for β_1, β_2 such that the line $y = \beta_1 + \beta_2 x$ best fits the data.

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

The projection of \mathbf{b} on the column space of A is then just $A\hat{\mathbf{x}}$.

Example

Find the least squares line for the data points
 $(2, 1), (5, 2), (7, 3), (8, 3)$.

Solution

Looking for β_1, β_2 such that the line $y = \beta_1 + \beta_2 x$ best fits the data.

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

The projection of \mathbf{b} on the column space of A is then just $A\hat{\mathbf{x}}$.

Example

Find the least squares line for the data points
 $(2, 1), (5, 2), (7, 3), (8, 3)$.

Solution

Looking for β_1, β_2 such that the line $y = \beta_1 + \beta_2 x$ best fits the data. The equations $y_i = \beta_1 + \beta_2 x_i$ in matrix form:

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$.

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

$\iff A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

The projection of \mathbf{b} on the column space of A is then just $A\hat{\mathbf{x}}$.

Example

Find the least squares line for the data points
(2, 1), (5, 2), (7, 3), (8, 3).

Solution

Looking for β_1, β_2 such that the line $y = \beta_1 + \beta_2 x$ best fits the data. The equations $y_i = \beta_1 + \beta_2 x_i$ in matrix form:

$$\underbrace{\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \\ 1 & x_4 \end{bmatrix}}_{\text{design matrix } X} \underbrace{\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix}}_{\text{observation vector } y} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}}_{\text{observation vector } y}$$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

$$X^T X =$$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} =$$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix},$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}.$

Solution (continued)

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, *we find* $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}.$

Line of best fit: $y = 2/7 + 5/14x.$

Example

What is the projection of $\begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ onto the column space of $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix}$?

Example

What is the projection of $\begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ onto the column space of $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix}$?

Solution

We found the least squares solution to $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ to be

Example

What is the projection of $\begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ onto the column space of $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix}$?

Solution

We found the least squares solution to $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ to be

Example

What is the projection of $\begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ onto the column space of $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix}$?

Solution

We found the least squares solution to $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ to be

$$\begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$$

Example

What is the projection of $\begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ onto the column space of $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix}$?

Solution

We found the least squares solution to $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ to be

$$\begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$$

So, the projection is $A\hat{\mathbf{x}} = \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix} =$

Example

What is the projection of $\begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ onto the column space of $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix}$?

Solution

We found the least squares solution to $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$ to be

$$\begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$$

So, the projection is $A\hat{\mathbf{x}} = \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix} = \begin{bmatrix} 1 \\ 29/14 \\ 39/14 \\ 22/7 \end{bmatrix}$

Gram-Schmidt

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1, \quad \mathbf{q}_1 = \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned}\mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|}\end{aligned}$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned}\mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|}\end{aligned}$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned} \mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ & \dots & \dots \end{aligned}$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned}\mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ &\dots & \dots\end{aligned}$$

- An **orthogonal matrix** is a square matrix Q with orthonormal columns.

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned}\mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ &\dots & \dots\end{aligned}$$

- An **orthogonal matrix** is a square matrix Q with orthonormal columns.

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned} \mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ & \dots & \dots \end{aligned}$$

- An **orthogonal matrix** is a square matrix Q with orthonormal columns. Equivalently, $Q^T Q =$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned}\mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ &\dots & \dots\end{aligned}$$

- An **orthogonal matrix** is a square matrix Q with orthonormal columns. Equivalently, $Q^T Q = I$ (also true for non-square matrices).

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned}\mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ & \dots & \dots\end{aligned}$$

- An **orthogonal matrix** is a square matrix Q with orthonormal columns. Equivalently, $Q^T Q = I$ (also true for non-square matrices).
- Apply Gram-Schmidt to the (independent) columns of A to obtain the **QR decomposition** $A = QR$.

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned}\mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ & \dots & \dots\end{aligned}$$

- An **orthogonal matrix** is a square matrix Q with orthonormal columns. Equivalently, $Q^T Q = I$ (also true for non-square matrices).
- Apply Gram-Schmidt to the (independent) columns of A to obtain the **QR decomposition** $A = QR$.
 - Q has orthonormal columns (the output vectors of Gram-Schmidt).

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce an orthonormal basis $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\begin{aligned}\mathbf{b}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1, & \mathbf{q}_2 &= \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|} \\ \mathbf{b}_3 &= \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2, & \mathbf{q}_3 &= \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|} \\ & \dots & \dots\end{aligned}$$

- An **orthogonal matrix** is a square matrix Q with orthonormal columns. Equivalently, $Q^T Q = I$ (also true for non-square matrices).
- Apply Gram-Schmidt to the (independent) columns of A to obtain the **QR decomposition** $A = QR$.
 - Q has orthonormal columns (the output vectors of Gram-Schmidt).
 - $R = Q^T A$ is upper triangular.

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

q₁

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

$$\mathbf{q}_1 = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

$$\mathbf{q}_1 = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

$$\mathbf{b}_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 =$$

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

$$\mathbf{q}_1 = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

$$\mathbf{b}_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \frac{1}{\sqrt{5}} \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} =$$

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

$$\mathbf{q}_1 = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

$$\mathbf{b}_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \frac{1}{\sqrt{5}} \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 4/5 \\ -2/5 \\ 1 \end{bmatrix},$$

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

$$\mathbf{q}_1 = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

$$\mathbf{b}_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \frac{1}{\sqrt{5}} \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 4/5 \\ -2/5 \\ 1 \end{bmatrix}, \mathbf{q}_2 =$$

Example

Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}$.

Solution

We apply Gram-Schmidt to the columns of A:

$$\mathbf{q}_1 = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

$$\mathbf{b}_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} - \frac{1}{\sqrt{5}} \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 4/5 \\ -2/5 \\ 1 \end{bmatrix}, \mathbf{q}_2 = \frac{1}{\sqrt{9/5}} \begin{bmatrix} 4/5 \\ -2/5 \\ 1 \end{bmatrix}$$

Solution (continued)

$$\text{Hence: } Q = [\mathbf{q}_1, \mathbf{q}_2] = \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{4}{\sqrt{45}} \\ \frac{2}{\sqrt{5}} & -\frac{2}{\sqrt{45}} \\ 0 & \frac{5}{\sqrt{45}} \end{bmatrix}$$

Solution (continued)

$$\text{Hence: } Q = [\mathbf{q}_1, \mathbf{q}_2] = \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{4}{\sqrt{45}} \\ \frac{2}{\sqrt{5}} & -\frac{2}{\sqrt{45}} \\ 0 & \frac{5}{\sqrt{45}} \end{bmatrix}$$

Solution (continued)

$$\text{Hence: } Q = [\mathbf{q}_1, \mathbf{q}_2] = \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{4}{\sqrt{45}} \\ \frac{2}{\sqrt{5}} & -\frac{2}{\sqrt{45}} \\ 0 & \frac{5}{\sqrt{45}} \end{bmatrix}$$

$$\text{And: } R = Q^T A =$$

Solution (continued)

$$\text{Hence: } Q = [\mathbf{q}_1, \mathbf{q}_2] = \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{4}{\sqrt{45}} \\ \frac{2}{\sqrt{5}} & -\frac{2}{\sqrt{45}} \\ 0 & \frac{5}{\sqrt{45}} \end{bmatrix}$$

$$\text{And: } R = Q^T A = \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} & 0 \\ \frac{4}{\sqrt{45}} & -\frac{2}{\sqrt{45}} & \frac{5}{\sqrt{45}} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix} =$$

Solution (continued)

$$\text{Hence: } Q = [\mathbf{q}_1, \mathbf{q}_2] = \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{4}{\sqrt{45}} \\ \frac{2}{\sqrt{5}} & -\frac{2}{\sqrt{45}} \\ 0 & \frac{5}{\sqrt{45}} \end{bmatrix}$$

$$\text{And: } R = Q^T A = \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} & 0 \\ \frac{4}{\sqrt{45}} & -\frac{2}{\sqrt{45}} & \frac{5}{\sqrt{45}} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \frac{5}{\sqrt{5}} & \frac{1}{\sqrt{5}} \\ 0 & \frac{9}{45} \end{bmatrix}$$

Determinants

- A is invertible $\iff \det(A) \neq 0$

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.
Reverses the sign of the determinant.

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.
Reverses the sign of the determinant.
 - **(scaling)** Multiply all entries in a row by s .

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.
Reverses the sign of the determinant.
 - **(scaling)** Multiply all entries in a row by s .

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.
Reverses the sign of the determinant.
 - **(scaling)** Multiply all entries in a row by s .
Multiplies the determinant by s .

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.
Reverses the sign of the determinant.
 - **(scaling)** Multiply all entries in a row by s .
Multiplies the determinant by s .

$$\begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{vmatrix} \quad \begin{array}{l} R4 \rightarrow R4 - \frac{3}{2}R3 \\ = \end{array}$$

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.
Reverses the sign of the determinant.
 - **(scaling)** Multiply all entries in a row by s .
Multiplies the determinant by s .

$$\begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{vmatrix} \xrightarrow[R4 \rightarrow R4 - \frac{3}{2}R3]{=} \begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & \frac{7}{2} \end{vmatrix} =$$

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.
Reverses the sign of the determinant.
 - **(scaling)** Multiply all entries in a row by s .
Multiplies the determinant by s .

$$\begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{vmatrix} \xrightarrow[R4 \rightarrow R4 - \frac{3}{2}R3]{=} \begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & \frac{7}{2} \end{vmatrix} = 1 \cdot 2 \cdot 2 \cdot \frac{7}{2} =$$

- A is invertible $\iff \det(A) \neq 0$
- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- The **determinant** is characterized by:
 - the normalization $\det I = 1$,
 - and how it is affected by elementary row operations:
 - **(replacement)** Add a multiple of one row to another row.
Does not change the determinant.
 - **(interchange)** Interchange two rows.
Reverses the sign of the determinant.
 - **(scaling)** Multiply all entries in a row by s .
Multiplies the determinant by s .

$$\begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{vmatrix} \xrightarrow[R4 \rightarrow R4 - \frac{3}{2}R3]{=} \begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & \frac{7}{2} \end{vmatrix} = 1 \cdot 2 \cdot 2 \cdot \frac{7}{2} = 14$$

Example

Cofactor expansion is another way to compute determinants.

Example

Cofactor expansion is another way to compute determinants.

Example

Cofactor expansion is another way to compute determinants.

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = -2 \cdot \begin{vmatrix} & - & \\ 3 & & 2 \\ 2 & & 1 \end{vmatrix} + (-1) \cdot \begin{vmatrix} 1 & & 0 \\ & + & \\ 2 & & 1 \end{vmatrix} - 0 \cdot \begin{vmatrix} 1 & & 0 \\ 3 & & 2 \\ & - & \end{vmatrix} \\
 = -2 \cdot (-1) + (-1) \cdot 1 - 0 = 1$$

Example

What is $\begin{vmatrix} 1 & 1 & 1 & 4 \\ -1 & 2 & 2 & 5 \\ 0 & 3 & 3 & 1 \\ 2 & 0 & 0 & 5 \end{vmatrix}$?

Example

What is $\begin{vmatrix} 1 & 1 & 1 & 4 \\ -1 & 2 & 2 & 5 \\ 0 & 3 & 3 & 1 \\ 2 & 0 & 0 & 5 \end{vmatrix}$?

Solution

The determinant is 0 because the matrix is not invertible (second and third columns are the same).

Eigenvalues and eigenvectors

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
It consists of all vectors (plus $\mathbf{0}$) with eigenvalue λ .

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
It consists of all vectors (plus $\mathbf{0}$) with eigenvalue λ .
- Eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_m$ of A corresponding to different eigenvalues are independent.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
It consists of all vectors (plus $\mathbf{0}$) with eigenvalue λ .
- Eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_m$ of A corresponding to different eigenvalues are independent.
- Useful when checking: sum of eigenvalues = sum of diagonal entries.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \det(A - \lambda I) = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
It consists of all vectors (plus $\mathbf{0}$) with eigenvalue λ .
- Eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_m$ of A corresponding to different eigenvalues are independent.
- Useful when checking: sum of eigenvalues = sum of diagonal entries.
- Useful when checking: product of eigenvalues = determinant of A .

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \det(A - \lambda I) = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
It consists of all vectors (plus $\mathbf{0}$) with eigenvalue λ .
- Eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_m$ of A corresponding to different eigenvalues are independent.
- Useful when checking: sum of eigenvalues = sum of diagonal entries.
- Useful when checking: product of eigenvalues = determinant of A .
- Eigenvalues can be complex numbers, even if A is a real matrix.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \det(A - \lambda I) = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
It consists of all vectors (plus $\mathbf{0}$) with eigenvalue λ .
- Eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_m$ of A corresponding to different eigenvalues are independent.
- Useful when checking: sum of eigenvalues = sum of diagonal entries.
- Useful when checking: product of eigenvalues = determinant of A .
- Eigenvalues can be complex numbers, even if A is a real matrix.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
It consists of all vectors (plus $\mathbf{0}$) with eigenvalue λ .
- Eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_m$ of A corresponding to different eigenvalues are independent.
- Useful when checking: sum of eigenvalues = sum of diagonal entries.
- Useful when checking: product of eigenvalues = determinant of A .
- Eigenvalues can be complex numbers, even if A is a real matrix. In that case eigenvectors will be complex, too.

- If $A\mathbf{x} = \lambda\mathbf{x}$, then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
- λ is an eigenvalue of $A \iff \det(A - \lambda I) = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
It consists of all vectors (plus $\mathbf{0}$) with eigenvalue λ .
- Eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_m$ of A corresponding to different eigenvalues are independent.
- Useful when checking: sum of eigenvalues = sum of diagonal entries.
- Useful when checking: product of eigenvalues = determinant of A .
- Eigenvalues can be complex numbers, even if A is a real matrix. In that case eigenvectors will be complex, too. Works just like real vectors.

Diagonalization

- If A is an $n \times n$ matrix, $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **eigenbasis** if

- If A is an $n \times n$ matrix, $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **eigenbasis** if

- If A is an $n \times n$ matrix, $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **eigenbasis** if $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$ for each i (and it is a basis!).

- If A is an $n \times n$ matrix, $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **eigenbasis** if $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$ for each i (and it is a basis!).
- Not all matrices have eigenbases!

- If A is an $n \times n$ matrix, $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **eigenbasis** if $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$ for each i (and it is a basis!).
- Not all matrices have eigenbases!

- If A is an $n \times n$ matrix, $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **eigenbasis** if $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$ for each i (and it is a basis!).
- Not all matrices have eigenbases! But if they do, write:

$$\mathbf{x} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_n\mathbf{v}_n.$$

- If A is an $n \times n$ matrix, $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an **eigenbasis** if $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$ for each i (and it is a basis!).
- Not all matrices have eigenbases! But if they do, write:

$$\mathbf{x} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_n\mathbf{v}_n.$$

- Then it is easy to calculate the action of powers of A on \mathbf{x} :

$$A^k\mathbf{x} = c_1\lambda_1^k\mathbf{v}_1 + c_2\lambda_2^k\mathbf{v}_2 + \cdots + c_n\lambda_n^k\mathbf{v}_n.$$

GOOD LUCK!