

Math 415 - Lecture 17

Linear Transformations

Monday October 5th 2015

Math 415 - Lecture 17

Linear Transformations

Monday October 5th 2015

Textbook reading: Chapter 2.6

Textbook reading: Chapter 2.6

Suggested practice exercises: same as lecture 16

Review

- A map $T : V \rightarrow W$ between vector spaces is **linear** if

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.
 - $T(c\mathbf{x}) = cT(\mathbf{x})$.

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.
 - $T(c\mathbf{x}) = cT(\mathbf{x})$.
- If $\mathbf{x}_1, \dots, \mathbf{x}_n$ is a basis for V , then T is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$:

$$T(\mathbf{v}) = T(c_1\mathbf{x}_1 + \cdots + c_n\mathbf{x}_n) = c_1T(\mathbf{x}_1) + \cdots + c_nT(\mathbf{x}_n).$$

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.
 - $T(c\mathbf{x}) = cT(\mathbf{x})$.
- If $\mathbf{x}_1, \dots, \mathbf{x}_n$ is a basis for V , then T is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$:

$$T(\mathbf{v}) = T(c_1\mathbf{x}_1 + \cdots + c_n\mathbf{x}_n) = c_1T(\mathbf{x}_1) + \cdots + c_nT(\mathbf{x}_n).$$

- Let A be an $m \times n$ matrix.

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.
 - $T(c\mathbf{x}) = cT(\mathbf{x})$.
- If $\mathbf{x}_1, \dots, \mathbf{x}_n$ is a basis for V , then T is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$:

$$T(\mathbf{v}) = T(c_1\mathbf{x}_1 + \cdots + c_n\mathbf{x}_n) = c_1T(\mathbf{x}_1) + \cdots + c_nT(\mathbf{x}_n).$$

- Let A be an $m \times n$ matrix.
 - $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by $T(\mathbf{x}) = A\mathbf{x}$ is linear.

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.
 - $T(c\mathbf{x}) = cT(\mathbf{x})$.
- If $\mathbf{x}_1, \dots, \mathbf{x}_n$ is a basis for V , then T is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$:

$$T(\mathbf{v}) = T(c_1\mathbf{x}_1 + \cdots + c_n\mathbf{x}_n) = c_1T(\mathbf{x}_1) + \cdots + c_nT(\mathbf{x}_n).$$

- Let A be an $m \times n$ matrix.
 - $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by $T(\mathbf{x}) = A\mathbf{x}$ is linear.
 - Every linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is of the form $T(\mathbf{x}) = A\mathbf{x}$.

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.
 - $T(c\mathbf{x}) = cT(\mathbf{x})$.
- If $\mathbf{x}_1, \dots, \mathbf{x}_n$ is a basis for V , then T is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$:

$$T(\mathbf{v}) = T(c_1\mathbf{x}_1 + \cdots + c_n\mathbf{x}_n) = c_1T(\mathbf{x}_1) + \cdots + c_nT(\mathbf{x}_n).$$

- Let A be an $m \times n$ matrix.
 - $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by $T(\mathbf{x}) = A\mathbf{x}$ is linear.
 - Every linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is of the form $T(\mathbf{x}) = A\mathbf{x}$.
- $T : \mathbb{P}_n \rightarrow \mathbb{P}_{n-1}$ defined by $T(p(t)) = p'(t)$ is linear.

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.
 - $T(c\mathbf{x}) = cT(\mathbf{x})$.
- If $\mathbf{x}_1, \dots, \mathbf{x}_n$ is a basis for V , then T is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$:

$$T(\mathbf{v}) = T(c_1\mathbf{x}_1 + \dots + c_n\mathbf{x}_n) = c_1T(\mathbf{x}_1) + \dots + c_nT(\mathbf{x}_n).$$

- Let A be an $m \times n$ matrix.
 - $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by $T(\mathbf{x}) = A\mathbf{x}$ is linear.
 - Every linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is of the form $T(\mathbf{x}) = A\mathbf{x}$.
- $T : \mathbb{P}_n \rightarrow \mathbb{P}_{n-1}$ defined by $T(p(t)) = p'(t)$ is linear. What is its “matrix”?

- A map $T : V \rightarrow W$ between vector spaces is **linear** if
 - $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.
 - $T(c\mathbf{x}) = cT(\mathbf{x})$.
- If $\mathbf{x}_1, \dots, \mathbf{x}_n$ is a basis for V , then T is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$:

$$T(\mathbf{v}) = T(c_1\mathbf{x}_1 + \cdots + c_n\mathbf{x}_n) = c_1T(\mathbf{x}_1) + \cdots + c_nT(\mathbf{x}_n).$$

- Let A be an $m \times n$ matrix.
 - $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by $T(\mathbf{x}) = A\mathbf{x}$ is linear.
 - Every linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is of the form $T(\mathbf{x}) = A\mathbf{x}$.
- $T : \mathbb{P}_n \rightarrow \mathbb{P}_{n-1}$ defined by $T(p(t)) = p'(t)$ is linear. What is its “matrix”?

Nonstandard Bases

Until now we have used the standard bases to describe $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$. Often it is useful to use other bases.

Until now we have used the standard bases to describe $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$. Often it is useful to use other bases.

Theorem (Linear Transformation is Matrix Multiplication)

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let $\mathcal{B} := (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of \mathbb{R}^n and let $\mathcal{C} := (\mathbf{w}_1, \dots, \mathbf{w}_m)$ be a basis of \mathbb{R}^m . Then there is a matrix B such that

$$T(\mathbf{x})_{\mathcal{C}} = B\mathbf{x}_{\mathcal{B}}, \quad \text{for all } \mathbf{x} \in \mathbb{R}^n.$$

Until now we have used the standard bases to describe $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$. Often it is useful to use other bases.

Theorem (Linear Transformation is Matrix Multiplication)

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let $\mathcal{B} := (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of \mathbb{R}^n and let $\mathcal{C} := (\mathbf{w}_1, \dots, \mathbf{w}_m)$ be a basis of \mathbb{R}^m . Then there is a matrix B such that

$$T(\mathbf{x})_{\mathcal{C}} = B\mathbf{x}_{\mathcal{B}}, \quad \text{for all } \mathbf{x} \in \mathbb{R}^n.$$

Until now we have used the standard bases to describe $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$. Often it is useful to use other bases.

Theorem (Linear Transformation is Matrix Multiplication)

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let $\mathcal{B} := (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of \mathbb{R}^n and let $\mathcal{C} := (\mathbf{w}_1, \dots, \mathbf{w}_m)$ be a basis of \mathbb{R}^m . Then there is a matrix B such that

$$T(\mathbf{x})_{\mathcal{C}} = B\mathbf{x}_{\mathcal{B}}, \quad \text{for all } \mathbf{x} \in \mathbb{R}^n.$$

Explicitly,

$$B = [T(\mathbf{v}_1)_{\mathcal{C}} \quad \dots \quad T(\mathbf{v}_n)_{\mathcal{C}}],$$

Example

Let $T \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3a + 1b \\ 1a + 3b \end{bmatrix}$. Then the matrix of T is $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$.

Example

Let $T \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3a + 1b \\ 1a + 3b \end{bmatrix}$. Then the matrix of T is $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$.

Example

Let $T \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3a + 1b \\ 1a + 3b \end{bmatrix}$. Then the matrix of T is $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$.

But let us use, instead of the standard basis, another basis adapted to T . Put

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

Example

Let $T \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3a + 1b \\ 1a + 3b \end{bmatrix}$. Then the matrix of T is $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$.

But let us use, instead of the standard basis, another basis adapted to T . Put

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

What is the coordinate matrix for T with respect to $\mathcal{B} = (\mathbf{b}_1, \mathbf{b}_2)$?

Solution

What do we want?

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$.

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) =$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} =$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} =$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2\mathbf{b}_1,$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2\mathbf{b}_1,$$

$$T(\mathbf{b}_2) = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} =$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2\mathbf{b}_1,$$

$$T(\mathbf{b}_2) = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} =$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2\mathbf{b}_1,$$

$$T(\mathbf{b}_2) = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ 1 \end{bmatrix} =$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2\mathbf{b}_1,$$

$$T(\mathbf{b}_2) = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 4\mathbf{b}_2$$

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2\mathbf{b}_1,$$

$$T(\mathbf{b}_2) = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 4\mathbf{b}_2$$

This means that the coordinate matrix with respect to \mathcal{B} is simply

Solution

What do we want? We want to find a matrix B that relates the coordinate vectors (w.r.t. basis \mathcal{B}) of input vector \mathbf{x} and output vector $T(\mathbf{x})$:

$$T(\mathbf{x})_{\mathcal{B}} = B\mathbf{x}_{\mathcal{B}}.$$

This matrix B has columns $T(\mathbf{b}_1)_{\mathcal{B}}$ and $T(\mathbf{b}_2)_{\mathcal{B}}$. So let us calculate

$$T(\mathbf{b}_1) = T \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 2\mathbf{b}_1,$$

$$T(\mathbf{b}_2) = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 4\mathbf{b}_2$$

This means that the coordinate matrix with respect to \mathcal{B} is simply

$$B = \begin{bmatrix} 2 & 0 \\ 0 & 4 \end{bmatrix}$$

Summary: The linear transformation $T \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3a + 1b \\ 1a + 3b \end{bmatrix}$ has with respect to the standard basis the coordinate matrix A , but with respect to the other basis \mathcal{B} the coordinate B :

Summary: The linear transformation $T \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3a + 1b \\ 1a + 3b \end{bmatrix}$ has with respect to the standard basis the coordinate matrix A , but with respect to the other basis \mathcal{B} the coordinate B :

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 0 \\ 0 & 4 \end{bmatrix}$$

Summary: The linear transformation $T \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3a + 1b \\ 1a + 3b \end{bmatrix}$ has with respect to the standard basis the coordinate matrix A , but with respect to the other basis \mathcal{B} the coordinate B :

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 0 \\ 0 & 4 \end{bmatrix}$$

The linear transformation T is geometrically clear in the \mathcal{B} basis: T is just stretching vectors by a factor 2 along \mathbf{b}_1 and by a factor 4 along \mathbf{b}_2 .

Summary: The linear transformation $T \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3a + 1b \\ 1a + 3b \end{bmatrix}$ has with respect to the standard basis the coordinate matrix A , but with respect to the other basis \mathcal{B} the coordinate B :

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 0 \\ 0 & 4 \end{bmatrix}$$

The linear transformation T is geometrically clear in the \mathcal{B} basis: T is just stretching vectors by a factor 2 along \mathbf{b}_1 and by a factor 4 along \mathbf{b}_2 . So using the standard basis T is an obscure operation on vectors, but using the basis \mathcal{B} it becomes clear. You can say that \mathcal{B} is a basis adapted to T .

Matrices for... Polynomials?

Let P_n be the vector space of polynomials of degree at most n .

Let P_n be the vector space of polynomials of degree at most n .

Example

Consider the map $T : P_2 \rightarrow P_1$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

Describe T by a matrix.

Let P_n be the vector space of polynomials of degree at most n .

Example

Consider the map $T : P_2 \rightarrow P_1$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

Describe T by a matrix.

Solution

Let P_n be the vector space of polynomials of degree at most n .

Example

Consider the map $T : P_2 \rightarrow P_1$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

Describe T by a matrix.

Solution

Wait, what?! We can't multiply a polynomial by a matrix!

Let P_n be the vector space of polynomials of degree at most n .

Example

Consider the map $T : P_2 \rightarrow P_1$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

Describe T by a matrix.

Solution

Wait, what?! We can't multiply a polynomial by a matrix! Use coordinate vectors instead.

Pick bases $\mathcal{A} = (1, t, t^2)$ for P_2 and $\mathcal{B} = (1, t)$ for P_1 . Find a matrix D that does to the coordinate vectors what T does to the polynomials.

Pick bases $\mathcal{A} = (1, t, t^2)$ for P_2 and $\mathcal{B} = (1, t)$ for P_1 . Find a matrix D that does to the coordinate vectors what T does to the polynomials.

$$T(2 + 3t + 4t^2) = 3 + 8t$$

Pick bases $\mathcal{A} = (1, t, t^2)$ for P_2 and $\mathcal{B} = (1, t)$ for P_1 . Find a matrix D that does to the coordinate vectors what T does to the polynomials.

$$T(2 + 3t + 4t^2) = 3 + 8t$$

$$D \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 8 \end{bmatrix}$$

Pick bases $\mathcal{A} = (1, t, t^2)$ for P_2 and $\mathcal{B} = (1, t)$ for P_1 . Find a matrix D that does to the coordinate vectors what T does to the polynomials.

$$T(2 + 3t + 4t^2) = 3 + 8t$$

$$D \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 8 \end{bmatrix}$$

$$T(t^2) = 2t$$

Pick bases $\mathcal{A} = (1, t, t^2)$ for P_2 and $\mathcal{B} = (1, t)$ for P_1 . Find a matrix D that does to the coordinate vectors what T does to the polynomials.

$$T(2 + 3t + 4t^2) = 3 + 8t$$

$$D \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 8 \end{bmatrix}$$

$$T(t^2) = 2t$$

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

Pick bases $\mathcal{A} = (1, t, t^2)$ for P_2 and $\mathcal{B} = (1, t)$ for P_1 . Find a matrix D that does to the coordinate vectors what T does to the polynomials.

$$T(2 + 3t + 4t^2) = 3 + 8t$$

$$D \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 8 \end{bmatrix}$$

$$T(t^2) = 2t$$

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

Formally,

$$D \cdot (f_{\mathcal{A}}) = T(f)_{\mathcal{B}}$$

From the equation

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

From the equation

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

The third column of D is $\begin{bmatrix} 0 \\ 2 \end{bmatrix}$.

From the equation

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

The third column of D is $\begin{bmatrix} 0 \\ 2 \end{bmatrix}$. What are the remaining two columns?

From the equation

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

The third column of D is $\begin{bmatrix} 0 \\ 2 \end{bmatrix}$. What are the remaining two columns?

$$T(1) = 0 \implies D \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

From the equation

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

The third column of D is $\begin{bmatrix} 0 \\ 2 \end{bmatrix}$. What are the remaining two columns?

$$T(1) = 0 \implies$$

$$D \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix},$$

From the equation

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

The third column of D is $\begin{bmatrix} 0 \\ 2 \end{bmatrix}$. What are the remaining two columns?

$$T(1) = 0 \implies$$

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}.$$

From the equation

$$D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

The third column of D is $\begin{bmatrix} 0 \\ 2 \end{bmatrix}$. What are the remaining two columns?

$$T(1) = 0 \implies D \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

$$T(t) = 1 \implies D \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix},$$

$$T(t^2) = 2t \implies D \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}.$$

Hence $D = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

Hence $D = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

Check Take $f(t) = 2 - t + 3t^2$.

Hence $D = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

Check Take $f(t) = 2 - t + 3t^2$. Then the coordinate vector for $f(t)$ is

$$\begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}.$$

Hence $D = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

Check Take $f(t) = 2 - t + 3t^2$. Then the coordinate vector for $f(t)$ is

$$\begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}.$$

Then

$$D \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} = \begin{bmatrix} -1 \\ 6 \end{bmatrix}.$$

Hence $D = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

Check Take $f(t) = 2 - t + 3t^2$. Then the coordinate vector for $f(t)$ is

$$\begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}.$$

Then

$$D \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} = \begin{bmatrix} -1 \\ 6 \end{bmatrix}.$$

On the other hand $T(f(t)) = f'(t) = -1 + 6t$, with coordinate vector $\begin{bmatrix} -1 \\ 6 \end{bmatrix}$.


Matrices for Linear Transformations

Let's organize this.


Let's organize this. Let $T : V \rightarrow W$ be a linear transformation, $\mathcal{A} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be an *input basis* for V , and $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_m\}$ an *output basis* for W .

Let's organize this. Let $T : V \rightarrow W$ be a linear transformation, $\mathcal{A} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be an *input basis* for V , and $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_m\}$ an *output basis* for W . Each vector in V has a coordinate vector in \mathbb{R}^n , each vector in W has a coordinate vector in \mathbb{R}^m . T now corresponds to a matrix from \mathbb{R}^n to \mathbb{R}^m .

Let's organize this. Let $T : V \rightarrow W$ be a linear transformation, $\mathcal{A} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be an *input basis* for V , and $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_m\}$ an *output basis* for W . Each vector in V has a coordinate vector in \mathbb{R}^n , each vector in W has a coordinate vector in \mathbb{R}^m . T now corresponds to a matrix from \mathbb{R}^n to \mathbb{R}^m .



Let's organize this. Let $T : V \rightarrow W$ be a linear transformation, $\mathcal{A} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be an *input basis* for V , and $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_m\}$ an *output basis* for W . Each vector in V has a coordinate vector in \mathbb{R}^n , each vector in W has a coordinate vector in \mathbb{R}^m . T now corresponds to a matrix from \mathbb{R}^n to \mathbb{R}^m .



In the last example this was

$$T(2 + 3t + 4t^2) = 3 + 8t$$

$$A \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 8 \end{bmatrix}$$

Definition

Let $\mathcal{A} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be a basis for V , and $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_m\}$ a basis for W .

The matrix $T_{\mathcal{B}\mathcal{A}}$ representing T with respect to these bases

Definition

Let $\mathcal{A} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be a basis for V , and $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_m\}$ a basis for W .

The matrix $T_{\mathcal{B}\mathcal{A}}$ representing T with respect to these bases

- has n columns (one for each of the \mathbf{x}_j),

Definition

Let $\mathcal{A} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be a basis for V , and $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_m\}$ a basis for W .

The matrix $T_{\mathcal{B}\mathcal{A}}$ representing T with respect to these bases

- has n columns (one for each of the \mathbf{x}_j),
- the j -th column is the coordinate vector of $T(\mathbf{x}_j)$ in the basis \mathcal{B} .

Definition

Let $\mathcal{A} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ be a basis for V , and $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_m\}$ a basis for W .

The matrix $T_{\mathcal{B}\mathcal{A}}$ representing T with respect to these bases

- has n columns (one for each of the \mathbf{x}_j),
- the j -th column is the coordinate vector of $T(\mathbf{x}_j)$ in the basis \mathcal{B} .

$$T_{\mathcal{B}\mathcal{A}} = [T(\mathbf{x}_1)_{\mathcal{B}} \quad T(\mathbf{x}_2)_{\mathcal{B}} \quad \dots \quad T(\mathbf{x}_n)_{\mathcal{B}}]$$

Example

Give the matrix for $T : P_2 \rightarrow P_1$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

in the bases $\mathcal{A} = (1, t, t^2)$ and $\mathcal{B} = (1, t)$.

Example

Give the matrix for $T : P_2 \rightarrow P_1$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

in the bases $\mathcal{A} = (1, t, t^2)$ and $\mathcal{B} = (1, t)$.

Solution

$$T_{\mathcal{B}\mathcal{A}} = [T(1)_{\mathcal{B}} \quad T(t)_{\mathcal{B}} \quad T(t^2)_{\mathcal{B}}] = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

Example

Recall the map T given by $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} y \\ x \end{bmatrix}$. (It reflects every vector in \mathbb{R}^2 across the line $y = x$.)

Example

Recall the map T given by $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} y \\ x \end{bmatrix}$. (It reflects every vector in \mathbb{R}^2 across the line $y = x$.)

- (a) Which matrix A represents T with respect to the standard bases?

Example

Recall the map T given by $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} y \\ x \end{bmatrix}$. (It reflects every vector in \mathbb{R}^2 across the line $y = x$.)

- (a) Which matrix A represents T with respect to the standard bases?
- (b) Which matrix B represents T with respect to the basis $\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix}$?

Solution

$$(a) \quad T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Solution

(a) $T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. So $A = \begin{bmatrix} 0 & * \\ 1 & * \end{bmatrix}$.

Solution

$$(a) \quad T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & * \\ 1 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

Solution

$$(a) \quad T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & * \\ 1 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Solution

$$(a) \quad T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & * \\ 1 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

$$(b) \quad T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

Solution

$$(a) \quad T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & * \\ 1 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

$$(b) \quad T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} -1 \\ 1 \end{bmatrix}. \text{ So } B = \begin{bmatrix} 1 & * \\ 0 & * \end{bmatrix}.$$

Solution

$$(a) \quad T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & * \\ 1 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

$$(b) \quad T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} -1 \\ 1 \end{bmatrix}. \text{ So } B = \begin{bmatrix} 1 & * \\ 0 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} -1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 0 \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

Solution

$$(a) \quad T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & * \\ 1 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

$$(b) \quad T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} -1 \\ 1 \end{bmatrix}. \text{ So } B = \begin{bmatrix} 1 & * \\ 0 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} -1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 0 \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \begin{bmatrix} -1 \\ 1 \end{bmatrix}. \text{ So } B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Solution

$$(a) \quad T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & * \\ 1 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \text{ So } A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

$$(b) \quad T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} -1 \\ 1 \end{bmatrix}. \text{ So } B = \begin{bmatrix} 1 & * \\ 0 & * \end{bmatrix}.$$

$$T\left(\begin{bmatrix} -1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 0 \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \begin{bmatrix} -1 \\ 1 \end{bmatrix}. \text{ So } B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Draw a picture!

Representing Linear Maps by Matrices

Remark

If a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is represented by the matrix A with respect to the standard bases, then $T(\mathbf{x}) = A\mathbf{x}$.

Representing Linear Maps by Matrices

Remark

If a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is represented by the matrix A with respect to the standard bases, then $T(\mathbf{x}) = A\mathbf{x}$.

Representing Linear Maps by Matrices

Remark

If a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is represented by the matrix A with respect to the standard bases, then $T(\mathbf{x}) = A\mathbf{x}$.

Matrix multiplication corresponds to function composition! That is, if T_1, T_2 are represented by A_1, A_2 , then $T_1(T_2(\mathbf{x})) = (A_1A_2)\mathbf{x}$.

Example

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be the linear map such that

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix}.$$

What is the matrix B representing T with respect to the following bases?

$$\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} -1 \\ 2 \end{bmatrix} \text{ for } \mathbb{R}^2,$$

$$\mathbf{y}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{y}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \mathbf{y}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \text{ for } \mathbb{R}^3.$$

Solution

$$T(\mathbf{x}_1) =$$

Solution

$$T(\mathbf{x}_1) = T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right)$$

Solution

$$T(\mathbf{x}_1) = T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$$

Solution

$$\begin{aligned} T(\mathbf{x}_1) = T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) &= T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \\ &= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \\ 10 \end{bmatrix} \end{aligned}$$

Solution

$$\begin{aligned}T(\mathbf{x}_1) &= T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \\&= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \\ 10 \end{bmatrix} \\&= 5 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\end{aligned}$$

Solution

$$\begin{aligned}T(\mathbf{x}_1) &= T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \\&= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \\ 10 \end{bmatrix} \\&= 5 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\&\Rightarrow B = \begin{bmatrix} 5 & * \\ -3 & * \\ 5 & * \end{bmatrix}\end{aligned}$$

$$T(\mathbf{x}_2) =$$

$$T(\mathbf{x}_2) = T\left(\begin{bmatrix} -1 \\ 2 \end{bmatrix}\right)$$

$$T(\mathbf{x}_2) = T\left(\begin{bmatrix} -1 \\ 2 \end{bmatrix}\right) = -T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + 2T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$$

$$\begin{aligned} T(\mathbf{x}_2) = T\left(\begin{bmatrix} -1 \\ 2 \end{bmatrix}\right) &= -T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + 2T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \\ &= -\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 2\begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = \begin{bmatrix} 7 \\ -2 \\ 11 \end{bmatrix} \end{aligned}$$

$$\begin{aligned}T(\mathbf{x}_2) &= T\left(\begin{bmatrix} -1 \\ 2 \end{bmatrix}\right) = -T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + 2T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \\&= -\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 2\begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = \begin{bmatrix} 7 \\ -2 \\ 11 \end{bmatrix} \\&= 7\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - 9\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 4\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\end{aligned}$$

$$\begin{aligned}T(\mathbf{x}_2) &= T\left(\begin{bmatrix} -1 \\ 2 \end{bmatrix}\right) = -T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + 2T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \\&= -\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 2\begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = \begin{bmatrix} 7 \\ -2 \\ 11 \end{bmatrix} \\&= 7\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - 9\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 4\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\&\implies B = \begin{bmatrix} 5 & 7 \\ -3 & -9 \\ 5 & 4 \end{bmatrix}\end{aligned}$$

Remark

A matrix representing T encodes in column j the coefficients of $T(\mathbf{x}_j)$ expressed as a linear combination of $\mathbf{y}_1, \dots, \mathbf{y}_m$.

Recap

What is the Point? Why write $T: V \rightarrow W$ as a matrix?

What is the Point? Why write $T: V \rightarrow W$ as a matrix?

- Replace obscure computations in V and W by transparent computations with matrices.

What is the Point? Why write $T: V \rightarrow W$ as a matrix?

- Replace obscure computations in V and W by transparent computations with matrices.
- Even if $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ (already have standard coordinates), T may be simpler in a different coordinate system.

Summary: Given \mathbf{v} in V , want to calculate $T(\mathbf{v})$ in W .

Summary: Given \mathbf{v} in V , want to calculate $T(\mathbf{v})$ in W . Take an input basis $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$ and an output basis $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$.

Summary: Given \mathbf{v} in V , want to calculate $T(\mathbf{v})$ in W . Take an input basis $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$ and an output basis $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$.

- We know \mathbf{v} if we know the coordinate vector $\mathbf{v}_{\mathcal{A}}$.

Summary: Given \mathbf{v} in V , want to calculate $T(\mathbf{v})$ in W . Take an input basis $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$ and an output basis $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$.

- We know \mathbf{v} if we know the coordinate vector $\mathbf{v}_{\mathcal{A}}$.
- We know $T(\mathbf{v})$ if we know the coordinate vector $T(\mathbf{v})_{\mathcal{B}}$.

Summary: Given \mathbf{v} in V , want to calculate $T(\mathbf{v})$ in W . Take an input basis $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$ and an output basis $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$.

- We know \mathbf{v} if we know the coordinate vector $\mathbf{v}_{\mathcal{A}}$.
- We know $T(\mathbf{v})$ if we know the coordinate vector $T(\mathbf{v})_{\mathcal{B}}$.
- So we know T if we know the matrix $T_{\mathcal{B}\mathcal{A}}$:

$$T(\mathbf{v})_{\mathcal{B}} = T_{\mathcal{B}\mathcal{A}}\mathbf{v}_{\mathcal{A}}.$$

Summary: Given \mathbf{v} in V , want to calculate $T(\mathbf{v})$ in W . Take an input basis $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$ and an output basis $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$.

- We know \mathbf{v} if we know the coordinate vector $\mathbf{v}_{\mathcal{A}}$.
- We know $T(\mathbf{v})$ if we know the coordinate vector $T(\mathbf{v})_{\mathcal{B}}$.
- So we know T if we know the matrix $T_{\mathcal{B}\mathcal{A}}$:

$$T(\mathbf{v})_{\mathcal{B}} = T_{\mathcal{B}\mathcal{A}}\mathbf{v}_{\mathcal{A}}.$$

The output coordinate vector equals the matrix for T times the input coordinate vector.

Example

Let $V = \mathbb{R}^2$ and $W = \mathbb{R}^3$. Let T be the linear map such that

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix}.$$

Example

Let $V = \mathbb{R}^2$ and $W = \mathbb{R}^3$. Let T be the linear map such that

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix}.$$

What is the matrix A representing T with respect to the standard bases? Use that to calculate $T\begin{bmatrix} 2 \\ 3 \end{bmatrix}$.

Example

Let $V = \mathbb{R}^2$ and $W = \mathbb{R}^3$. Let T be the linear map such that

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix}.$$

What is the matrix A representing T with respect to the standard bases? Use that to calculate $T\begin{bmatrix} 2 \\ 3 \end{bmatrix}$.

Solution

The standard bases are

$$\mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad \mathbf{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{y}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \mathbf{y}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$T(\mathbf{x}_1) =$$

$$T(\mathbf{x}_1) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} =$$

$$T(\mathbf{x}_1) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\begin{aligned}T(\mathbf{x}_1) &= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\ &= 1\mathbf{y}_1 + 2\mathbf{y}_2 + 3\mathbf{y}_3\end{aligned}$$

$$T(\mathbf{x}_2) =$$

$$\begin{aligned}T(\mathbf{x}_1) &= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\ &= 1\mathbf{y}_1 + 2\mathbf{y}_2 + 3\mathbf{y}_3\end{aligned}$$

$$T(\mathbf{x}_2) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} =$$

$$\begin{aligned}T(\mathbf{x}_1) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} &= 1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\ &= 1\mathbf{y}_1 + 2\mathbf{y}_2 + 3\mathbf{y}_3\end{aligned}$$

$$T(\mathbf{x}_2) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = 4\mathbf{y}_1 + 0\mathbf{y}_2 + 7\mathbf{y}_3$$

$$\begin{aligned}T(\mathbf{x}_1) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} &= 1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\ &= 1\mathbf{y}_1 + 2\mathbf{y}_2 + 3\mathbf{y}_3\end{aligned}$$

$$T(\mathbf{x}_2) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = 4\mathbf{y}_1 + 0\mathbf{y}_2 + 7\mathbf{y}_3$$

$$\implies A = \begin{bmatrix} 1 & 4 \\ 2 & 0 \\ 3 & 7 \end{bmatrix}$$

$$\text{So } T \begin{bmatrix} 2 \\ 3 \end{bmatrix} = A \begin{bmatrix} 2 \\ 3 \end{bmatrix} =$$

$$\begin{aligned}T(\mathbf{x}_1) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} &= 1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\ &= 1\mathbf{y}_1 + 2\mathbf{y}_2 + 3\mathbf{y}_3\end{aligned}$$

$$T(\mathbf{x}_2) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix} = 4\mathbf{y}_1 + 0\mathbf{y}_2 + 7\mathbf{y}_3$$

$$\implies A = \begin{bmatrix} 1 & 4 \\ 2 & 0 \\ 3 & 7 \end{bmatrix}$$

$$\text{So } T \begin{bmatrix} 2 \\ 3 \end{bmatrix} = A \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 14 \\ 4 \\ 27 \end{bmatrix}$$

Additional Problems

- Suppose $A = \begin{bmatrix} 1 & 2 & 3 & 4 & 1 \\ 2 & 4 & 7 & 8 & 1 \end{bmatrix}$. Find the dimensions and a basis for all four fundamental subspaces of A .

- Suppose $A = \begin{bmatrix} 1 & 2 & 3 & 4 & 1 \\ 2 & 4 & 7 & 8 & 1 \end{bmatrix}$. Find the dimensions and a basis for all four fundamental subspaces of A .
- Suppose A is 5×5 and \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = \mathbf{0}$?

- Suppose $A = \begin{bmatrix} 1 & 2 & 3 & 4 & 1 \\ 2 & 4 & 7 & 8 & 1 \end{bmatrix}$. Find the dimensions and a basis for all four fundamental subspaces of A .
- Suppose A is 5×5 and \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = \mathbf{0}$?
- Let T be the linear map such that

$$T\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix}, \quad T\left(\begin{bmatrix} -1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix}.$$

What is $T\left(\begin{bmatrix} 0 \\ 4 \end{bmatrix}\right)$?