

Proposal to Measure the Cross Section of NC Inclusive Pi0 Interaction Channel

Ariana Hackenburg
Yale University
`ariana.hackenburg@yale.edu`

December 4, 2014

Abstract

Llamas be where it's at.

Contents

List of Figures

List of Tables

1 Overview and History

Definitive proof for neutrino oscillation has been delivered over and again in the last 15 years. Alongside this proof however, have come several notoriously anomalous results. Liquid Scintillator Neutrino Detector (LSND), an oscillation experiment from the 90s, observed an excess of low energy electron anti-neutrino events for $L/E \sim 1\text{m/MeV}$ (Fig 1), explained at the time with a simple 2-neutrino oscillation model (Fig 2). These unexpected results led directly to Mini Booster Neutrino Experiment (MiniBooNE). After 10 years of running, MiniBooNEs data revealed an excess of low energy neutrino events in both neutrino and anti-neutrino mode at energies below and incorporating LSNDs data (Fig 3). These results did several important things: first, MiniBooNEs data supported LSND in anti-neutrino mode, while at the same time refuted the previous 2-neutrino oscillation explanation. MiniBooNE also observed different amounts of excess in neutrino and anti-neutrino mode, leading to more questions about nuclear interactions and cross sections; lastly, it called into question the nature of the observed excess. MiniBooNE is a Cherenkov detector (Fig 4). In a Cherenkov detector, an electron and single photon are indistinguishable; how can we determine if MiniBooNEs excess was due to neutrinos or to photon background?

1.1 The Beginning

Any sort of historical review of neutrino physics at some point comes to Ray Davis, and so we might as well start our story here. In the late 1960s, Ray Davis of Brookhaven National Lab (BNL) ventured 4850ft underground in the Homestake mine with a plan to measure various contributions to solar neutrino flux [?]. Working closely with John Bahcall's theory team at CalTech, Davis' group built a tetrachloroethylene detector sensitive down to 0.814 MeV and primarily to B^8 solar neutrinos[?] a mile underground in Homestake Mine. In 1968, the group released results which revealed that a fraction of the predicted solar neutrinos were missing (it was confirmed by a special task force that the mine's mole people did not eat the missing 2/3 neutrinos). This result became known as "the solar neutrino problem".

set the stage for the next half century drive towards uncovering and resolving anomalies in neutrino physics.

The Homestake experiment ran for 20 years, with results through the duration pointing to the same conclusion: 2/3 of the neutrinos predicted by Bahcall's Standard Solar Model were not accounted for in the data[?]. Throughout the 80's and 90's, a series of experimental results from various collaborations confirmed the mysterious lack of solar neutrinos [?] [?] [?]. Kamiokande II¹, a 3 kton water Cherenkov detector built in Japan in the 80s, was only sensitive to B^8 neutrinos above 6 MeV[?] because of the surrounding radioactivity. Despite its inability to probe the keV energies of the Homestake experiment, Kamiokande II was a highly appealing detector due to its real time depiction of events and its ability to reconstruct both the energies and directions of those events[?]. By the 1990s, the theory of neutrino oscillation as a solution to the solar neutrino problem had become popular, but

¹The original Kamiokande focused primarily on proton decay. After initial data taking, adjustments were made to the original detector in order to make it sensitive enough to study electron recoils from elastic neutrino scatters in water. One example addition was a surrounding layer of water which was intended to decrease background radiation. This "new" experiment was called Kamiokande II

remained unproven. This would change in the later 90s and early 2000's when a series of experimental results flooded the physics community .

n experiment in Hida, Japan called Kamiokande

2) SNO/Kamland/Super K

3) Other current neutrino experiments?

1.2 Experimental Anomalies

Despite the resolution of the solar neutrino problem, further anomalies arose. Two notable examples of this are the results of the Ga radiochemical experiments GALLEX and SAGE. GALLEX was a reactor experiment which took place in the 90s at Gran Sasso, Italy. This radiochemical experiment was sensitive to the pp neutrino flux energies, with a threshold of 233 keV [?]; GALLEX was thus able to probe uncharted waters (Homestake threshold had only been 0.814 keV, making it insensitive to pp neutrinos). After 6 years of running, results showed solar pp flux at ~40% of that expected by the SSM². The validity of this result was confirmed with a ⁵¹Cr source experiment in the detector[?]. SAGE, another Ga experiment set in Russia in the 90s observed a deficit comparable to GALLEX's. A similar test of the SAGE detector's efficiency with a ⁵¹Cr source confirmed it was operating near 100% on the ⁵¹Cr neutrinos[?]; these 2 pieces of the puzzle are examples of the reactor anomaly.

2) Reactor experiments

3) Conclude with oscillation anomalies–LSND, MiniBooNE

Summarize what these things collectively can imply–transition from this into a technology we can use which will help answer some of these questions

2 Theory

Go into some detail about the neutrino oscillations–maybe do a derivation in the appendix. Make sure you quote all relevant resources. Include in this section thus far collection information on :

1) neutrino oscillation frequency Δm^2

2) oscillation amplitude $\sin^2 2\theta$

3 Liquid Argon Time Projection Chambers (LArTPCs)

There are a number of properties that make Liquid Argon (LAr) an appealing detector medium for particle physicists: it is cheap, easy to cool, and transparent to its own scintillation light. When a charged particle travels through a TPC in LAr, it leaves a trail of ionization electrons that are drifted by an applied field to wire planes. Readout from the wires make the Y and Z coordinates of an interaction accessible, while scintillation signals seen by an array of PMTs make the drift direction accessible. Thus a LArTPC with 3 (or

²This only showed conclusive signs for deficit of Be^7 neutrino flux

even 2) planes give us the capability to reconstruct 3 dimensional configurations of events in the detector. It is important to note that LAr scintillates outside the visible spectra at 128nm. In order to see events in LAr, our Photo Multiplier Tubes (PMTs) must be coated in a layer of wavelength shifting material, currently Tetraphenyl Butadiene (TPB).

Liquid Argon Time Projection Chambers (LArTPCs) are ideal detectors for neutrino oscillation experiments with long baselines.

3.1 MicroBooNE

MicroBooNE, the latest in a series of Booster Beam experiments located at Fermilab, is a Liquid Argon Time Projection Chamber (LArTPC) that will investigate the low energy neutrino excess seen by its predecessor, MiniBooNE. Cherenkov detectors, such as MiniBooNE, are limited by their inability to distinguish between single electrons and photons, a task LArTPCs are well suited for. With the high precision reconstruction capabilities of a LArTPC, MicroBooNE will be able to determine with high statistical certainty whether electrons or photons caused the anomalous MiniBooNE low energy excess. Of further interest to MicroBooNE are neutrino-nucleon cross-sections. Cross sections have accounted for much of the uncertainty in recent results from a variety of neutrino experiments, including **** and ***** (reference) and sensitive measurements by MicroBooNE will lead to improved nuclear models and rate predictions. Beyond MicroBooNE, LArTPCs will continue to play a notable role in oscillation physics. LAr1-ND will act as a baseline for improving systematic uncertainties in MicroBooNE and investigating the nature of the MiniBooNE excess, while also acting as a small-scale phase experiment for future, bigger LArTPCs such as LAr1 and LBNE.

4 Current Efforts and Proposal

MicroBooNE, a surface dweller, is subject to a constant bombardment of cosmic radiation from the upper atmosphere. Electromagnetic showers originating from cosmic μ^+ and μ^- are our primary source of background in this case, carrying the ability to mimic MicroBooNE's single electron shower signal (with vertex activity). There are two subsets of events we can break these backgrounds into in order to maximize our rejection efficiency and minimize our signal rejection. The first is the case in which the muon parent is seen in the beam window. This class of shower backgrounds can be minimized fairly effectively with a series of cuts described in DocDB # [technote]. The second and more concerning class of showers are those who either originate from a neutral particle (ie, neutron, π^0), or those whose charged particle parent does not enter the detector. While ionization electrons originating outside the detector can be rejected as "entering showers", gammas produced outside may not Compton scatter/ pair produce until they are inside the detector; this gives it a high probability to be tagged as a signal event if steps are not taken to minimize this background. To filter these background events from data, we can look at various parameters that may distinguish them from our desired signal. A dEdx cut can be used with 94% efficiency to separate pair producing gmmas from cosmic frmo the Compton scatters ****DocDB#[Andrzej technote]. Also useful is the idea that most cosmic radiation will come from above the detector; it is

thus reasonable to expect many of the Compton scattered (and pair produced) showers to have a downward trajectory. In addition, the interaction length in Liquid Argon (14cm) limits the distance we can expect a Compton gamma to travel before interacting (DocDB 3693). We explore these and other parameters in further detail in this study.

In the first, MCC5 generated BNB files are used with (Kazu's) PDG filter to select single electron events. Information such as energy, vertex activity, vertex coordinates and distances along electron trajectory to detector edge (both forward and backward) are stored for 121 electron events / 20,000 total BNB events. This is intended to act as a baseline for the second study.

In the second, the cosmogenic sample described above is examined. In such a case, there are 2 subsets of interactions to sift through which might mimic our primary signal: Compton scatters and pair production. A fiducial volume cut guarantees the vertex is inside the detector volume and dE/dx information from the first 2.4cm of a shower can help distinguish gamma from electron activity; we are then able to distinguish between these two interactions in a true sample. Our first mode of action beyond this is thus to introduce a tag which stores properties of each interaction mode separately (I'm pretty sure I'm going into unnecessary detail here?). Using the selected sample of Compton scatters from Cosmic data, we then investigate properties of the interactions. In particular, we (will) use geometric algorithms to calculate the distance back along the trajectory from vertex to detector edge. Combining this information with interaction length, energy and angle, we are able to successfully (hopefully-not done yet) reduce background Compton scatters while minimally reducing events in the true single electron sample.

Ultimately, these reconstruction effort will lead to a series of interaction channel cross section measurements. To do a cross section measurement, there are several pieces of information that we need to obtain. The cross section is given by:

$$N_{obs} = \sigma * \Phi * \epsilon,$$

where N_{obs} is number of interactions , Φ is the flux of your interaction of interest and ϵ is the efficiency with which you are able to select events. The flux in BNB has been well characterized by the MiniBooNE collaboration and is given in DocBB # []. N is the number of interactions we select in this study. This means that the real lifting comes in estimating your efficiency. There are a few aspects of estimating your efficiency of event selection. The first is that efficiency is energy dependent. One way to overlook this issue is to calculate a differential cross section of -channel of interest- on Ar with respect to energy; we will make this measurement for completeness. Another issue is that estimating the efficiency of our automated reconstruction is tricky. With a limited sample of un-blinded data, we don't have very high statistic and have to rely on our MC truth estimates to estimate our efficiency.

A Aaaarrrrppendix number 1

References

- [1] Raymond Davis Jr and Don S. Harmer, *Solar Neutrinos*. BNL, April 30, 1965
- [2] MEASUREMENT OF THE SOLAR ELECTRON NEUTRINO FLUX WITH THE HOMESTAKE CHLORINE DETECTOR
- [3] Real-time, directional measurement of 8 solar neutrinos in the Kamiokande II detector
- [4] SNO paper
- [5] Results of the whole GALLEX experiment
- [6] GALLEX solar neutrino observations: results for GALLEX IV
- [7] Final results of the ^{51}Cr neutrino source experiments in GALLEX
- [8] The Russian-American Gallium Experiment (SAGE) Cr Neutrino Source Measurement
- [9] Evidence for Oscillation of Atmospheric Neutrinos