

# Homework 6

## MATH 476

Ahad Jiva

May 15, 2025

### Exercise 72

Show that  $\mathbb{E}[S_n] = np$ .

First note that we can break up  $S_n$  Bernoulli trials into  $n$   $S_1$  trials, ie the number of successes in  $n$  Bernoulli trials is the same as the number of successes in one Bernoulli trial  $n$  times, ie  $S_n = S_1 + S_1 + \dots + S_1$ . By the linearity of expected value, we can write

$$\mathbb{E}[S_n] = \mathbb{E}[S_1] + \mathbb{E}[S_1] + \mathbb{E}[S_1] + \dots + \mathbb{E}[S_1].$$

Let's assign a value of 1 to a success in a Bernoulli trial and a value of 0 to a failure. We know the expected value for  $S_1$  is

$$\sum_{x \in \Omega} x \cdot p(x)$$

In this case, there are only two possible outcomes, with a success having probability  $p$  and failure  $1 - p$ . So the expected value is  $(1 \cdot p) + (0 \cdot (1 - p)) = p$ . Then we have

$$\mathbb{E}[S_n] = \mathbb{E}[S_1] + \mathbb{E}[S_1] + \mathbb{E}[S_1] + \dots + \mathbb{E}[S_1] = p + p + p + \dots + p = np$$

as desired.  $\square$

### Exercise 74

Show that  $\mathbb{V}[S_n] = \sigma^2 = npq$ .

First note  $\mathbb{E}[X^2] = ((1^2)p + (0^2)(1 - p))$ . Observe that for a Bernoulli trial,  $\mu = p$ , or  $\mu^2 = p^2$ . So then  $\text{Var}(X) = p - p^2$  since variance  $= \mathbb{E}[X^2] - (\mathbb{E}[X])^2$ . Then we have

$$\text{Var}(S_n) = \text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i) = n(p - p^2) = np(1 - p) = npq$$

as desired.  $\square$

### Exercise 75

Show that  $\int_{-\infty}^{\infty} \phi(x) dx = 1$ , where  $\phi(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2}$ .

Suppose that  $I$  is equal to this integral. We will switch to polar coordinates to prove the identity.

$$I^2 = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} dx \cdot \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}y^2} dy = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-\frac{1}{2}(x^2+y^2)} dx dy$$

Now we can make the substitution  $x^2 + y^2 = r^2$  and change the bounds of integration accordingly.

$$\frac{1}{2\pi} \int_0^{2\pi} \int_0^{\infty} e^{-\frac{1}{2}r^2} r dr d\theta$$

Since the integrand is entirely in  $r$ , we can separate the integrals and cancel the  $d\theta$  integral from 0 to  $2\pi$  with the  $\frac{1}{2\pi}$  outside.

$$\int_0^\infty e^{-\frac{1}{2}r^2} r dr$$

Now we can use u-substitution to finish the improper integral.

$$u = \frac{1}{2}r^2, du = r dr \rightarrow \int_0^\infty e^{-u} du$$

$$-e^{-u} \Big|_0^\infty = 0 + 1 = 1$$

as desired.  $\square$