

FWAM Session B: Function Approximation and Differential Equations

Alex Barnett¹ and **Keaton Burns²**

Wednesday afternoon, 10/30/19

¹Center for Computational Mathematics, Flatiron Institute

²Center for Computational Astrophysics, Flatiron Institute, and Department of Mathematics, MIT

LECTURE 1

interpolation, integration, differentiation, spectral methods

Goals and plan

Overall: graph of $f(x)$ needs ∞ number of points to describe, so how handle f to user-specified accuracy in computer w/ least cost? (bytes/flops)

Goals and plan

Overall: graph of $f(x)$ needs ∞ number of points to describe, so how handle f to user-specified accuracy in computer w/ least cost? (bytes/flops)

- Interpolation: also key to numerical ODE/PDEs...
task: given exact $f(x_j)$ at some x_j , model $f(x)$ at other points x ?
App: cheap but accurate “look-up table” for possibly expensive func.
Contrast: fit noisy data = learning (pdf for) params in model, via likelihood/prior
- Numerical integration:
App: computing expectation values, given a pdf or quantum wavefunc.
App: integral equation methods for PDEs (Jun Wang’s talk)
- Numerical differentiation:
App: build a matrix (linear system) to approximate an ODE/PDE (Lecture II)
App: get gradient ∇f , eg for optimization (cf adjoint methods)

Goals and plan

Overall: graph of $f(x)$ needs ∞ number of points to describe, so how handle f to user-specified accuracy in computer w/ least cost? (bytes/flops)

- Interpolation: also key to numerical ODE/PDEs...
task: given exact $f(x_j)$ at some x_j , model $f(x)$ at other points x ?
App: cheap but accurate “look-up table” for possibly expensive func.
Contrast: fit noisy data = learning (pdf for) params in model, via likelihood/prior
- Numerical integration:
App: computing expectation values, given a pdf or quantum wavefunc.
App: integral equation methods for PDEs (Jun Wang’s talk)
- Numerical differentiation:
App: build a matrix (linear system) to approximate an ODE/PDE (Lecture II)
App: get gradient ∇f , eg for optimization (cf adjoint methods)

Key concepts:

convergence rate, degree of smoothness of f , global vs local,
spectral methods, adaptivity, rounding error & catastrophic cancellation

Goals and plan

Overall: graph of $f(x)$ needs ∞ number of points to describe, so how handle f to user-specified accuracy in computer w/ least cost? (bytes/flops)

- Interpolation: also key to numerical ODE/PDEs...
task: given exact $f(x_j)$ at some x_j , model $f(x)$ at other points x ?
App: cheap but accurate “look-up table” for possibly expensive func.
Contrast: fit noisy data = learning (pdf for) params in model, via likelihood/prior
- Numerical integration:
App: computing expectation values, given a pdf or quantum wavefunc.
App: integral equation methods for PDEs (Jun Wang’s talk)
- Numerical differentiation:
App: build a matrix (linear system) to approximate an ODE/PDE (Lecture II)
App: get gradient ∇f , eg for optimization (cf adjoint methods)

Key concepts:

convergence rate, degree of smoothness of f , global vs local, spectral methods, adaptivity, rounding error & catastrophic cancellation

Plus: good 1D tools, pointers to codes, higher dim methods, opinions!


Interpolation in 1D ($d = 1$)

Say $y_j = f(x_j)$ known at nodes $\{x_j\}$

exact data, not noisy

want interpolant $\tilde{f}(x)$, s.t. $\tilde{f}(x_j) = y_j$

N -pt "grid"




Interpolation in 1D ($d = 1$)

Say $y_j = f(x_j)$ known at nodes $\{x_j\}$ N-pt "grid"


exact data, not noisy

want interpolant $\tilde{f}(x)$, s.t. $\tilde{f}(x_j) = y_j$



hopeless w/o assumptions on f , eg smoothness, otherwise...

- extra info helps, eg f periodic, or $f(x) = \text{smooth} \cdot |x|^{-1/2}$




Interpolation in 1D ($d = 1$)

Say $y_j = f(x_j)$ known at nodes $\{x_j\}$ N-pt "grid"


exact data, not noisy

want interpolant $\tilde{f}(x)$, s.t. $\tilde{f}(x_j) = y_j$



hopeless w/o assumptions on f , eg smoothness, otherwise...

- extra info helps, eg f periodic, or $f(x) = \text{smooth} \cdot |x|^{-1/2}$




Simplest: use value at x_j nearest to x

"snap to grid"

Error $\max_x |\tilde{f}(x) - f(x)| = \mathcal{O}(h)$ as $h \rightarrow 0$

holds if f' bounded; ie f can be nonsmooth but not crazy




Recall notation " $\mathcal{O}(h)$ ": exists $C, h_0 > 0$ s.t. error $\leq Ch$ for all $h < h_0$

Interpolation in 1D ($d = 1$)

Say $y_j = f(x_j)$ known at nodes $\{x_j\}$ N-pt "grid"


exact data, not noisy

want interpolant $\tilde{f}(x)$, s.t. $\tilde{f}(x_j) = y_j$



hopeless w/o assumptions on f , eg smoothness, otherwise...

- extra info helps, eg f periodic, or $f(x) = \text{smooth} \cdot |x|^{-1/2}$




Simplest: use value at x_j nearest to x

"snap to grid"

Error $\max_x |\tilde{f}(x) - f(x)| = \mathcal{O}(h)$ as $h \rightarrow 0$

holds if f' bounded; ie f can be nonsmooth but not crazy




Recall notation " $\mathcal{O}(h)$ ": exists $C, h_0 > 0$ s.t. error $\leq Ch$ for all $h < h_0$

Piecewise linear:

"connect the dots"

max error = $\mathcal{O}(h^2)$ as $h \rightarrow 0$

needs f'' bounded, ie smoother than before



Message: a higher order method is *only* higher order if f smooth enough

Interlude: convergence rates

Should know or measure convergence rate of any method you use

- “effort” parameter N eg # grid-points = $1/h^d$ where h = grid spacing, d = dim

We just saw algebraic conv. error = $\mathcal{O}(N^{-p})$, for order $p = 1, 2$


Interlude: convergence rates

Should know or measure convergence rate of any method you use

- “effort” parameter N eg # grid-points = $1/h^d$ where h = grid spacing, d = dim

We just saw algebraic conv. error = $\mathcal{O}(N^{-p})$, for order $p = 1, 2$

There's only one graph in numerical analysis: “relative error vs effort”





Interlude: convergence rates

Should know or measure convergence rate of any method you use

- “effort” parameter N eg # grid-points = $1/h^d$ where h = grid spacing, d = dim

We just saw algebraic conv. error = $\mathcal{O}(N^{-p})$, for order $p = 1, 2$

There's only one graph in numerical analysis: “relative error vs effort”





Interlude: convergence rates

Should know or measure convergence rate of any method you use

- “effort” parameter N eg # grid-points = $1/h^d$ where h = grid spacing, d = dim

We just saw algebraic conv. error = $\mathcal{O}(N^{-p})$, for order $p = 1, 2$

There's only one graph in numerical analysis: “relative error vs effort”



Note how spectral gets many digits for small N crucial for eg 3D prob.

“spectral” = “superalgebraic”, beats $\mathcal{O}(N^{-p})$ for any p

- how many digits to you want? for 1-digit (10% error), low order ok, easier to code

<rant> test your code w/ *known exact soln* to check error conv. <\rant>

How big is prefactor C in error $\leq Ch^p$? Has asymp. rate even kicked in yet? :)


Higher-order interpolation for smooth f : the local idea

For any target x , use only set of nearest p nodes:

Exists unique degree- $(p-1)$ poly, $\sum_{k=0}^{p-1} c_k x^k$
which matches local data $(x_j, y_j)_{j=1}^p$

generalizes piecewise lin. idea

do **not** eval poly outside its central region!




Higher-order interpolation for smooth f : the local idea

For any target x , use only set of nearest p nodes:

Exists unique degree- $(p-1)$ poly, $\sum_{k=0}^{p-1} c_k x^k$
which matches local data $(x_j, y_j)_{j=1}^p$

generalizes piecewise lin. idea

do **not** eval poly outside its central region!




- error $\mathcal{O}(h^p)$, ie high order, but \tilde{f} not continuous ($\tilde{f} \notin C$) has small jumps
if must have cont, recommend splines, eg cubic $p = 3$: $\tilde{f} \in C^2$, meaning \tilde{f}'' is cont.

Higher-order interpolation for smooth f : the local idea

For any target x , use only set of nearest p nodes:

Exists unique degree- $(p-1)$ poly, $\sum_{k=0}^{p-1} c_k x^k$
which matches local data $(x_j, y_j)_{j=1}^p$



generalizes piecewise lin. idea

do **not** eval poly outside its central region!

- error $\mathcal{O}(h^p)$, ie high order, but \tilde{f} not continuous ($\tilde{f} \notin C$) has small jumps
if must have cont, recommend splines, eg cubic $p = 3$: $\tilde{f} \in C^2$, meaning \tilde{f}'' is cont.


How to find this degree- $(p-1)$ poly?

- 1) crafty: solve square lin sys for coeffs $\sum_{k < p} x_j^k c_k = y_j \quad j = 1, \dots, p$
ie, $V\mathbf{c} = \mathbf{y}$ V ="Vandermonde" matrix, is ill-cond. but works

Higher-order interpolation for smooth f : the local idea

For any target x , use only set of nearest p nodes:

Exists unique degree- $(p-1)$ poly, $\sum_{k=0}^{p-1} c_k x^k$
which matches local data $(x_j, y_j)_{j=1}^p$



generalizes piecewise lin. idea

do **not** eval poly outside its central region!


- error $\mathcal{O}(h^p)$, ie high order, but \tilde{f} not continuous ($\tilde{f} \notin C$) has small jumps
if must have cont, recommend splines, eg cubic $p = 3$: $\tilde{f} \in C^2$, meaning \tilde{f}'' is cont.

How to find this degree- $(p-1)$ poly?

- 1) crafty: solve square lin sys for coeffs $\sum_{k < p} x_j^k c_k = y_j \quad j = 1, \dots, p$
ie, $V\mathbf{c} = \mathbf{y}$ V ="Vandermonde" matrix, is ill-cond. but works


- 2) traditional: barycentric formula $\tilde{f}(x) = \frac{\sum_{j=1}^p \frac{y_j}{x-x_j} w_j}{\sum_{j=1}^p \frac{1}{x-x_j} w_j} \quad w_j = \frac{1}{\prod_{i \neq j} (x_j - x_i)}$
[Tre13, Ch. 5]

Either way, $\tilde{f}(x) = \sum_{j=1}^p y_j \ell_j(x)$ where $\ell_j(x)$ is j th Lagrange basis func:




Global polynomial (Lagrange) interpolation?

Want increase order p . Use *all* data, get single $\tilde{f}(x)$, so $p = N$? “global”
 $p = N = 32$, smooth (analytic) $f(x) = \frac{1}{1+9x^2}$ on $[-1, 1]$: (Runge 1901)



Global polynomial (Lagrange) interpolation?



Want increase order p . Use *all* data, get single $\tilde{f}(x)$, so $p = N$? “global”
 $p = N = 32$, smooth (analytic) $f(x) = \frac{1}{1+9x^2}$ on $[-1, 1]$: (Runge 1901)



- warning: **unif. grid, global interp. fails** → only use locally in central region

Global polynomial (Lagrange) interpolation?

Want increase order p . Use *all* data, get single $\tilde{f}(x)$, so $p = N$ “global”
 $p = N = 32$, smooth (analytic) $f(x) = \frac{1}{1+9x^2}$ on $[-1, 1]$: (Runge 1901)



- warning: **unif. grid, global interp. fails** → only use locally in central region

But exists good choice of nodes...


“Chebyshev”: means non-unif. grid density $\sim \frac{1}{\sqrt{1-x^2}}$

- our first spectral method

$$\max \text{err} = \mathcal{O}(\rho^{-N})$$

exponential conv!

$\rho > 1$ “radius” of largest ellipse in which f analytic



Node choice and adaptivity

Recap: poly approx. $f(x)$ on $[a, b]$ has good & bad node sets $\{x_j\}_{j=1}^N$

Question: Do *you* get to choose the set of nodes at which f known?

- data fitting applications: No (or noisy variants: kriging, Gaussian processes, etc)
use local poly (central region only!), or something stable (eg splines) [GC12]
- almost all else, interp., quadrature, PDE solvers: Yes so pick good nodes!

Node choice and adaptivity

Recap: poly approx. $f(x)$ on $[a, b]$ has good & bad node sets $\{x_j\}_{j=1}^N$

Question: Do *you* get to choose the set of nodes at which f known?

- data fitting applications: No (or noisy variants: kriging, Gaussian processes, etc)
use local poly (central region only!), or something stable (eg splines) [GC12]
- almost all else, interp., quadrature, PDE solvers: Yes so pick good nodes!

Adaptivity idea global is inefficient if f smooth in most places, but not everywhere

Node choice and adaptivity


Recap: poly approx. $f(x)$ on $[a, b]$ has good & bad node sets $\{x_j\}_{j=1}^N$

Question: Do *you* get to choose the set of nodes at which f known?

- data fitting applications: No (or noisy variants: kriging, Gaussian processes, etc)
use local poly (central region only!), or something stable (eg splines) [GC12]
- almost all else, interp., quadrature, PDE solvers: Yes so pick good nodes!

Adaptivity idea

global is inefficient if f smooth in most places, but not everywhere



Node choice and adaptivity


Recap: poly approx. $f(x)$ on $[a, b]$ has good & bad node sets $\{x_j\}_{j=1}^N$

Question: Do *you* get to choose the set of nodes at which f known?

- data fitting applications: No (or noisy variants: kriging, Gaussian processes, etc)
use local poly (central region only!), or something stable (eg splines) [GC12]
- almost all else, interp., quadrature, PDE solvers: Yes so pick good nodes!

Adaptivity idea

global is inefficient if f smooth in most places, but not everywhere



automatically split
(recursively) panels
until $\max \text{err} \leq \epsilon$

via test for local error

1D adaptive interpolator codes to try:

- `github:dbstein/function_generator` py+numba, fast (Stein '19)
- `chebfun` for MATLAB (Trefethen et al.)

App.: replace nasty expensive $f(x)$ by cheap one!

optimal "look-up table"

Global interpolation of periodic functions I

Just did f on intervals $[a, b]$. global interp. (& integr., etc.) of smooth *periodic* f differs!

Periodic: $f(x + 2\pi) = f(x)$ for all x , $f(x) = \sum_{n \in \mathbb{Z}} \hat{f}_k e^{ikx}$ Fourier series

Global interpolation of periodic functions I

Just did f on intervals $[a, b]$. global interp. (& integr., etc.) of smooth *periodic* f differs!

Periodic: $f(x + 2\pi) = f(x)$ for all x , $f(x) = \sum_{n \in \mathbb{Z}} \hat{f}_k e^{ikx}$ Fourier series

Instead of poly's, use **truncated** series $\tilde{f}(x) = \sum_{|k| < N/2} c_k e^{ikx}$ "trig. poly"

Global interpolation of periodic functions I

Just did f on intervals $[a, b]$. global interp. (& integr., etc.) of smooth *periodic* f differs!


Periodic: $f(x + 2\pi) = f(x)$ for all x , $f(x) = \sum_{n \in \mathbb{Z}} \hat{f}_k e^{ikx}$ Fourier series

Instead of poly's, use **truncated** series $\tilde{f}(x) = \sum_{|k| < N/2} c_k e^{ikx}$ "trig. poly"

What's best you can do?

get N coeffs right $c_k = \hat{f}_k$

error \sim size of tail $\{\hat{f}_k\}_{|k| \geq N/2}$



Global interpolation of periodic functions I

Just did f on intervals $[a, b]$. global interp. (& integr., etc.) of smooth *periodic* f differs!


Periodic: $f(x + 2\pi) = f(x)$ for all x , $f(x) = \sum_{n \in \mathbb{Z}} \hat{f}_k e^{ikx}$ Fourier series

Instead of poly's, use truncated series $\tilde{f}(x) = \sum_{|k| < N/2} c_k e^{ikx}$ "trig. poly"

What's best you can do?

get N coeffs right $c_k = \hat{f}_k$

error \sim size of tail $\{\hat{f}_k\}_{|k| \geq N/2}$



How read off c_k from *samples* of f on a grid?

uniform grid best (unlike for poly's!); non-uniform needs linear solve, slow $\mathcal{O}(N^3)$ effort

Uniform grid $x_j = \frac{2\pi j}{N}$, set $c_k = \frac{1}{N} \sum_{j=1}^N e^{ikx_j} f(x_j)$ simply $\mathbf{c} = \text{FFT}[\mathbf{f}]$

Global interpolation of periodic functions I

Just did f on intervals $[a, b]$. global interp. (& integr., etc.) of smooth *periodic* f differs!


Periodic: $f(x + 2\pi) = f(x)$ for all x , $f(x) = \sum_{n \in \mathbb{Z}} \hat{f}_k e^{ikx}$ Fourier series

Instead of poly's, use **truncated** series $\tilde{f}(x) = \sum_{|k| < N/2} c_k e^{ikx}$ "trig. poly"

What's best you can do?

get N coeffs right $c_k = \hat{f}_k$

error \sim size of tail $\{\hat{f}_k\}_{|k| \geq N/2}$



How read off c_k from *samples* of f on a grid?

uniform grid best (unlike for poly's!); non-uniform needs linear solve, slow $\mathcal{O}(N^3)$ effort

Uniform grid $x_j = \frac{2\pi j}{N}$, set $c_k = \frac{1}{N} \sum_{j=1}^N e^{ikx_j} f(x_j)$ simply $\mathbf{c} = \text{FFT}[\mathbf{f}]$

easy to show $c_k = \dots + \hat{f}_{k-N} + \hat{f}_k + \hat{f}_{k+N} + \hat{f}_{k+2N} + \dots$

$= \hat{f}_k$ desired $+ \sum_{m \neq 0} \hat{f}_{k+mN}$ aliasing error, small if tail small

Global interpolation of periodic functions I

Just did f on intervals $[a, b]$. global interp. (& integr., etc.) of smooth *periodic* f differs!


Periodic: $f(x + 2\pi) = f(x)$ for all x , $f(x) = \sum_{n \in \mathbb{Z}} \hat{f}_k e^{ikx}$ Fourier series

Instead of poly's, use **truncated** series $\tilde{f}(x) = \sum_{|k| < N/2} c_k e^{ikx}$ "trig. poly"

What's best you can do?

get N coeffs right $c_k = \hat{f}_k$

error \sim size of tail $\{\hat{f}_k\}_{|k| \geq N/2}$



How read off c_k from *samples* of f on a grid?

uniform grid best (unlike for poly's!); non-uniform needs linear solve, slow $\mathcal{O}(N^3)$ effort

Uniform grid $x_j = \frac{2\pi j}{N}$, set $c_k = \frac{1}{N} \sum_{j=1}^N e^{ikx_j} f(x_j)$ simply $\mathbf{c} = \text{FFT}[\mathbf{f}]$

easy to show $c_k = \dots + \hat{f}_{k-N} + \hat{f}_k + \hat{f}_{k+N} + \hat{f}_{k+2N} + \dots$

$= \hat{f}_k$ desired $+ \sum_{m \neq 0} \hat{f}_{k+mN}$ aliasing error, small if tail small

Summary: given N samples $f(x_j)$, interp. error = truncation + aliasing

a crude bound is $\max_{x \in [0, 2\pi)} |\tilde{f}(x) - f(x)| \leq 2 \sum_{|k| \geq N/2} |\hat{f}_k|$

ie error controlled by sum of tail

Global interpolation of periodic functions II

As grow grid N , how accurate is it? just derived err \sim sum of $|\hat{f}_k|$ in tail $|k| \geq N/2$

Now $\hat{f}_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx = \frac{1}{2\pi} \int_0^{2\pi} f^{(p)}(x) \frac{e^{-ikx}}{(ik)^p} dx$ integr. by parts p times

So for a periodic $f \in C^p$, recall first p derivs of f bounded

$$\hat{f}_k = \mathcal{O}(k^{-p}), \text{ tail sum } \mathcal{O}(N^{1-p}) \quad (p-1)\text{th order acc.} \quad (\text{better: [Tre00]})$$

Global interpolation of periodic functions II

As grow grid N , how accurate is it? just derived $\text{err} \sim \text{sum of } |\hat{f}_k| \text{ in tail } |k| \geq N/2$

Now $\hat{f}_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx = \frac{1}{2\pi} \int_0^{2\pi} f^{(p)}(x) \frac{e^{-ikx}}{(ik)^p} dx$ integr. by parts p times

So for a periodic $f \in C^p$, recall first p derivs of f bounded

$$\hat{f}_k = \mathcal{O}(k^{-p}), \text{ tail sum } \mathcal{O}(N^{1-p}) \quad (p-1)\text{th order acc.} \quad (\text{better: [Tre00]})$$

Example of: f smoother \leftrightarrow faster \hat{f}_k tail decay \leftrightarrow faster convergence

Global interpolation of periodic functions II

As grow grid N , how accurate is it? just derived $\text{err} \sim \text{sum of } |\hat{f}_k| \text{ in tail } |k| \geq N/2$

Now $\hat{f}_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx = \frac{1}{2\pi} \int_0^{2\pi} f^{(p)}(x) \frac{e^{-ikx}}{(ik)^p} dx$ integr. by parts p times

So for a periodic $f \in C^p$, recall first p derivs of f bounded

$$\hat{f}_k = \mathcal{O}(k^{-p}), \text{ tail sum } \mathcal{O}(N^{1-p}) \quad (p-1)\text{th order acc.} \quad (\text{better: [Tre00]})$$

Example of: f smoother \leftrightarrow faster \hat{f}_k tail decay \leftrightarrow faster convergence

Even smoother case: f analytic, so $f(x)$ analytic in some complex strip $|\text{Im } x| \leq \alpha$
then $\hat{f}_k = \mathcal{O}(e^{-\alpha|k|})$, exp. conv. $\mathcal{O}(e^{-\alpha N/2})$ (fun proof: shift the contour)
as with Bernstein ellipse, to get exp. conv. rate need understand f off its real axis (wild!)

Global interpolation of periodic functions II

As grow grid N , how accurate is it? just derived $\text{err} \sim \text{sum of } |\hat{f}_k| \text{ in tail } |k| \geq N/2$

Now $\hat{f}_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx = \frac{1}{2\pi} \int_0^{2\pi} f^{(p)}(x) \frac{e^{-ikx}}{(ik)^p} dx$ integr. by parts p times

So for a periodic $f \in C^p$, recall first p derivs of f bounded

$$\hat{f}_k = \mathcal{O}(k^{-p}), \text{ tail sum } \mathcal{O}(N^{1-p}) \quad (p-1)\text{th order acc.} \quad (\text{better: [Tre00]})$$

Example of: f smoother \leftrightarrow faster \hat{f}_k tail decay \leftrightarrow faster convergence

Even smoother case: f analytic, so $f(x)$ analytic in some complex strip $|\text{Im } x| \leq \alpha$
then $\hat{f}_k = \mathcal{O}(e^{-\alpha|k|})$, exp. conv. $\mathcal{O}(e^{-\alpha N/2})$ (fun proof: shift the contour)
as with Bernstein ellipse, to get exp. conv. rate need understand f off its real axis (wild!)

Smoothest case: “band-limited” f with $\hat{f}_k = 0, |k| > k_{\max}$,
then interpolant exact once $N > 2k_{\max}$

Global interpolation of periodic functions II

As grow grid N , how accurate is it? just derived $\text{err} \sim \text{sum of } |\hat{f}_k| \text{ in tail } |k| \geq N/2$

Now $\hat{f}_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx = \frac{1}{2\pi} \int_0^{2\pi} f^{(p)}(x) \frac{e^{-ikx}}{(ik)^p} dx$ integr. by parts p times

So for a periodic $f \in C^p$, recall first p derivs of f bounded

$$\hat{f}_k = \mathcal{O}(k^{-p}), \text{ tail sum } \mathcal{O}(N^{1-p}) \quad (p-1)\text{th order acc.} \quad (\text{better: [Tre00]})$$

Example of: f smoother \leftrightarrow faster \hat{f}_k tail decay \leftrightarrow faster convergence

Even smoother case: f analytic, so $f(x)$ analytic in some complex strip $|\text{Im } x| \leq \alpha$
then $\hat{f}_k = \mathcal{O}(e^{-\alpha|k|})$, exp. conv. $\mathcal{O}(e^{-\alpha N/2})$ (fun proof: shift the contour)
as with Bernstein ellipse, to get exp. conv. rate need understand f off its real axis (wild!)

Smoothest case: “band-limited” f with $\hat{f}_k = 0, |k| > k_{\max}$,
then interpolant exact once $N > 2k_{\max}$


That’s theory. In real life you always **measure** your conv. order/rate!

Take-home:


- f smooth, periodic, global interpolation w/ uniform grid: spectral acc.
- key to spectral methods. FFT cost $\mathcal{O}(N \log N)$ swaps from $f(x_j)$ grid to \hat{f}_k

Flavor of interpolation in higher dims $d > 1$

If you *can* choose the nodes:
tensor product of 1D interp.
either global
or adaptively refined boxes




periodic, global




adaptive $p = 6 \times 6$ Cheby

Flavor of interpolation in higher dims $d > 1$

If you *can* choose the nodes:
tensor product of 1D interp.
either global
or adaptively refined boxes



periodic, global



adaptive $p = 6 \times 6$ Chebyshev


If *cannot* choose the nodes:
interp. $f(\mathbf{x})$ from scattered data
 $\{\mathbf{x}_i\}$ is hard

Eg google terrain: $f(\mathbf{x})$ rough \rightarrow v. low ord,
see amusing transition in interp. type:

Or if know f smooth

fit local multivariate polynomial

If f noisy and smooth, many
methods
kriging, kernels, ***



height $f(\mathbf{x})$
interp from
unstructured
points in 2D,
kernel method

pock-marks!

interp from
Cartesian grid,
more accurate

Numerical integration (back to $d = 1$)

Task: eval. $\int_a^b f(x)dx$ accurately w/ least number of func. evals, N

“quadrature”: nodes $\{x_j\}$, weights $\{w_j\}$, s.t. $\int_a^b f(x)dx \approx \sum_{j=1}^N w_j f(x_j)$

Idea: get interpolant \tilde{f} thru data $f(x_j) \rightarrow$ *integrate that exactly*

“intepolatory quadrature”

Numerical integration (back to $d = 1$)

Task: eval. $\int_a^b f(x)dx$ accurately w/ least number of func. evals, N

“quadrature”: nodes $\{x_j\}$, weights $\{w_j\}$, s.t. $\int_a^b f(x)dx \approx \sum_{j=1}^N w_j f(x_j)$

Idea: get interpolant \tilde{f} thru data $f(x_j) \rightarrow$ *integrate that exactly*

Examples:

- local piecewise linear \rightarrow composite trapezoid rule

$w_j = h$ except $h/2$ at ends. low-order, err $\mathcal{O}(N^{-2})$, avoid!



- N -node global poly \rightarrow gives $\{w_j\}$ integrating degree $N-1$ exactly

err $\mathcal{O}(\rho^{-N})$

solve lin sys $V^T \mathbf{w} = \{\int_a^b x^k dx\}_{k=0}^{N-1}$ (Newton–Coates)

- better: “Gaussian” $\{x_j, w_j\}$ integrates deg. $2N-1$ exactly! err $\mathcal{O}(\rho^{-2N})$

Adaptive quadrature (Gauss in each panel) excellent: codes `quadgk`, `scipy`, etc

Numerical integration (back to $d = 1$)

Task: eval. $\int_a^b f(x)dx$ accurately w/ least number of func. evals, N

“quadrature”: nodes $\{x_j\}$, weights $\{w_j\}$, s.t. $\int_a^b f(x)dx \approx \sum_{j=1}^N w_j f(x_j)$

Idea: get interpolant \tilde{f} thru data $f(x_j) \rightarrow$ *integrate that exactly*

Examples:

“interpolatory quadrature”

- local piecewise linear \rightarrow composite trapezoid rule

$w_j = h$ except $h/2$ at ends. low-order, err $\mathcal{O}(N^{-2})$, avoid!



- N -node global poly \rightarrow gives $\{w_j\}$ integrating degree $N-1$ exactly

err $\mathcal{O}(\rho^{-N})$ solve lin sys $V^T \mathbf{w} = \{\int_a^b x^k dx\}_{k=0}^{N-1}$ (Newton–Coates)

- better: “Gaussian” $\{x_j, w_j\}$ integrates deg. $2N-1$ exactly! err $\mathcal{O}(\rho^{-2N})$

Adaptive quadrature (Gauss in each panel) excellent: codes `quadgk`, `scipy`, etc

- periodic case: $x_j = \frac{2\pi j}{N}$, $w_j = \frac{2\pi}{N}$ excellent “periodic trap. rule”

easy to check integrates e^{ikx} exactly for $|k| < N$, “Gaussian”

f analytic in $|\operatorname{Im} x| < \alpha$ gives exp. conv. $\mathcal{O}(e^{-\alpha N})$, twice as good as interp!

Numerical integration (back to $d = 1$)

Task: eval. $\int_a^b f(x)dx$ accurately w/ least number of func. evals, N

“quadrature”: nodes $\{x_j\}$, weights $\{w_j\}$, s.t. $\int_a^b f(x)dx \approx \sum_{j=1}^N w_j f(x_j)$

Idea: get interpolant \tilde{f} thru data $f(x_j) \rightarrow$ *integrate that exactly*

Examples:

“interpolatory quadrature”

- local piecewise linear \rightarrow composite trapezoid rule

$w_j = h$ except $h/2$ at ends. low-order, err $\mathcal{O}(N^{-2})$, avoid!



- N -node global poly \rightarrow gives $\{w_j\}$ integrating degree $N-1$ exactly

err $\mathcal{O}(\rho^{-N})$

solve lin sys $V^T \mathbf{w} = \{\int_a^b x^k dx\}_{k=0}^{N-1}$ (Newton–Coates)

- better: “Gaussian” $\{x_j, w_j\}$ integrates deg. $2N-1$ exactly! err $\mathcal{O}(\rho^{-2N})$

Adaptive quadrature (Gauss in each panel) excellent: codes `quadgk`, `scipy`, etc

- periodic case: $x_j = \frac{2\pi j}{N}$, $w_j = \frac{2\pi}{N}$ excellent “periodic trap. rule”

easy to check integrates e^{ikx} exactly for $|k| < N$, “Gaussian”

f analytic in $|\operatorname{Im} x| < \alpha$ gives exp. conv. $\mathcal{O}(e^{-\alpha N})$, twice as good as interp!

demo: `N=14; sum(exp(cos(2*pi*(1:N)/N)))/N - besseli(0,1)`
`ans = 1.3e-15`

Advanced integration

- custom quadr. for singularity eg $f(x) = \text{smooth} \cdot |x|^{-1/2}$
or for arb. set of funcs. “generalized Gaussian quad.” (CCM: Manas Rachh)
- high-order end-corrections to uniform trap. rule (Alpert '99)
-

Advanced integration

- custom quadr. for singularity eg $f(x) = \text{smooth} \cdot |x|^{-1/2}$
or for arb. set of funcs. “generalized Gaussian quad.” (CCM: Manas Rachh)
- high-order end-corrections to uniform trap. rule (Alpert '99)
-

Higher dimensions $d > 1$

For d a few, tensor products of 1D n -node grids in each dim

other coord systems, eg sphere, can use tensor product in (θ, ϕ)

adaptivity works: subdivide into boxes same pic as for interp.

Much higher $d \gg 1$

Now exponential cost $N = n^d$ kills you :(

Are “sparse grids” scaling better rely on funcs aligning w/ axes

Monte Carlo methods: sum N values of $f(\mathbf{x}_j)$ for \mathbf{x}_j random points

error = $\mathcal{O}(N^{-1/2})$ 1/2-order acc, indep of dim d

*** check

Numerical differentiation

Task: given ability to eval. $f(\mathbf{x})$ anywhere, how get $\nabla f(\mathbf{x})$? assume smooth

Numerical differentiation

Task: given ability to eval. $f(\mathbf{x})$ anywhere, how get $\nabla f(\mathbf{x})$? assume smooth

Finite differencing idea, 1D: $f'(x) = \frac{f(x+h)-f(x-h)}{2h} + \mathcal{O}(h^2)$ Taylor's thm

Want smallest error:

suggests taking $h \rightarrow 0$?

Let's see how that goes...


Numerical differentiation

Task: given ability to eval. $f(\mathbf{x})$ anywhere, how get $\nabla f(\mathbf{x})$? assume smooth

Finite differencing idea, 1D: $f'(x) = \frac{f(x+h)-f(x-h)}{2h} + \mathcal{O}(h^2)$ Taylor's thm

Want smallest error:
suggests taking $h \rightarrow 0$?

Let's see how that goes. . .



- shrinking $\mathcal{O}(h^2)$ error gets swamped by a new growing error. . . what?


Numerical differentiation

Task: given ability to eval. $f(\mathbf{x})$ anywhere, how get $\nabla f(\mathbf{x})$? assume smooth

Finite differencing idea, 1D: $f'(x) = \frac{f(x+h)-f(x-h)}{2h} + \mathcal{O}(h^2)$ Taylor's thm

Want smallest error:
suggests taking $h \rightarrow 0$?

Let's see how that goes...



- shrinking $\mathcal{O}(h^2)$ error gets swamped by a new growing error... what?
- CPU arithmetic done only to relative "rounding error" $\epsilon_{\text{mach}} \sim 10^{-16}$
- subtracting v. close $f(x+h)$ and $f(x-h)$: "catastrophic cancellation"
- balance two error types: $h_{\text{best}} \sim \epsilon_{\text{mach}}^{1/3} \sim 10^{-5}$

Essential reading: floating point, backward stability [GC12, Ch. 5–6] [TBI97, Ch. 12–15]

Better approach to differentiation ($d = 1$)

As w/ integration: get interpolant \rightarrow differentiate it exactly [Tre00, Ch. 6]

Get $N \times N$ matrix D acting on func. values $\{f(x_j)\}$ to give $\{f'(x_j)\}$. Has simple formula

Better approach to differentiation ($d = 1$)





As w/ integration: get interpolant \rightarrow differentiate it exactly [Tre00, Ch. 6]

Get $N \times N$ matrix D acting on func. values $\{f(x_j)\}$ to give $\{f'(x_j)\}$. Has simple formula

Examples:

N Chebychev nodes
in $[-1, 1]$

shown: max error in f'



- for N large, the dense D is never formed, merely applied via FFT

spectral solvers for ODE/PDEs. codes: `chebfun`, `PseudoPack`, `dedalus`... Lecture II

Summary: we scratched the surface

Can integrate & differentiate smooth funcs given only point values $f(x_j)$

Both follow from building a good (fast-converging) interpolant

For f smooth in 1D, can & should easily get many (10+) digits accuracy

Concepts:

convergence order/rate	how much effort will you have to spend to get more digits?
smoothness	smooth \Leftrightarrow fast convergence; nonsmooth needs custom methods
global	(one interpolation formula/basis for the whole domain)
vs local	(distinct formulae for x in different regions)
spectral method	global, can exploit FFT, converge very fast
adaptivity	automatically splitting boxes to put nodes only where they need to be
rounding error & catastrophic cancellation	how not shoot self in the foot
tensor products for 2D, 3D	for higher dims: randomized/NN/TN (Sessions C,D,E)

See recommended books at end, and come discuss stuff!

LECTURE II: numerical differential equations

GOALS (maybe not needed)

Reminder of types and applications of diff. eq.

- ODEs: eg pendulum $u''(t) + \sin(u(t)) = 0$ task: solve $u(t)$ efficiently & acc.
needs initial conditions to kick it off eg $u(0) = 1, u'(0) = 0$
eg kinetics of system of well-mixed chemicals, $\mathbf{u}(t)$ is vector
- time-independent PDEs, usually “elliptic” Mike will explain
eg Poisson eqn $\Delta u = g$, $u(\mathbf{x}) =$ chemical concentration or electric potential
 \mathbf{x} means (x, y, \dots) , Δu means Laplacian $\partial^2 u / \partial x^2 + \partial^2 u / \partial y^2 + \dots =$ curvature of u
 $g(\mathbf{x}) =$ volume source of chemical or charge
needs boundary conditions (BCs) on u , or decay at infinity
eg viscous fluid flow: \mathbf{u} is velocity field, Stokes eqns
eg what is ground state of quantum system? solving t-indep. Schrödinger $\Delta u = Eu$
- time-dependent (evolution) PDEs $u(\mathbf{x}, t)$ varies in space and time
eg heat/diffusion equation $\partial u / \partial t + \Delta u = g$ $g(\mathbf{x}, t) =$ source
needs initial and BCs BCs may vary in time...

BCs: simple (eg periodic cube) vs complicated ($u = 0$ on a nasty surface)

[Mike will in next talk overview the three flavors of PDE]

IVPs to BVPs

Motivate focus on BVPs by mentioning:

- Most people familiar with method of lines
- Equivalent “propagator” approach: discretize time, then solve BVP
- Most timesteppers are FD methods (some examples)

Finite difference methods

Basic viewpoint:

- Construct Taylor-series approximation to value at neighboring points
- For N points, expand to $(N-1)$ derivatives (error $\mathcal{O}(h^N)$)
- Eliminate system to get approximation to d -th derivative ($d < N$) (error $\mathcal{O}(h^{N-d})$)

Alternate viewpoint:

- Differentiate the unique interpolating polynomial of deg $N-1$
- Illustrate with centered forward 1st-order scheme, centered 2nd-order scheme (get extra degree for free).

Timestepping with finite difference methods

Consider temporal ODE $u'(t) = f(u(t))$

Explicit methods: simple, just require evaluating $f(u_n)$.

- E.g. Forward Euler: use 1st-order forward difference rule

Unstable for large timesteps

$$u'(t) = -\lambda u(t) \quad \lambda > 0 \quad (1)$$

$$\frac{u_{n+1} - u_n}{k} = -\lambda u_n \quad (2)$$

$$u_{n+1} = (1 - k\lambda)u_n \quad (3)$$

$$\boxed{k\lambda < 2} \quad (4)$$

Implicit methods: stable, but require solving $f(u^{n+1}) = \dots$


- Backward Euler: use 1st-order backward difference rule

Stability, but potentially at high cost

$$u'(t) = -\lambda u(t) \quad \lambda > 0$$

$$\frac{u_{n+1} - u_n}{k} = -\lambda u_{n+1}$$

Advanced finite difference techniques




- Selecting stencils term by term (e.g. upwinding)
- Adaptive stencil selection (e.g. WENO)

Resources: ...

Codes: ...

Finite element methods

Partition domain into elements, represent unknown within each element using basis functions (usually polynomials)



Convert equations to weak form

- Give example
 - Weak form advantages: lowers order, works if lower derivs are integrable (good for discontinuities, hyperbolic), can use tent functions
- Replace weak form with Galerkin/weighted-residual form to get algebraic system

Traditional FEM methods use tent functions, produce continuous solutions. Fluxes at cell boundaries can be computed from internal representation, but can be discontinuous (non conservative).

Finite volume methods

Case of FEM taking functions to be piecewise constant inside elements
Resulting system is exactly conservative (good for hyperbolic), but requires Riemann solve

- Order increased by using neighbors to reconstruct higher order internal representations or fluxes (slope/flux reconstruction)
- Reconstruction is nonlinear to control oscillations around discontinuities (TVD, ENO/WENO).

Very common in CFD, CCA

Advanced finite element methods

- Discontinuous Galerkin: allow discontinuities, need Riemann solvers again
- Spectral elements: move towards large p for better internal representations

Resources: ...

Codes: ...

Spectral methods

Limit of very few elements with very large p : exponential accuracy for smooth problems

Traditional techniques: Fourier spectral methods.

- Fast due to FFT: optimal complexity with exponential accuracy
- Limited to simple geometries / equations with symmetries

Polynomial spectral methods

- More flexible in terms of equations
- Still limited to simple geometries: cubes, spheres, cylinders, etc.
- Still a weak method, but don't integrate by parts. Apply Galerkin directly.

Modern research: sparse methods for arbitrary equations.

Resources: ...

Codes: ...

Boundary integral methods

Use knowledge of PDEs in constructing numerical solutions:

A Green's func G (fundamental soln) needed


eg $\Delta G = \delta$

For linear PDEs dominated by boundary rather than bulk terms, compute solution by forming integral equation of the fundamental solution/Green's function. Reduced dimensionality. Improved conditioning. Low-rank interactions and fast methods. Reconstruct solution in the bulk.

Examples: Stokes flow, Helmholtz, Maxwell, typically with homogeneous media

heavily uses high-order integration methods

- see Jun Wang's talk later today!



Summary

Recommended accessible reading

- [GC12] A Greenbaum and T P Chartier, *Numerical methods*, Princeton University Press, 2012.
- [TBI97] L. N. Trefethen and D. Bau III, *Numerical linear algebra*, SIAM, 1997.
- [Tre00] Lloyd N. Trefethen, *Spectral methods in MATLAB*, Software, Environments, and Tools, vol. 10, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2000.
- [Tre13] L. N. Trefethen, *Approximation theory and approximation practice*, SIAM, 2013,
<http://www.maths.ox.ac.uk/chebfun/ATAP>.