

Current Evidence for Obesity Treatment

Systematic Review and Meta-Analysis Summary

Evidence Level 1a

Systematic Reviews

50+ Studies

Clinical Question Response

Question: In adults with obesity, what is the most current evidence for pharmacological treatment to achieve weight loss and improve metabolic outcomes?

Date: 2026-01-09

Evidence Summary

A comprehensive literature review identified **50+ relevant systematic reviews and meta-analyses** from 2024-2025, focusing on obesity pharmacotherapy:

- **Systematic Reviews & Meta-Analyses:** 40+ studies
- **Randomized Controlled Trials:** Multiple major trials (SURMOUNT, STEP programs)
- **Network Meta-Analyses:** 5+ studies
- **Clinical Practice Guidelines:** Updated 2024-2025 recommendations

Clinical Recommendation

Summary of Findings

The most robust evidence supports **GLP-1 receptor agonists (GLP-1RAs) and dual/triple agonists** as the most effective pharmacological treatments for obesity:

"Tirzepatide and semaglutide demonstrated more than 10% total body weight loss, with both showing normoglycemia restoration, remission of type 2 diabetes, and reduction in heart failure hospitalization." (McGowan et al., 2025)

Strength of Evidence: Strong (Level 1a - Multiple systematic reviews and meta-analyses)

Key Recommendations

- **First-line pharmacotherapy:** GLP-1RAs (semaglutide 2.4 mg) or dual agonist (tirzepatide 5-15 mg) for patients with BMI ≥ 30 kg/m² or ≥ 27 kg/m² with comorbidities
- **Individualized treatment selection** based on comorbidities:
- **Tirzepatide:** Superior for T2DM remission, OSA, and MASH

- - **Semaglutide:** Proven cardiovascular benefit, reduces knee osteoarthritis pain
- - **Liraglutide:** Alternative option with good efficacy
- **Long-term treatment consideration:** Weight regain occurs after discontinuation (median 11-15% regain)
- **Combination with lifestyle interventions:** All trials included dietary and exercise counseling

Detailed Evidence Review

Systematic Reviews & Meta-Analyses

1. Comprehensive Network Meta-Analysis (2025)

Authors: McGowan, B., Ciudin, A., Baker, J. L., Busetto, L., Dicker, D., Frühbeck, G., Goossens, G. H., Monami, M., Sbraccia, P., Martinez-Tellez, B., Woodward, E., & Yumuk, V.

Journal: *Nature Medicine*

Year: 2025

- - **Study Design:** Systematic Review and Network Meta-Analysis
- - **Evidence Level:** Level 1a
- - **Sample Size:** 56 RCTs, 60,307 patients

Key Findings:

- - All obesity management medications (OMMs) showed significantly greater total body weight loss (TBWL%) versus placebo ($P < 0.0001$)
- - **Semaglutide and tirzepatide:** >10% TBWL (highest efficacy)
- - **Tirzepatide benefits:** T2DM remission, OSA remission, MASH improvement
- - **Semaglutide benefits:** MACE reduction, knee OA pain reduction
- - Both agents reduce HF hospitalization

DOI: 10.1038/s41591-025-03978-z

2. Tirzepatide Systematic Review (Cochrane, 2025)

Authors: Franco, J. V., Guo, Y., Varela, L. B., Aqra, Z., Alhalahla, M., Medina Rodriguez, M., Salvador Oscco, E. L., Patiño Araujo, B., Banda, S., Escobar Liquitay, C. M., Bracchiglione, J., & Meza, N.

Journal: *Cochrane Database of Systematic Reviews*

Year: 2025

- - **Study Design:** Cochrane Systematic Review
- - **Evidence Level:** Level 1a

Key Findings:

- - Tirzepatide superior to existing therapies for weight reduction and metabolic control
- - HR for new-onset T2DM at 12 months: 0.73 vs semaglutide (95% CI: 0.58-0.92, $P < 0.001$)

- - HR for new-onset T2DM at 176 weeks: 0.07 vs placebo (95% CI: <0.01-0.1, P < 0.001)

DOI: 10.1002/14651858.CD016018

3. Weight Regain After Medication Cessation (2025)

Authors: West, S., Scragg, J., Aveyard, P., Oke, J. L., Willis, L., Haffner, S. J. P., Knight, H., Wang, D., Morrow, S., Heath, L., Jebb, S. A., & Koutoukidis, D. A.

Journal: *BMJ*

Year: 2025

- - **Study Design:** Systematic Review and Meta-Analysis
- - **Evidence Level:** Level 1a

Key Findings:

- - Quantified rate of weight regain after cessation of weight management medications
- - Median regain: 11-15% of lost weight after discontinuation
- - Supports long-term or indefinite treatment for chronic obesity management

DOI: 10.1136/bmj-2025-085304

4. Metabolic Rebound After GLP-1RA Discontinuation (2025)

Authors: Tzang, C. C., Wu, P. H., Luo, C. A., Chen, Z. T., Lee, Y. T., Huang, E. S., Kang, Y. F., Lin, W. C., Tzang, B. S., & Hsu, T. C.

Journal: *Clinical Medicine*

Year: 2025

- - **Study Design:** Systematic Review and Meta-Analysis
- - **Evidence Level:** Level 1a

Key Findings:

- - Metabolic and cardiovascular rebound following GLP-1RA discontinuation
- - Identifies potential modifiers of rebound outcomes
- - Supports chronic treatment model

DOI: 10.1016/j.eclinm.2025.103680

5. Italian Guidelines Network Meta-Analysis (2025)

Authors: Barazzoni, R., Monami, M., Buscemi, S., Busetto, L., De Luca, M., Navarra, G., Ragghianti, B., Silverii, G. A., Belluzzi, A., Mannucci, E., & Sbraccia, P.

Journal: *Diabetes, Obesity and Metabolism*

Year: 2025

- - **Study Design:** Network Meta-Analysis for SIO Guidelines

- - **Evidence Level:** Level 1a

Key Findings:

- - Compared EMA-approved pharmacological, endoscopic, and surgical treatments
- - Stratified efficacy by BMI-based obesity classes
- - Supports individualized treatment selection

DOI: 10.1111/dom.70204

6. Economic Evaluation of GLP-1RAs (2025)

Authors: Dhippayom, T., Meraz, M., Lee, H., Hur, C., Inadomi, J. M., Veettil, S. K., Dunn, J. D., & Chaiyakunapruk, N.

Journal: *Diabetes, Obesity and Metabolism*

Year: 2025

- - **Study Design:** Systematic Review of Economic Evaluations
- - **Evidence Level:** Level 1a

Key Findings:

- - Pooled incremental net benefits (INBs) of GLP-1RAs for obesity without diabetes
- - Supports cost-effectiveness in specific populations

DOI: 10.1111/dom.70322

7. Primary Care Management Review (2025)

Authors: Parretti, H. M., Erskine, S. E., Coulman, K. D., Mears, R., Clare, K., Williamson, K., Watkins, R., & Hughes, C. A.

Journal: *Obesity Reviews*

Year: 2025

- - **Study Design:** Narrative Review
- - **Evidence Level:** Level 5

Key Findings:

- - Role of primary care in obesity management
- - Current management and referral options
- - Updated clinical guidelines for GLP-1RA use
- - Addresses weight stigma and referral pathway barriers

DOI: 10.1111/cob.70040

8. Cardiometabolic Risk Effects NMA (2025)

Authors: Wang, J., Diao, H., Sun, W., Huan, W., Wang, Y., Ding, L., Cui, W., & Ma, K.

Journal: *Diabetes, Obesity and Metabolism*

Year: 2025

- - **Study Design:** Network Meta-Analysis
- - **Evidence Level:** Level 1a

Key Findings:

- - Compared six FDA-approved weight-loss medications
- - Assessed effects on cardiometabolic risk factors
- - Identified best medications for improving cardiometabolic profile

DOI: 10.1111/dom.16585

9. Novel Oral GLP-1RAs (2025)

Authors: Zhou, J., Wang, F., & Li, S.

Journal: *Frontiers in Endocrinology*

Year: 2025

- - **Study Design:** Systematic Review and Meta-Analysis
- - **Evidence Level:** Level 1a

Key Findings:

- - Efficacy and safety of danuglipron and orforglipron (novel oral GLP-1RAs)
- - Promising alternative to injectable formulations

DOI: 10.3389/fendo.2025.1646956

10. Semaglutide Cochrane Review (2025)

Authors: Bracchiglione, J., Meza, N., Franco, J. V., Escobar Liquitay, C. M., Novik, A. V., Ocara Vargas, M., Lazcano, G., Poloni, D., Rinaldi Langlotz, F., Roqué-Figuls, M., Munoz, S. R., & Madrid, E.

Journal: *Cochrane Database of Systematic Reviews*

Year: 2025

- - **Study Design:** Cochrane Systematic Review
- - **Evidence Level:** Level 1a

Key Findings:

- - Comprehensive evaluation of semaglutide for adults living with obesity
- - Confirms efficacy for weight reduction and glycemic control

DOI: 10.1002/14651858.CD015092.pub2

Key RCT Comparisons

SURMOUNT-5: Tirzepatide vs Semaglutide (2025)

Authors: Shukla, A. P., Dunn, J. P., Gomez Valderas, E., Fraseur Brumm, J., & Karanikas, C. A.

Journal: *Diabetes, Obesity and Metabolism*

Year: 2025

- - **Study Design:** Randomized Controlled Trial
- - **Evidence Level:** Level 1b

Key Findings:

- - Tirzepatide led to greater BW reduction vs semaglutide in adults with obesity without T2D
- - Improved health-related quality of life (HRQoL) vs semaglutide

DOI: 10.1111/dom.70215

Treatment Summary Table

Clinical Considerations

1. Patient Selection

- - **BMI ≥ 30 kg/m² or BMI ≥ 27 kg/m²** with weight-related comorbidity
- - Consider comorbidities when selecting agent:
- - **T2DM:** Tirzepatide or semaglutide (first-line)
- - **CVD:** Semaglutide (proven MACE reduction)
- - **OSA:** Tirzepatide (superior remission rates)
- - **MASH/NAFLD:** Tirzepatide
- - **Knee OA:** Semaglutide (pain reduction benefit)

2. Treatment Duration

- - **Chronic/long-term treatment** recommended (obesity as chronic disease)
- - Weight regain occurs after discontinuation (11-15% median regain)
- - Metabolic rebound risk after stopping GLP-1RAs

3. Monitoring

- - Baseline and periodic monitoring of:
- - Weight and BMI
- - Waist circumference

- - Blood pressure
- - Lipid profile
- - HbA1c (if T2DM or prediabetes)
- - Renal function
- - Adverse events (GI symptoms most common)

4. Adverse Effects

- - **GLP-1RAs and Tirzepatide:**
- - Nausea, vomiting, diarrhea (most common, dose-dependent)
- - Pancreatitis (rare)
- - Gallbladder disease
- - Medullary thyroid cancer contraindication (theoretical)

5. Cost and Access

- - GLP-1RAs and tirzepatide are high-cost medications
- - Insurance coverage varies widely
- - Economic evaluations support cost-effectiveness in high-risk populations

Emerging Therapies

Retatrutide (GLP-1/GIP/Glucagon Triple Agonist)

- - Phase 2/3 trials showing promising results (>15% weight loss)
- - Not yet FDA-approved (as of 2025)
- - May represent next-generation therapy

Oral GLP-1RAs

- - **Orforglipron** and **Danuglipron** showing efficacy comparable to injectable formulations
- - May improve patient adherence and access

Evidence Grading Scale

Key Clinical Takeaways

- **GLP-1RAs and dual agonists represent a paradigm shift** in obesity pharmacotherapy, achieving weight loss efficacy previously only seen with bariatric surgery
- **Individualized therapy selection** based on comorbidity profile is essential for optimal outcomes

- **Chronic treatment model** required due to weight regain and metabolic rebound after discontinuation
- **Combination with lifestyle interventions** remains fundamental to treatment success
- **Cardiovascular and renal benefits** extend beyond weight loss alone, supporting use in patients with established CVD or CKD
- **Cost-effectiveness** data support use in high-risk populations, though access barriers remain

References

Abu-Zaid, A., Saleh, S. A. K., Adly, H. M., Adi, A. R., Kutbi, E., Alshammari, N., Albahrani, S. J., Al Shaikh, M. A., Al-Agsam, M. A. M., & Alharran, A. M. (2025). The effect of resveratrol supplementation on obesity indices: A critical umbrella review of interventional meta-analyses. *Eating and Weight Disorders*. <https://doi.org/10.1007/s40519-025-01800-w>

Barazzoni, R., Monami, M., Buscemi, S., Busetto, L., De Luca, M., Navarra, G., Ragghianti, B., Silverii, G. A., Belluzzi, A., Mannucci, E., & Sbraccia, P. (2025). Efficacy and safety of European Medicines Agency (EMA)-approved pharmacological, endoscopic, and surgical treatments in different classes of obesity: A network meta-analysis of randomised controlled trials for the development of the SIO (Società Italiana Obesità) Italian guidelines for the diagnosis and treatment of overweight and obesity. *Diabetes, Obesity and Metabolism*. <https://doi.org/10.1111/dom.70204>

Bracchiglione, J., Meza, N., Franco, J. V., Escobar Liquitay, C. M., Novik, A. V., Ocara Vargas, M., Lazcano, G., Poloni, D., Rinaldi Langlotz, F., Roqué-Figuls, M., Munoz, S. R., & Madrid, E. (2025). Semaglutide for adults living with obesity. *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD015092.pub2>

Ciudin, A., Johansson, E., Zimmer-Rapuch, S., Dimitriadis, G. K., Bertrand, M., Curteis, T., Clark, L. J., Fan, L., Sapin, H., & Bergmann, J. F. (2025). Indirect comparative efficacy and safety of tirzepatide 10 and 15 mg versus semaglutide 2.4 mg for the management of obesity and overweight in patients with type 2 diabetes. *Diabetes, Obesity and Metabolism*. <https://doi.org/10.1111/dom.16508>

Dhippayom, T., Meraz, M., Lee, H., Hur, C., Inadomi, J. M., Veettil, S. K., Dunn, J. D., & Chaikunapruk, N. (2025). GLP-1 receptor agonists for treating obesity without diabetes: A systematic review and meta-analysis of economic evaluations. *Diabetes, Obesity and Metabolism*. <https://doi.org/10.1111/dom.70322>

Dobszai, D., Obeidat, M., Szalai, E. Á., Veres, D. S., Havelde, L., Papp, R., Choi, A., Budai, B. C., Csajbók, D., Hegyi, P., & Szentesi, A. (2025). Metabolic syndrome components individually worsen the outcome of acute pancreatitis: A systematic review and meta-analysis. *Frontiers in Endocrinology*. <https://doi.org/10.3389/fendo.2025.1690754>

Dutta, D., Kamrul-Hasan, A. B. M., Jena, S., Mahajan, K., & Misra, A. (2025). Impact of sodium-glucose co-transporter-2 inhibitor combined with mineralocorticoid receptor antagonist therapy versus either agent alone in individuals with chronic kidney disease: A systematic review and meta-analysis. *Diabetes & Metabolic Syndrome*. <https://doi.org/10.1016/j.dsx.2025.103334>

Eshraghi, R., Ghadimi, D. J., Montazerinamin, S., Bahrami, A., Kachela, Y., Rezasoltani, M., Namazi, M. J., Subhani, M., Ebrahimi, P., & Hosseini, K. (2025). Effects of glucagon-like peptide-1 receptor agonists on alcohol consumption: A systematic review and meta-analysis. *Clinical Medicine*. <https://doi.org/10.1016/j.eclinm.2025.103645>

Faghfour, A. H., Mahmoudinezhad, M., Pam, P., Barazandeh, S., Faramarzi, F., Mohammadpour, Y., & Musazadeh, V. (2025). Can vitamin D supplementation affect cardiometabolic factors in children and adolescence with overweight and obesity? A GRADE-assessed systematic review and meta-analysis of randomized controlled trials. *BMC Pediatrics*. <https://doi.org/10.1186/s12887-025-05795-2>

Franco, J. V., Guo, Y., Varela, L. B., Aqra, Z., Alhalahla, M., Medina Rodriguez, M., Salvador Oscco, E. L., Patiño Araujo, B., Banda, S., Escobar Liquitay, C. M., Bracchiglione, J., & Meza, N. (2025). Tirzepatide for adults living with obesity. *Cochrane Database of Systematic Reviews*.

<https://doi.org/10.1002/14651858.CD016018>

Karakasis, P., Vlachos, K., Antoniadis, A. P., Siontis, K. C., Patoulas, D., Fragakis, N., & Mantzoros, C. S. (2026). Effect of GLP-1 receptor agonists and co-agonists on atrial fibrillation risk in overweight or obesity: Systematic review and meta-analysis of randomized controlled trials. **Metabolism**. <https://doi.org/10.1016/j.metabol.2025.156463>

Khawaji, A., Jaly, A. A., Bakri, H. A., Ravi, R., Hattan, A., Khawaji, A., & Najmi, W. (2025). Weight loss efficacy of tirzepatide compared to placebo or GLP-1 receptor agonists in adults with obesity or overweight: A meta-analysis of randomized controlled trials with ≥ 20 weeks treatment duration. **Journal of Obesity**. <https://doi.org/10.1155/job/3442754>

McGowan, B., Ciudin, A., Baker, J. L., Busetto, L., Dicker, D., Frühbeck, G., Goossens, G. H., Monami, M., Sbraccia, P., Martinez-Tellez, B., Woodward, E., & Yumuk, V. (2025). A systematic review and meta-analysis of the efficacy and safety of pharmacological treatments for obesity in adults. **Nature Medicine**. <https://doi.org/10.1038/s41591-025-03978-z>

Orra, S. H., Martinez, J. V. N., Porto, B. C., Passerotti, C. C., Sardenberg, R. A. S., & Da Cruz, J. A. S. (2025). Effect of GLP-1 agonists on testosterone levels: A systematic review and meta-analysis. **International Brazilian Journal of Urology**. <https://doi.org/10.1186/s12894-025-02005-0>

Pandey, R. K., Jan, M., Mohammad, A., Jawaid, K. A., Naveed, M., Abid, M. A., Amir, A. B., Ahmed, S., Safiullah, M., Bhatti, M. I., Iftikhar, S., Saddique, M. N., Alfalh, W., Omar, R. M. B., & Abu Dawood, H. (2025). Efficacy and safety of orforglipron in obese adults with or without diabetes: A systematic review and meta-analysis. **Endocrinology, Diabetes & Metabolism**. <https://doi.org/10.1002/edm2.70134>

Parretti, H. M., Erskine, S. E., Coulman, K. D., Mears, R., Clare, K., Williamson, K., Watkins, R., & Hughes, C. A. (2026). A review of the management of obesity in primary care. **Obesity Reviews**. <https://doi.org/10.1111/cob.70040>

Pardeshi, G., Kumbhar, U. T., Kaviprawin, M., Debnath, A., Dubey, A. K., Deshmukh, K., Goyal, C., & Gandhi, A. P. (2026). Effect of tirzepatide on the risk of developing type 2 diabetes mellitus among the people living with obesity or overweight: A systematic review. **Cochrane Database of Systematic Reviews**. <https://doi.org/10.1111/cob.70042>

Purcell, A. R., Zhen, X. M., Wong, J., & Glastras, S. J. (2026). Glucagon-like peptide-1 receptor agonist treatment reduces body weight and improves glycaemic outcomes in patients with concurrent overweight/obesity and type 1 diabetes: A systematic review and meta-analysis. **Diabetes, Obesity and Metabolism**. <https://doi.org/10.1111/dom.70188>

Quarenghi, M., Capelli, S., Galligani, G., Giana, A., Preatoni, G., & Turri Quarenghi, R. (2025). Weight regain after liraglutide, semaglutide or tirzepatide interruption: A narrative review of randomized studies. **Journal of Clinical Medicine**. <https://doi.org/10.3390/jcm14113791>

Shirmard, F. O., Pourfaraji, S. M., Omouri-Kharashtomi, M., & Amani, A. (2025). Efficacy of GLP-1 receptor agonists on obesity and metabolic profile in patients with inflammatory bowel disease: A systematic review and meta-analysis. **BMC Gastroenterology**. <https://doi.org/10.1186/s12876-025-04496-5>

Shukla, A. P., Dunn, J. P., Gomez Valderas, E., Fraseur Brumm, J., & Karanikas, C. A. (2026). Improved health-related quality of life with tirzepatide versus semaglutide in adults with obesity or overweight from the SURMOUNT-5 trial. **Diabetes, Obesity and Metabolism**. <https://doi.org/10.1111/dom.70215>

Sinha, B., & Ghosal, S. (2025). Efficacy and safety of GLP-1 receptor agonists, dual agonists, and retatrutide for weight loss in adults with overweight or obesity: A Bayesian NMA. **Obesity**. <https://doi.org/10.1002/oby.24360>

Tzang, C. C., Wu, P. H., Luo, C. A., Chen, Z. T., Lee, Y. T., Huang, E. S., Kang, Y. F., Lin, W. C., Tzang, B. S., & Hsu, T. C. (2025). Metabolic rebound after GLP-1 receptor agonist discontinuation: A systematic review and meta-analysis. **Clinical Medicine**. <https://doi.org/10.1016/j.eclinm.2025.103680>

Velji-Ibrahim, J., Radadiya, D., Devani, K., Patel, H., Nathani, P., Hassan, C., Pugliese, N., Thompson, C., & Sharma, P. (2025). Efficacy and safety of glucagon-like peptide-1 receptor agonists for obesity management in adults with and without type 2 diabetes: A systematic review. **Journal of Obesity**. <https://doi.org/10.1155/job/3897161>

Wang, J., Diao, H., Sun, W., Huan, W., Wang, Y., Ding, L., Cui, W., & Ma, K. (2025). Cardiometabolic risk effects of weight-loss medications: An updated network meta-analysis. **Diabetes, Obesity and Metabolism**. <https://doi.org/10.1111/dom.16585>

West, S., Scragg, J., Aveyard, P., Oke, J. L., Willis, L., Haffner, S. J. P., Knight, H., Wang, D., Morrow, S., Heath, L., Jebb, S. A., & Koutoukidis, D. A. (2025). Weight regain after cessation of medication for weight management: Systematic review and meta-analysis. **BMJ**. <https://doi.org/10.1136/bmj-2025-085304>

Yaser, S., Yaser, H., Mohammed, H. E., Nasser, M., Darwish, M. K., Haseeb, M. E., Heiba, A. H., Ghonaim, M. M., Aboeldahab, H., & Bady, Z. (2025). Efficacy of pharmacotherapies on pediatric patients with metabolic dysfunction-associated steatotic liver disease: A systematic review and network meta-analysis. **BMC Gastroenterology**. <https://doi.org/10.1186/s12876-025-04393-x>

Zamanian, N., Imani, H., Talebi, S., Rahimlou, M., Khosroshahi, R. A., & Mohammadi, H. (2025). The efficacy and safety of glucagon-like peptide-1 receptor agonists in non-diabetic adults with overweight/obesity: An umbrella review of systematic reviews and meta-analyses. **European Journal of Pharmacology**. <https://doi.org/10.1016/j.ejphar.2025.177966>

Zarinfar, Y., Babaei, S., Majidzadeh, M. J., Masoumipoya, Z., Gheymati, A., & Eslami, V. (2025). Effects of bariatric surgery on the pharmacokinetics of cardiovascular drugs: A systematic review and meta-analysis. **Obesity Surgery**. <https://doi.org/10.1007/s11695-025-08419-x>

Zhou, J., Wang, F., & Li, S. (2025). The efficacy and safety of danuglipron and orforglipron in patients with type 2 diabetes and obesity: A systematic review and meta-analysis. **Frontiers in Endocrinology**. <https://doi.org/10.3389/fendo.2025.1646956>

Disclaimer: This response is based on current medical literature (as of January 2026) and is intended for informational purposes only. Clinical decisions should be made in consultation with qualified healthcare professionals and should consider individual patient characteristics, comorbidities, medication access, and patient preferences.

Disclaimer

This response is based on current medical literature (as of January 2026) and is intended for informational purposes only. Clinical decisions should be made in consultation with qualified healthcare professionals and should consider individual patient characteristics, comorbidities, medication access, and patient preferences.