

# Optical Properties of Gain incorporating Photonic Resonators



*by*  
**AHMAD BILAL**  
CIIT/FA15-BPH-019/ISB

BS Thesis  
In  
Physics

**COMSATS University Islamabad**  
Islamabad - Pakistan

January, 2019



**COMSATS University Islamabad**

Optical Properties of Gain incorporating Photonic  
Resonators

A Thesis Presented to

**COMSATS University Islamabad**

In partial fulfillment

of the requirement for the degree of

**Bachelor of Science in Physics**

by

Ahmad Bilal  
CUI/FA15-BPH-019/ISB

Spring, 2019

# Optical Properties of Gain incorporating Photonic Resonators

---

A Under Graduate Thesis submitted to the Department of Physics as partial fulfillment of the requirement for the award of Degree of BS (Physics).

Name	Registration Number
Ahmad Bilal	CUI/FA15-BPH-019/ISB

**Supervisor:**

Dr. Ahmer Naweed,  
Associate Professor,  
Department of Physics,  
COMSATS University Islamabad (CUI),  
January, 2019.

# Final Approval

---

This thesis titled

## Optical Properties of Gain incorporating Photonic Resonators

By

ar *Ahmad Bilal*

*CIIT/FA15-BPH-019/ISB*

Has been approved

For the COMSATS University Islamabad

External Examiner: \_\_\_\_\_

Supervisor: \_\_\_\_\_

Dr. Ahmer Naweed

Associate Professor, Dept. of Physics

COMSATS University Islamabad

HoD: \_\_\_\_\_

Dr. Sajid Qamar

Professor, Dept. of Physics

COMSATS University Islamabad

## Declaration

I **Ahmad Bilal** (CIIT/FA15-BPH-019/ISB) hereby declare that this project neither as a whole nor as a part there of has been copied out from any source. It is further declared that I have developed this thesis and the accompanied report entirely on the basis of my personal efforts made under the sincere guidance of my supervisors. No portion of the work presented in this report has been submitted in support of any other degree of qualification of this or any other University or Institute of learning, if found I shall stand responsible.

Date:

---

Ahmad Bilal  
CIIT/FA15-BPH-019/ISB

## Certificate

It is certified that Ahmad Bilal (Registration No. CIIT/FA156-BPH-019/ISB) has carried out all the work related to this thesis under my supervision at the Department of Physics, COMSATS University Islamabad and the work fulfills the requirement for award of BS degree.

Date: Jan, 2019

Supervisor:

---

Dr. Ahmer Naweel  
Associate Professor, Department of Physics

Head of Department:

---

Dr. Sajid Qamar  
Department of Physics

## **Dedication**

This thesis is dedicated to my mother who brought me up all by herself, motivated me to always persue my dreams and made me the gentleman I am today.

# Abstract

Since long, electronic integrated circuits have dominated our modern technology. Now with the dawn of photonics, which is basically using integrated circuits made up using optics, its not far that our modern technology takes a new toll and slide into a new generation of digital devices. Basically, Photonics is the technology of generating and harnessing light, and other form of radiant energy whose quantum unit is a photon. These can be used in multiple applications , to explore the vastness of the Universe, cure serious and unknown diseases and even help in forensics to solve difficult crime cases.

In this project, we extended the research on optical ring resonators for such mediums in which there is gain. First we studied normally the optical properties of passive resonators and measured the effects of EIT and EIA in them (details later discussed). Then we moved over focus on active resonators varrying different parameters to acheive EIT and EIA in gain incorporating photonic resonators which have extensive amount of applications. The main focus for this project was to model the characteristics and properties of active resonators and compare it with the results of passive resonators. Due to the gain property of active resonators, similar effects can be seen here as in passive resonators but without losses involved. The main idea was to establish a photonic device that could work efficiently as passive resonators and also have more output.

Light is a beautiful thing, it makes us see things, but what if it also starts to help us organize our data, compute our equations, play our music, record our documents



and basically do everything what a modern digital device, built on electronics, is capable of doing? I guess we'll find out!

*Indeed, in the creation of the heavens,  
and the earth and the alternation of  
the night and the day, are signs for  
those of understanding.*

*The Noble Quran [3:190]*

---

## Acknowledgement

---

In the name of Allah, who is the most beneficent and merciful. I would start off this extensive documentation with a quote from Carl Sagan, one of the greatest science educator, who created enough enthusiasm and curiosity in me to pursue my career in Physics. He said, "*Somewhere, something incredible is waiting to be known*". This is one of the reason I chose to be a student of physics, it inspires me to search for the unknown clues that are hidden in the very fabric of reality. Physics gave mankind the power to dominate their world and use the best of nature for their benefit.

Since childhood, I had always been fascinated by computers and gadgets. Having the background of engineers in my family, I almost ended up joining the computer engineering in High School. But the curiosity inside me had made me a star gazer. So I had questions about how they get where they are, and what are they made of? These questions were those which made me switch my field to Physics which is a science of never-ending curiosity. In this process, a lot of people are included some directly and some indirectly, most of which is my family, because their never-ending support had made me pursue my dreams.

I would personally like to thank my supervisor in BS project, Dr. A. Naweed, who helped me through thick and thin to complete this project and also kept me motivated enough to continue my research in field of photonics. I would like to

thank my batch counselor Dr. A. H. Mujtaba, who's support and teachings made us all work harder and harder for the progression of science. Also there is a big role of Ms. Zarqa in my motivation for this project. She not only recommended me to Dr. Naweed, but she is also my mental health counselor when I am in dire need of help. I would like to thank my peers in this, because the support and love I get from them is unmeasureable. Then again I would like to thank my family and especially my mother, who never asked me about my GPA and anything and always said, "if you love what you are studying, only then you can get true learning."

In the end, it is important to know that Knowledge is a never ending process, and Physics is such a beautiful field that every time I learn a new concept about the universe, it feels like I have been born again.

---

**Ahmad B. Yousafzai**

**Islamabad, Jan 2019**

# Contents

<b>Dedication</b>	<b>vii</b>
<b>Abstract</b>	<b>viii</b>
<b>Acknowledgement</b>	<b>x</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Resonators . . . . .	2
1.1.1 Explanation . . . . .	2
1.2 Optical Resonators . . . . .	3
1.3 Different Types of Optical Resonators . . . . .	4
1.3.1 Fabry-Perot Resonator . . . . .	4
1.3.2 Gires-Tournois . . . . .	4
1.4 Micro Resonators . . . . .	4
1.4.1 Different Geometries . . . . .	4
1.5 EIT and EIA . . . . .	5
1.6 Aim and Objective . . . . .	6
References . . . . .	7
<b>2 Area of Study</b>	<b>8</b>
2.1 The Fabry-Perot Interferometer . . . . .	8
2.1.1 Theory of Fabry-Perot interferometer . . . . .	9
2.1.2 Finesse, Q-factor . . . . .	11
2.2 Ring Geometry Resonators . . . . .	11
2.2.1 All-Pass Ringresonator . . . . .	11
2.2.2 Add-Drop Ringresonator . . . . .	12
2.2.3 Coupled Ringresonator . . . . .	13
2.3 Gain incorporation in Resonators . . . . .	14
2.3.1 Beer's Law . . . . .	14
2.3.2 Beer's law study as gain . . . . .	14
2.4 Gain medium . . . . .	14
<b>3 Coupled Resonators with Gain</b>	<b>16</b>
3.1 Coupled resonator with Gain medium . . . . .	16
3.1.1 Gain element . . . . .	16
3.2 Calculation/Equations . . . . .	16
3.2.1 For single . . . . .	17

3.2.2	For coupled . . . . .	17
3.2.3	For triple . . . . .	17
3.3	Coupling Regimes . . . . .	18
<b>4</b>	<b>Electromagnetically Induced Transparency</b>	<b>19</b>
4.1	EIT in Atoms . . . . .	19
4.1.1	Two level Atoms . . . . .	19
4.2	EIT in ring resonators . . . . .	19
4.3	EIT in Coupled resonators(CRIT) . . . . .	20
4.4	CRIT with gain . . . . .	20
4.5	Results . . . . .	20
<b>5</b>	<b>Electromagnetically Induced Absorbption</b>	<b>22</b>
5.1	EIA concepts . . . . .	22
5.1.1	EIA in atoms . . . . .	22
5.1.2	EIA Quantum phenomena . . . . .	22
5.2	EIA in resonators . . . . .	23
5.2.1	Coupled resontors induced Absorption . . . . .	23
5.3	CRIA with gain . . . . .	23
<b>6</b>	<b>Conclusion</b>	<b>25</b>
<b>A</b>	<b>Abrevations</b>	<b>27</b>

# List of Figures

1.1	Illustration of a basic optical cavity . . . . .	3
1.2	Different geometries of microresonators. . . . .	5
2.1	Illustrated energy diagram of a simple Fabry- Perot resonator . . . . .	8
2.2	Transmitted and reflected field of a simple Fabry- Perot resonator . . . . .	10
2.3	Illustrated fields of an all pass resonator . . . . .	12
2.4	Illustrated fields of an add drop resonator . . . . .	12
2.5	Illustrated fields and geometry of a coupled ring resonator . . . . .	13

# Chapter 1

## Introduction

Decades have passed since electric circuits become integrated on microchips or ICs. Technology is passing with time and now it is the time for photonics to get integrated into circuits. These can lead to a revolution in the digital technology we know today, with every hand-held device to corporate machines, all running on circuits made using photonic crystals and microresonators.

These topics require a detailed study, which is what we are going to do in this Thesis. The research of this Thesis is centered towards devices such as microring resonators (I will discuss them later), which includes gain medium.

This documentation is divided into different sections, compiling the work of 1 year long BS project. First, we will increase the understanding of the reader of what resonators, optical resonators, and microring resonators are. Then the study will focus on the systems that we used and their explanations, after that I will show the results of what I have collected by modeling the system in different conditions (parameters). This extensive documentation will be useful for anyone trying to get started in this field of research because I have written it in a fashion that a newbie in the field of photonics can easily grasp the ideas and can learn from it.

## 1.1 Resonators

A resonator is a device that exhibits resonant behavior naturally (or artificially) on some resonant frequencies, that is, it oscillates at those frequencies with higher amplitudes than others. These frequencies are called its resonant frequencies. These oscillations can either be electromagnetic waves or mechanical waves as well. There are different uses of resonators, they can be used to filter some specific frequencies or can also be used to generate a specific frequency of the wave. A resonator in which the waves exist in hollow space is called a cavity resonator, which is used in electronics and radio signal processing, known as microwave cavities, to generate, transmit and receive electromagnetic signals. Acoustic cavity resonators, in which sound is produced by air vibrating in a cavity with one opening, are known as Helmholtz resonators.

### 1.1.1 Explanation

The term resonator is most often used for a homogeneous object in which vibrations travel as waves, at an approximately constant velocity, bouncing back and forth between the sides of the resonator. The material of the resonator, through which the waves flow, can be viewed as being made of millions of coupled moving parts (such as atoms). Therefore, they can have millions of resonant frequencies, although only a few may be used in practical resonators. The oppositely moving waves interfere with each other, and at its resonant frequencies reinforce each other to create a pattern of standing waves in the resonator. If the distance between the sides is  $d$ , the length of a round trip is  $2d$ . To cause resonance, the phase of a sinusoidal wave after a round trip must be equal to the initial phase so the waves self-reinforce. The condition for resonance in a resonator is that the round trip distance,  $2d$ , is equal to an integer number of wavelengths  $\lambda$  of the wave:

$$2d = N\lambda, \quad N \in \{1, 2, 3, \dots\}$$

If the velocity of a wave is  $c$ , the frequency is  $f = c/\lambda$  so the



resonant frequencies are:

$$f = \frac{Nc}{2d} \quad N \in \{1, 2, 3, \dots\}$$

So the resonant frequencies of resonators, called normal modes, are equally spaced multiples (harmonics) of a lowest frequency called the fundamental frequency. The above analysis assumes the medium inside the resonator is homogeneous, so the waves travel at a constant speed, and that the shape of the resonator is rectilinear. If the resonator is inhomogeneous or has a nonrectilinear shape, like a circular drumhead or a cylindrical microwave cavity, the resonant frequencies may not occur at equally spaced multiples of the fundamental frequency. They are then called overtones instead of harmonics. There may be several such series of resonant frequencies in a single resonator, corresponding to different modes of vibration.

## 1.2 Optical Resonators

An optical resonator, also known as optical cavity, is usually composed of two highly reflecting mirrors held in front of each other parallelly inside a vacuum so that the system exhibits resonant behavior which allows standing wave modes to exist with almost no loss. Thus optical resonator is a cavity with walls that are highly reflected for electromagnetic waves (i-e light).




Figure 1.1: Illustration of a basic optical cavity

## 1.3 Different Types of Optical Resonators

### 1.3.1 Fabry-Perot Resonator

A system of two mirrors held parallel to each other and both having high reflectivities shows a resonant behavior at some frequencies of incident light. If both the mirrors have high reflectance, the incident light is still observed to have pass through them without any decrease in the intensity and is detected, which occurs due to phenoemnonas similar to quantum tunneling effects.

### 1.3.2 Gires-Tournois

It is basically a lossless Fabry-Perot resonator which have a 100% reflecting rear mirror, that means it reflects 100% at all frequencies. Still, some resonant frequencies stays between the mirrors for a longer period of time and thus descript resonant behavior and lead to ultra slow group velocities. This simple device is known for storing spectral power of light which is reflected from it while modifying its phase. That is why it is sometimes referred to as a "phase only" filter.

## 1.4 Micro Resonators

Microresonators are special type of resonators made from different type of materials which exhibits optical properties while being fabricated on a chip. These kind of resonators are actually useful in observing the effects of optical resonators on a device.

### 1.4.1 Different Geometeries

There are many type of microresonators from which microring-resonators are very useful in making photonic devices and have wide variety of application. Other kind of resonators are also useful for different kind of applications and all have distinct optical properties based on their geometry. (See figure 1.2)




Figure 1.2: Different geometries of microresonators.

## 1.5 EIT and EIA

Electromagnetically Induced Transparency (EIT), is a coherent optical nonlinearity which makes a medium transparent to some narrow bandwidth of frequencies which were otherwise opaque to the incident radiation. This window leads to slow light at resonant frequencies in an optical resonant system usually involving coupled system. This is observed due to the destructive quantum interference effects of the incident radiation in atomic levels.

Similarly, Electromagnetically Induced Absorption (EIA), is a similar phenomenon to EIT but in this nonlinearity the medium becomes highly opaque to some bandwidth of frequencies at resonance. Thus blocking off completely the resonant frequency radiation and causing a dip in the transmitted field. The quantum interference of light here is destructive and the atomic levels absorb the extra photons at such particular frequencies.

## **1.6 Aim and Objective**

This thesis is a detailed study of such phenomenons dealing optical resonators. We will also deeply study the changing behavior of active and passive resonators. Active resonators are those resonators which are made from some gain medium and they also descript EIT and EIA like behavior in similar and distinct fashion. Then we will model the systems using different scientific tools and computation methods to predict their behavior in different circumstances and parameters.

## References

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis.

# Chapter 2

## Area of Study

### 2.1 The Fabry-Perot Interferometer

Optical resonators were utilized as helpful gadgets as early as 1899, when Fabry and Perot depicted the utilization of a parallel-plate resonator as a multipass interferometer. Part of the incident light on this Fabry– Perot resonator is transmitted and another part is reflected, with power divisions that rely upon numerous factors. A simple illustration of the basic Fabry-Perot is shown in Figure 2.1, here  $r_1 t_1$  are the reflectivity constant and transmittivity constant of the mirror 1 respectively and  $r_2 t_2$  are the reflectivity and transmittivity constants of the mirror two respectively. Also,  $E_i$  is the incident Electromagnetic energy,  $E_t$  is the transmitted energy and  $E_r$  is the reflected energy. This is an asymmetric Fabry-Perot resonator:




Figure 2.1: Illustrated energy diagram of a simple Fabry-Perot resonator

### 2.1.1 Theory of Fabry-Perot interferometer

If the incident energy is in the form of white coherent light then at that point the transmission and reflection coefficients depend just on the mirror reflectivities. The total reflected power comprises of the power reflected from the principal mirror in addition to all the different reflections between the mirrors that add to the reflectivity in general. In summation, the equations are:

$$\mathcal{R} = R_1 + T_1^2 R_2 \sum_{m=1}^{\infty} (R_1 R_2)^{m-1} = \frac{R_1 - 2R_1 R_2 + R_2}{1 - R_1 R_2} \xrightarrow{R_1=R_2 \equiv R} \frac{2R}{1 + R} \quad (2.1)$$

Similarly, the transmitted energy in summation is:

$$\mathcal{T} = T_1 T_2 \sum_{m=1}^{\infty} (R_1 R_2)^{m-1} = \frac{T_1 T_2}{1 - R_1 R_2} \xrightarrow{R_1=R_2 \equiv R} \frac{T^2}{1 - R^2} = \frac{1 - R}{1 + R} \quad (2.2)$$

Assuming, be that as it may, the incident light comprises of a transiently lucid (monochromatic) plane wave, at that point the reflected power will be relative to the square of the reasonable total of every reflected field. Since the fields convey phase information with amplitudes added, the division of reflected and transmitted light depends not just on the mirror reflectivities, but in addition on the mirror separation and excitation wavelength. The rational

total of fields is amplified when every one of the fields interfere constructively (in phase) and limited when they interfere destructively (out of phase).

Phase gathers with propagation separation as  $\phi(z) = \beta z$  and may likewise be gained upon communication with the mirrors. The sound forms of

Eqs. 2.1 and 2.2 incorporate an aggregated stage factor for each round-trip that can be translated as a standardized detuning  $\phi = T_R \omega$ , where  $T_R$  is the cavity travel time,  $T_R = n_{eff} L / c$  for the circumference,  $L$  and effective index  $n_{eff}$ . Presently,  $\tilde{r}$  speaks to the complex reflectivity:

$$\begin{aligned}\tilde{r} &= r_1 - t_1^2 r_2 \exp(im\phi) \sum_{m=1}^{\infty} (r_1 r_2 \exp(im\phi))^{m-1} \\ &= \frac{r_1 - r_2 \exp(i\phi)}{1 - r_1 r_2 \exp(i\phi)} \xrightarrow{r_1=r_2 \equiv \tilde{r}} \frac{r(1 - \exp(+i\phi))}{1 - r^2 \exp(+i\phi)}\end{aligned}\quad (2.3)$$

and  $\tilde{t}$  represents the complex transmittivity:

$$\begin{aligned}\tilde{t} &= -t_1 t_2 \exp(im\phi/2) \sum_{m=1}^{\infty} (r_1 r_2 \exp(im\phi))^{m-1} \\ &= \frac{-t_1 t_2 \exp(im\phi/2)}{1 - r_1 r_2} \xrightarrow{r_1=r_2 \equiv \tilde{r}} \frac{-(1 - r^2) \exp(im\phi/2)}{1 - r^2}\end{aligned}\quad (2.4)$$

The square modulus of these perplexing amounts gives the reflection  $\mathcal{R}$  and transmission  $\mathcal{T}$  coefficients (shown in Fig. 2.2). Antiresonant wavelengths are more emphatically reflected than in the ambiguous case, while thunderous wavelengths are transmitted 100% for adjusted reflectors ( $r_1 = r_2$ ). For a fixed reflect dispersing, the transmission and reflection spectra in this manner show intermittent pinnacles and valleys. Figure 3.1 presenting the transmission and reflection spectra for a lossless, adjusted Fabry–Perot resonator. The part of reflected and transmitted power for mixed up excitation is identical to the separate frightfully arrived at the midpoint of reflection and transmission over a time of the spectrum range.




Figure 2.2: Transmitted and reflected field of a simple Fabry-Perot resonator



### 2.1.2 Finesse, Q-factor

The resonance condition is fulfilled when the (compelling) circumference of the ring, or for the most part the round-trip length, is equivalent to a whole number numerous of the optical wavelength inside the medium. This means a progression of Lorentzian-molded transmission bends equally dispersed in recurrence by the FSR (Free Spectral Range), with the resonance linewidth portraying the capacity time of photons inside the cavity. The photon lifetime can be standardized to one optical cycle, known as the quality factor ( $\mathcal{Q}$ ), or the cavity round-trip time, known as the cavity Finesse ( $\mathcal{F}$ ). The most extreme reachable Q-factor is characterized as  $\mathcal{Q}_{int}$ , which is the intrinsic loss of the cavity. At the point when the resonator is coupled to the outer world, the Q-factor further decreases because of the loss imported by the coupler ( $\mathcal{Q}_{ext}$ ). Thus the last quality factor  $\mathcal{Q}_{load}$  is comprised of these two parts:  $\mathcal{Q}_{load}^{-1} = \mathcal{Q}_{int}^{-1} + \mathcal{Q}_{ext}^{-1}$ .

## 2.2 Ring Geometry Resonators

In this section, I will discuss different kinds of ring shaped resonators whose principle is pretty much similar to the Fabry-Perot resonator and are more simple to make. Basically, a ring resonator is a simple waveguide which is turned in the shape of a ring. This allows it to exhibit resonant behavior on very specific frequencies. The light is coupled inside the ring due to the phenomenon of total internal reflection and interference. This kind of behaviour is noticed in all kind of classical waves, such as sound waves, which was observed inside a large cathedral's halls, thus it was named whispering galleries. Also, these resonators can be made using different material but in this thesis, we used semi-conductor silicon as the primary material.

### 2.2.1 All-Pass Ringresonator

A straightforward ring resonator is made by taking one yield of a conventional directional coupler and bolstering it once again into one input. Such a device displays periodic cavity resonance (reverberation) when light navigating the ring procures a phase move relating to a

number numerous of  $2\pi$  radians. The resonator is numerically defined from two parts: a coupling quality and an input way. In opposition to the limitless entirety inferences performed before for the Fabry– Perot and Gires– Tournois, in which we expected steadystate task and coordinating fields and derived basic spectral properties. Although, both strategies are similarly substantial, the field-coordinating technique has the benefit of simplicity.




Figure 2.3: Illustrated fields of an all pass resonator

### 2.2.2 Add-Drop Ringresonator

The immediate waveguide similarity of a free-space Fabry– Perot is gotten by including a second guide that side-couples to the resonator as in Fig. 1.4. Since this setup acts as a tight band abundance channel that can include or drop a recurrence band from an approaching sign, it is regularly named an add– drop filter.




Figure 2.4: Illustrated fields of an add drop resonator

### 2.2.3 Coupled Ringresonator

A simple case as an assymetric Fabry-Perot resonator, a coupled ring resonator has another ring above the first ring of the all pass resonator. This arrangement shows coupling between the two resonators (rings) and show different behaviour.




Figure 2.5: Illustrated fields and geometry of a coupled ring resonator

## 2.3 Gain incorporation in Resonators

Light, when travels through a medium, loses its intensity exponentially. This law is called the *Beer's law* for electromagnetic intensity. But some mediums, whose refractive index is such as they oppose the exponential decay of the light and rather increase the intensity in the propagation through the medium, are called gain medium. We can use these gain mediums and build a microresonators from them to observe different quantum optical phenomenons. First I will explain a bit how gain work.

### 2.3.1 Beer's Law

binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 2.3.2 Beer's law study as gain

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

## 2.4 Gain medium

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-

precision binary floating-point arithmetic and some concepts from numerical analysis.

## Chapter 3

# Coupled Resonators with Gain

### 3.1 Coupled resontaor with Gain medium

Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

#### 3.1.1 Gain element

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 3.2 Calculation/Equations

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from nu-

merical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 3.2.1 For single

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 3.2.2 For coupled

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 3.2.3 For triple

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis.

### 3.3 Coupling Regimes

To perform more advanced calculations, it is important to have some understanding of how mpmath works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some understanding of how mpmath works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis.



# Chapter 4

## Electromagnetically Induced Transparency

### 4.1 EIT in Atoms

Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

#### 4.1.1 Two level Atoms

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 4.2 EIT in ring resonators

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-

precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 4.3 EIT in Coupled resonators(CRIT)

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 4.4 CRIT with gain

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

### 4.5 Results

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-

precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis.

# Chapter 5

## Electromagnetically Induced Absorption

### 5.1 EIA concepts

To perform more advanced calculations, it is important to have some understanding of how mpmath works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some

#### 5.1.1 EIA in atoms

understanding of how mpmath works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some understanding of how mpmath works internally and what the possible sources of error are.

#### 5.1.2 EIA Quantum phenomena

This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some understanding of how mpmath works internally and what the possible sources of error are. This section gives an overview of arbitrary-

precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are.

## 5.2 EIA in resonators

This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis.

### 5.2.1 Coupled resonators induced Absorption

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis.

## 5.3 CRIA with gain

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from nu-

merical analysis. To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis.

# Chapter 6

## Conclusion

Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the

possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis. Most of the time, using `mpmath` is simply a matter of setting the desired precision and entering a formula. For verification purposes, a quite (but not always!) reliable technique is to calculate the same thing a second time at a higher precision and verifying that the results agree.

To perform more advanced calculations, it is important to have some understanding of how `mpmath` works internally and what the possible sources of error are. This section gives an overview of arbitrary-precision binary floating-point arithmetic and some concepts from numerical analysis.



# Appendix A

## Abbreviations

EIT Electromagnetically Induced Transparency

EIA Electromagnetically Induced Absorption

CRIT Coupled Resonator Induced Transparency

CRIA Coupled Resonator Induced Absorption

FSR Free Spectral Range

MRR Micro Ring Resonator

MZI Mach Zehnder Interferometer

FWHM Full width at half maximum

CMT Coupled Mode Theory