



Faculty of Engineering, Architecture and Science

Department of Electrical and Computer Engineering

Course Number	ELE532
Course Title	Signals and Systems I
Semester/Year	Fall 2020

Instructor	Javad Alirezaie
------------	------------------------

ASSIGNMENT No.	2
-----------------------	----------

Assignment Title	<i>System Properties and Convolution</i>
------------------	--

Submission Date	October 25th, 2020
Due Date	October 25th, 2020

Student Name	Nathan Agnew
Student ID	500819685
Signature*	<i>Nathan Agnew</i>
Student Name	Ahmad Fahmy
Student ID	500913092
Signature*	AF

**By signing above you attest that you have contributed to this written lab report and confirm that all work you have contributed to this lab report is your own work. Any suspicion of copying or plagiarism in this work will result in an investigation of Academic Misconduct and may result in a "0" on the work, an "F" in the course, or possibly more severe penalties, as well as a Disciplinary Notice on your academic record under the Student Code of Academic Conduct, which can be found online at: www.ryerson.ca/senate/current/pol60.pdf.*

Table of Contents

A. Impulse response	3
B. Convolution	5
C. System behavior and stability	9
D. Discussion	11

A Impulse response

Problem A.1 [0.5 Marks] Complete *Lathi, Section 2.7-1 Script Files*, page 213. Use MATLAB command **poly** to generate the characteristic polynomial from the characteristic values specified by **lambda**.

```
% Problem A.1


% CH2MP1.m : Chapter 2, MATLAB Program 1
% Script M-file determines characteristic roots of op-amp circuit.
% Set component values:
R = [1e4, 1e4, 1e4];
C = [1e-6, 1e-6];
% Determine coefficients for characteristic equation:
A = [1, (1/R(1)+1/R(2)+1/R(3))/C(2), 1/(R(1)*R(2)*C(1)*C(2))];
% Determine characteristic roots:
lambda = roots(A)
eq1 = poly(lambda);

lambda =

-261.8034
-38.1966
```

Problem A.2 [2 Marks] Plot the impulse response of the system in Problem A.1 for $t = [0:0.0005:0.1]$.

```
AssignmentB.m AssignmentA.m CH2MP1.m CH2MP2.m lab2conv.m +
1 %Assignment A.2
2 lab2conv;
3 CH2MP1;
4
5 f = @(t) poly(lambda);
6 t = (0:0.0005:0.1);
7
8 %plot results
9 figure('Name', 'Impulse Response of Lambda','NumberTitle','off');
10 plot(t, polyval(f(t), t));
11 xlabel('t'); ylabel('f(t)'); grid;
12
```



Problem A.3 [0.5 Marks] Complete *Lathi, Section 2.7-2 Function M-Files*, page 214.

```


lab2conv.m  CH2MP1.m  CH2MP2.m  AssignmentA.m  AssignmentB.m  ProblemA.m
1  function [lambda] = CH2MP2(R,C)
2  % CH2MP2.m : Chapter 2, MATLAB Program 2
3  % Function M-file finds characteristic roots of op-amp circuit.
4  % INPUTS: R = length-3 vector of resistances
5  % C = length-2 vector of capacitances
6  % OUTPUTS: lambda = characteristic roots
7  % Determine coefficients for characteristic equation:
8  A = [1, (1/R(1)+1/R(2)+1/R(3))/C(2), 1/(R(1)*R(2)*C(1)*C(2))];
9  % Determine characteristic roots:
10 lambda = roots(A);

```


B. Convolution

Problem B.1 [3 Marks] Exercise M2.4 Lathi, *Matlab Session 2*, page 232. Plot $y(t)$ at step $t = 2.25$ as shown in Figure M2.4 on page 233. Use the MATLAB command **pause** instead of **drawnow** to observe the steps of the convolution operation slowly.

```
% CH2MP4.m : Chapter 2, MATLAB Program 4
% Script M-file graphically demonstrates the convolution process.
figure(1) % Create figure window and make visible on screen
u = @(t) 1.0*(t>=0);
x = @(t) 1.5*sin(pi*t).*(u(t)-u(t-1));
h = @(t) 1.5*(u(t)-u(t-1.5))-u(t-2)+u(t-2.5);
dtau = 0.005;
tau = -1:dtau:4;
ti = 0;
tvec = -.25:.1:3.75;
y = NaN*zeros(1,length(tvec));
% Pre-allocate memory
for t = tvec
    ti = ti+1; % Time index
    xh = x(t-tau).*h(tau);
    lxh = length(xh);
    y(ti) = sum(xh.*dtau); % Trapezoidal approximation of convolution integral
    subplot(2,1,1), plot(tau,h(tau), 'k-', tau,x(t-tau), 'k--', t,0, 'ok');
    axis([tau(1) tau(end) -2.0 2.5]);
    patch([tau(1:end-1);tau(1:end-1);tau(2:end);tau(2:end)],...
        [zeros(1,lxh-1);xh(1:end-1);xh(2:end);zeros(1,lxh-1)],...
        [.8 .8 .8], 'edgecolor', 'none');
    xlabel('\tau');
    title('B.1 h(\tau) [solid], x(t-\tau) [dashed], h(\tau)x(t-\tau) [gray]');
    c = get(gca, 'children'); set(gca, 'children', [c(2);c(3);c(4);c(1)]);
    subplot(2,1,2), plot(tvec,y, 'k', tvec(ti), y(ti), 'ok');
    xlabel('t'); ylabel('y(t) = \int h(\tau)x(t-\tau) d\tau');
    axis([tau(1) tau(end) -1.0 2.0]); grid;
    %drawnow;
    pause;
end
%pause;
```



Problem B.2 [4 Marks] Perform the convolution of the signal $x(t)$ in **Figure P2.4-19(a)** with $h(t)$ in **Figure P2.4-21**. Plot all signals and results.



```
clear;
t1=-10;
t2=10;
N=2000;
Delta_t=(t2-t1)/N;
t = (t1:Delta_t:t2);

x = zeros(size(t));
x(t>=-1 & t<=1)=1;

f = zeros(size(t));
f(t>=-1 & t<=0)=1;

for i=1:length(t)
    if (t(i) < -1 || t(i) > 0)
        h(i)=0;
    else
        h(i)=h(i-1)+.01;
    end
end

x1=x*Delta_t;

y=conv(x1,h);


figure('Name','Figure B.2','NumberTitle','off');

subplot(3,1,1);
plot(t,x); axis([t1 t2 -0.1 1.1])
title('x(t)');xlabel('t');

subplot (3,1,2);
plot(t,h); axis([t1 t2 -0.1 1.1])
title('h(t)');xlabel('t');

subplot (3,1,3);
plot(t,y(1000:3000)); axis([t1 t2 -0.1 1.1])
title('y(t)');xlabel('t');
```

Problem B.3 [5 Marks] Perform the convolution of the signal $x_1(t)$ and $x_2(t)$ in **Figure P2.4-18(a), (b) and (h)**. Plot all signals and results.



```
x_1 = zeros(size(t));
x_1(t>=4 & t<=6)=1;

x_2 = zeros(size(t));
x_2(t>=-5 & t<=-4)=1;
```

```
x_lb=x_1.*.1;
x_2b=x_2.*.1;
```

```
f=conv(x_lb,x_2b);
```

```
figure('Name','Figure B.3(a)','NumberTitle','off');
```

```
subplot(3,1,1);
plot(t,x_1); axis([t1 t2 -0.1 1.1])
title('x(t)');xlabel('t');
```

```
subplot (3,1,2);
plot(t,x_2); axis([t1 t2 -0.1 1.1])
title('h(t)');xlabel('t');
```

```
subplot (3,1,3);
plot(t,f(1000:3000));axis([t1 t2 -0.1 1.1])
title('h(t)');xlabel('t');
```

```
x_1 = zeros(size(t));
x_1(t>=3 & t<=5)=1;

x_2 = zeros(size(t));
x_2(t>=-5 & t<=-3)=1;
```

```
x_lb=x_1.*.1;
x_2b=x_2.*.1;
```


```
f=conv(x_lb,x_2b);
```

```
figure('Name','Figure B.3(b)','NumberTitle','off');
```

```
subplot(3,1,1);
plot(t,x_1); axis([t1 t2 -0.1 1.1])
title('x(t)');xlabel('t');
```

```
subplot (3,1,2);
plot(t,x_2); axis([t1 t2 -0.1 1.1])
title('h(t)');xlabel('t');
```

```
subplot (3,1,3);
plot(t,f(1000:3000));axis([t1 t2 -0.1 2.1])
title('h(t)');xlabel('t');
```



```

u = @(t) 1.0*(t>=0);
f_1 = @(t) exp(t).*(u(t+2)-u(t)); %x1
f_2 = @(t) exp(-2*t).*(u(t)-u(t-1)); %x2

f_1a=f_1(t);
f_2a=f_2(t);

h1=conv(f_1a,f_2a);

figure('Name','Figure B.3(h)','NumberTitle','off');

subplot(3,1,1);
plot(t,f_1a);
title('x_1(t)');xlabel('t');

subplot (3,1,2);
plot(t,f_2a);
title('x_2(t)');xlabel('t');

subplot (3,1,3);
plot(t,h1(1000:3000));
title('h(t)');xlabel('t');

```


C. System behavior and stability

Problem C.1 [4 Marks] Consider the LTI systems S1, S2, S3 and S4 represented by their respective unit impulse response functions given as follows:


$$h_1(t) = e^{\frac{t}{5}}u(t); \quad (3)$$

$$h_2(t) = 4e^{-\frac{t}{5}}u(t); \quad (4)$$

$$h_3(t) = 4e^{-t}u(t); \quad (5)$$

$$h_4(t) = 4(e^{-\frac{t}{5}} - e^{-t})u(t); \quad (6)$$


Plot each unit impulse response function for $t = [-1:0.001:5]$.



```
u = @(t) 1.0*(t>=0);
h_1 = @(t) exp(t/5).*u(t);
h_2 = @(t) 4*exp(-t/5).*u(t);
h_3 = @(t) 4*exp(-t).*u(t);
h_4 = @(t) 4*(exp(-t/5)-exp(-t)).*u(t);

t=[-1:0.001:5];

figure('Name','Figure C1 h_1','NumberTitle','off');
plot(t,h_1(t));
title('C.1 h_1(t)');
ylabel('h_1');
xlabel('t');
pause;
figure('Name','Figure C1 h_2','NumberTitle','off');
plot(t,h_2(t));
title('C.1 h_2(t)');
ylabel('h_2');
xlabel('t');
pause;
figure('Name','Figure C1 h_3','NumberTitle','off');
plot(t,h_3(t));
title('C.1 h_3(t)');
ylabel('h_3');
xlabel('t');
pause;
figure('Name','Figure C1 h_4','NumberTitle','off');
plot(t,h_4(t));
title('C.1 h_4(t)');
ylabel('h_4');
xlabel('t');
```



Problem C.2 [1 Mark] Determine the characteristic values (eigenvalues) of systems S1–S4.

$$\lambda h_1 = \frac{1}{5}$$

$$\lambda h_2 = -1/5$$


$$\lambda h_3 = -1$$

$$\lambda h_4 = -1/5, -1$$

Problem C.3 [5 Marks] Truncate the impulse response functions $h_1(t), \dots, h_4(t)$ such that they are nonzero only for $0 \leq t \leq 20$. Determine the convolution of the truncated impulse response functions with the input signal $x(t) = [u(t) - u(t-3)] \sin 5t$ using the M-file in **Problem B.1** with the following changes **tau** = [0:dttau:20] and **tvec** = [0:0.1:20]. Plot the output of each system. State and explain your observations. Is there any relationship between the outputs of systems S2, S3, and S4? Explain.

```
h2_1=@(t) h_1(t).*(u(t)-u(t-20));
h2_2=@(t) h_2(t).*(u(t)-u(t-20));
h2_3=@(t) h_3(t).*(u(t)-u(t-20));
h2_4=@(t) h_4(t).*(u(t)-u(t-20));

figure('Name','Figure C3 h_1','NumberTitle','off');
% Create figure window and make visible on screen
u = @(t) 1.0*(t>=0);
x = @(t) sin(5*t).*(u(t)-u(t-3));
h = @(t) h2_1(t);
dttau = 0.005;
tau = 0:dttau:20;
ti = 0;
tvec = 0:.1:20;
y = NaN*zeros(1,length(tvec));
% Pre-allocate memory
for t = tvec
    ti = ti+1; % Time index
    xh = x(t-tau).*h(tau);
    lxh = length(xh);
    y(ti) = sum(xh.*dttau); % Trapezoidal approximation of convolution integral
    subplot(2,1,1),plot(tau,h(tau),'k-',tau,x(t-tau),'k--',t,0,'ok');
    axis([tau(1) tau(end) -2.0 2.5]);
    patch([tau(1:end-1);tau(1:end-1);tau(2:end);tau(2:end)],...
        [zeros(1,lxh-1);xh(1:end-1);xh(2:end);zeros(1,lxh-1)],...
        [.8 .8 .8],'edgecolor','none');
    xlabel('\tau');
    title('C3(a) h(\tau) [solid], x(t-\tau) [dashed], h(\tau)x(t-\tau) [gray]');
    c = get(gca,'children'); set(gca,'children',[c(2);c(3);c(4);c(1)]);
    subplot(2,1,2),plot(tvec,y,'k',tvec(ti),y(ti),'ok');
    xlabel('t'); ylabel('y(t) = \int h(\tau)x(t-\tau) d\tau');
    axis([tau(1) tau(end) -1.0 5.0]); grid;
    drawnow;
    %pause;
end
pause;
```



After performing the convolution, you can see that h_2 , h_3 , and h_4 have similar graphs. Each one has a wave the length of $x(t)$ and then transitions to $h(t)$ which approaches 0.

D. Discussion

Problem D.1 [2 Marks] Calculate the results of **Problems A.1, A.2** and **A.3** by hand and compare to those obtained with your MATLAB code.

Problem D.2 [1 Mark] What can you say about the width/duration of the signal resulting from the convolution of two signals?

The Width/duration of the convolution of 2 signals is the addition of their individual widths/durations.