

Midterm I Review

Math HI85

Ahmed Shakil

Contents

1	Tools and Tricks	2
2	Imaginary Numbers and Complex Variables	3
2.1	Topology of \mathbb{C}	3
3	Complex Derivatives	4
4	Power Series	4
4.1	Differentiation of Power Series	5
4.2	Cauchy-Riemann Equations	6
5	Integrating Over Curves	7
5.1	Curves	7
5.1.1	Integration on Parametrized Curves	7
5.1.2	Cauchy's Theorem	8

I Tools and Tricks

I.1: Properties of the modulus

$$\begin{aligned} z\bar{z} &= |z|^2 \\ \overline{z \cdot w} &= \bar{z} \cdot \bar{w} \\ |z \cdot w| &= |z| \cdot |w| \end{aligned}$$

I.2: $z^n = \omega$

Let $\omega = re^{i\theta}$, with $r > 0$, and $n > 0$.

$$\begin{aligned} z^n &= \omega \\ z &= \sqrt[n]{r}e^{i(\frac{\theta+2\pi k}{n})}, k \in \{0, 1, \dots, n-1\} \end{aligned}$$

I.3: Standard nonholomorphic functions

The following functions are $\mathbb{C} \rightarrow \mathbb{C}$ and are not holomorphic at any point in the complex plane.

$$\begin{aligned} f_1(z) &= |z| \\ f_2(z) &= \bar{z} \end{aligned}$$

I.4: Expansion of e^x

$$e^z = \sum_{n \geq 0} \frac{z^n}{n!}$$

I.5: Some useful limits

$$\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$$

I.6: Bounding the sin function

$$\sin \theta \geq \frac{2\theta}{\pi}, \quad \theta \in \left[0, \frac{\pi}{2}\right]$$

Note to self: Look for a tighter bound. (Check my hw)

2 Imaginary Numbers and Complex Variables

We start with the concept of an imaginary number, we define $i := \sqrt{-1}$. A complex number is $z := a + bi$ where $a, b \in \mathbb{R}$. We regard \mathbb{C} as the algebraic closure of \mathbb{R} . We denote the "real part" of z as $\text{Re}(z)$, similarly we denote the "imaginary part" of z as $\text{Im}(z)$. We denote the complex conjugate as $\bar{z} := a - bi$, and the set of complex numbers as \mathbb{C} .

Operations on complex numbers are similar to operations on the real numbers. Addition is component wise, and multiplication distributes. Addition and multiplication satisfy the usual commutativity, associativity, and distributivity rules. Division is a bit nuanced:

2.1: Division in \mathbb{C}

If $z, w \in \mathbb{C}$, and $w \neq 0$, then $\frac{z}{w} \in \mathbb{C}$ is the unique complex number such that $(\frac{z}{w}) \cdot w = z$.

Geometrically we view \mathbb{C} as a two-dimensional plane where the y axis is the imaginary axis and the x axis is the real axis. We denote the absolute value (modulus) of $z = a + bi \in \mathbb{C}$ as $|z| := \sqrt{a^2 + b^2}$. Addition of complex numbers can be seen as addition of vectors in two dimensions.

We can translate $z \in \mathbb{C}$ by describing them in polar form instead.

2.2: Polar form of complex numbers

$$\begin{aligned} & \text{Polar} \quad \text{Cartesian} \\ & (r, \theta) \rightarrow (x, y) \\ & (\sqrt{x^2 + y^2}, \arctan(\frac{y}{x})) \leftarrow (x, y) \\ & re^{i\theta} \rightarrow r \cos \theta + ir \sin \theta \\ & |z|e^{i \arctan(\frac{y}{x})} \leftarrow z = x + iy \end{aligned}$$

We view θ as the angle from the x axis in the counter-clockwise direction, and r as the straight line distance from the point in the complex plane to the origin.

Theorem 2.3: Euler's Theorem

$$e^{iz} = \cos(z) + i \sin(z)$$

Proof. Follows from Taylor series expansion formulas. □

2.1 Topology of \mathbb{C}

As a metric space we make a sort of equivalence between \mathbb{C} and \mathbb{R}^2 . In a sense,

$$\begin{aligned} \mathbb{C} &\simeq \mathbb{R}^2 \\ z = x + iy &\leftrightarrow (x, y) \\ |z| = \sqrt{x^2 + y^2} &\quad |(x, y)| = \sqrt{x^2 + y^2} \end{aligned}$$

Example 2.4: Neighborhoods in \mathbb{C}

Let $z_0 \in \mathbb{C}$ and $0 < r \in \mathbb{R}$.

$$\overline{B_r(z_0)} := \{z \in \mathbb{C} : |z - z_0| \leq r\}$$

is the closed ball of radius r around z_0 .

Similarly,

$$B_r(z_0) := \{z \in \mathbb{C} : |z - z_0| < r\}$$

is the open ball.

A subset $U \subset \mathbb{C}$ is open if $\forall z \in U, \exists 0 < r \in \mathbb{R}$ such that $B_r(z) \subset U$.

It is important to note that a sequence $\{z_n = x_n + iy_n\} \subset \mathbb{C}$ converges to $z \in \mathbb{C}$ if and only if

$$\lim_{n \rightarrow \infty} x_n = x \text{ and } \lim_{n \rightarrow \infty} y_n = y.$$

3 Complex Derivatives

Let U be an open subset of \mathbb{C} and $f : U \rightarrow \mathbb{C}$ be a function.

Definition 3.1: Holomorphic at a point

We say f is holomorphic at some point z_0 if $\lim_{h \rightarrow 0} \frac{f(z_0+h) - f(z_0)}{h}$ exists in \mathbb{C} . Equivalently it is holomorphic at z_0 if the following limit exists

$$\lim_{z_0 \rightarrow z} \frac{f(z) - f(z_0)}{z - z_0}.$$

If these limits exists, then we denote the value as $f'(z_0)$.

Remark 3.2: About holomorphic functions

Holomorphicity is a property which is quite a lot stronger than typically differentiability in the real number world. Holomorphicity means we demand the limit to be the same along any path of approach to a particular point. A common example of a function which is not holomorphic due to limits not being consistent along different paths of approach is the conjugation function: $f(z) = \bar{z}$. This function is not holomorphic at any point in the complex plane.

Some cool properties of holomorphic functions

- If a function is holomorphic at a point, then it is automatically infinitely complex differentiable at this particular point
- If f, g are holomorphic on a connected $U \subset \mathbb{C}$ and $f = g$ on a line segment, then $f = g$ on all of U

Something important to note about complex differentiation is that the typical differentiation rules from the real numbers hold. So things like the product, quotient, chain, and addition rules hold. Also, the rules for differentiating polynomials are the same as well. Ratio of polynomials are holomorphic at all points in the complex plane other than the points at which the denominator is equal to 0.

4 Power Series

A power series is an expression of the following form:

$$\sum_{n \geq 0} a_n z^n = a_0 + a_1 z + a_2 z^2 + \dots$$

Remark 4.1: Manipulating power series

Addition:

$$\left(\sum_{n \geq 0} a_n z^n\right) + \left(\sum_{n \geq 0} b_n z^n\right) = \sum_{n \geq 0} (a_n + b_n) z^n$$

Multiplication:

$$\left(\sum_{n \geq 0} a_n z^n\right) \left(\sum_{n \geq 0} b_n z^n\right) = \sum_{n \geq 0} \left(\sum_{j+k=n} a_j b_k\right) z^n$$

A power series defines a function when it converges.

Example 4.2: Geometric SeriesLet $a \in \mathbb{C}$

$$\sum_{n \geq 0} a^n z^n$$

is called the geometric series.

The convergence of this power series can be contemplated by taking a look at the partial sums. For example,

$$\sum_{n \geq 0}^{N-1} a^n z^n = \frac{1 - (az)^N}{1 - az}.$$

Now based on this we see that if $|az| < 1$ then the series converges, on the other hand if $|az| > 1$ then the series diverges. If we have that $|az| = 1$ then it is less clear as to if the series converges or diverges. If $az = 1$ then the series diverges and $(az)^n$ amounts to moving around the unit circle without converging. Ultimately, the geometric series converges if and only if $|z| < \frac{1}{a}$.

Recall that we say $\sum z_n$ converges absolutely if $\sum |z_n|$ converges. Also recall that absolute convergence implies convergence.

Definition 4.3: Radius of Convergence

For a power series $\sum_{n \geq 0} a_n z^n$, we define the radius of convergence as

$$r := \left(\limsup_{n \rightarrow \infty} |a_n|^{\frac{1}{n}}\right)^{-1}$$

Theorem 4.4: Convergence of power series

1. If $|z| < r$, then the power series converges absolutely.
2. If $|z| > r$, then the power series diverges.
3. If $|z| = r$, then more examination is required, the series could converge or diverge.

4.1 Differentiation of Power Series

For $z_0 \in \mathbb{C}$, a power series centered at z_0 is an expression

$$f(z) := \sum_{n \geq 0} a_n (z - z_0)^n$$

Once again, it's radius of convergence is given by $r := \left(\limsup_{n \rightarrow \infty} |a_n|^{\frac{1}{n}}\right)^{-1}$, and it converges absolutely for $|z - z_0| < r$.

Theorem 4.5

1. $f(z)$ is holomorphic at all $z \in B_r(z_0)$, and
2. $f'(z) = \sum_{n \geq 0} n a_n z^{n-1}$ is also holomorphic on $B_r(z_0)$ with the same radius of convergence.

Definition 4.6: Analytic function

A function $f : U \rightarrow \mathbb{C}$ on an open subset $U \subset \mathbb{C}$ is analytic if for every $z_0 \in U$, $\exists r > 0$ such that f agrees with an absolutely convergent power series on $B_r(z_0)$.

Examples of analytic functions are $\sin(z)$ and $\cos(z)$.

4.2 Cauchy-Riemann Equations

As we mentioned before \mathbb{C} is basically isomorphic to \mathbb{R}^2 when viewed as metric spaces. However, in terms of differentiability they are not the same. Now say we have some open $\Omega \subset \mathbb{C}$ and $f : \Omega \rightarrow \mathbb{C}$. We can write f as the following function:

$$\begin{aligned} f &= u + iv, \quad \text{where} \\ u &= \operatorname{Re}(f) : \Omega \rightarrow \mathbb{R} \\ v &= \operatorname{Im}(f) : \Omega \rightarrow \mathbb{R} \end{aligned}$$

Using this, given some $z = x + iy$, we view $u(z) = u(x, y)$ and the same goes for v as well.

Theorem 4.7: Cauchy-Riemann Equations

If f is holomorphic at z_0 , then

$$\begin{aligned} \frac{\partial u}{\partial x}(z_0) &= \frac{\partial v}{\partial y}(z_0) \\ \frac{\partial u}{\partial y}(z_0) &= -\frac{\partial v}{\partial x}(z_0) \end{aligned}$$

These are called the Cauchy-Riemann equations.

Proof.

$$\begin{aligned} \lim_{\substack{x \rightarrow 0 \\ x \in \mathbb{R}}} \frac{f(z_0 + x) - f(z_0)}{x} &= f'(z_0) = \lim_{\substack{y \rightarrow 0 \\ y \in \mathbb{R}}} \frac{f(z_0 + iy) - f(z_0)}{iy} \\ \frac{\partial f}{\partial x}(z_0) &= \frac{1}{i} \frac{\partial f}{\partial y}(z_0) \\ \frac{\partial u}{\partial x}(z_0) + i \frac{\partial v}{\partial x}(z_0) &= \frac{1}{i} \left(\frac{\partial u}{\partial y}(z_0) + i \frac{\partial v}{\partial y}(z_0) \right) \\ \implies \begin{cases} \frac{\partial u}{\partial x}(z_0) &= \frac{\partial v}{\partial y}(z_0) \\ \frac{\partial u}{\partial y}(z_0) &= -\frac{\partial v}{\partial x}(z_0) \end{cases} \end{aligned}$$

□

It is also important to note that there exists a converse result. If f is in C^1 , meaning $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$ all exist and are continuous, and the Cauchy-Riemann equations hold, then f is holomorphic.

The partial derivative matrix of a holomorphic function has the following special form:

$$\begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix} = \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$$

It is important to note that holomorphic functions are conformal mappings, meaning they infinitesimally preserve angles or scale them to 0.

5 Integrating Over Curves

Integrating in complex analysis in the most basic sense amounts to the following.

$$\int f(z) dz = \int \operatorname{Re}(f(z)) dz + \int \operatorname{Im}(f(z)) dz$$

5.1 Curves

The idea is to connect \int_γ to \int_a^b using substitution, where γ is a curve.

Definition 5.1: Parametrized Curve

A parametrized curve is a continuous function

$$\gamma(t) : [a, b] \subset \mathbb{R} \rightarrow \mathbb{C}$$

We say a curve γ is piecewise smooth if we can divide the domain $[a, b]$ into finitely many subintervals on which γ is smooth (i.e. infinitely differentiable).

5.1.1 Integration on Parametrized Curves

Theorem 5.2: U-substitution

If $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise-smooth, and f is defined on the image of γ , then

$$\int_\gamma f(z) dz = \int_a^b f(\gamma(t)) \gamma'(t) dt.$$

An important property to note is that reversing the orientation of the curve is equal to taking the negative of the original curve. The reverse of γ is defined as follows.

$$\gamma^-(t) = \gamma(b + a - t)$$

Then $-\int_{\gamma^-} f(z) dz = \int_\gamma f(z) dz$. As a matter of definitions it is important to note that we refer to an antiderivative as a primitive, these terms should be taken as one and the same.

Theorem 5.3: Fundamental Theorem of Calculus

$$\int_\gamma f(z) dz = F(\gamma(b)) - F(\gamma(a))$$

Corollary:

If $\gamma : [a, b] \rightarrow \mathbb{C}$ is a closed loop, i.e. $\gamma(a) = \gamma(b)$, then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)) = 0$$

Example 5.4: The most fundamental example

Say $f(z) = z^n$ and $n \in \mathbb{Z}$. Consider the curve $\gamma(t) = re^{it}$ where $r \in \mathbb{R}_{>0}$ and $t \in [0, 2\pi]$. Calculate $\int_{\gamma} f(z) dz$ for all $n \in \mathbb{Z}$.

$$\begin{aligned} \int_{\gamma} f(z) dz &= \int_0^{2\pi} r^n e^{nit} (rie^{it}) dt \\ &= ir^{n+1} \int_0^{2\pi} e^{(n+1)it} dt \end{aligned}$$

If $n \neq -1$: A primitive for $e^{(n+1)it}$ is $\frac{1}{(n+1)i} e^{(n+1)it}$.

$$\implies \boxed{\int_{\gamma} f(z) dz = 0} \quad (n \neq -1)$$

If $n = -1$:

$$\begin{aligned} ir^0 \int_0^{2\pi} e^{(0)it} dt &= 2\pi i \\ \implies \boxed{\int_{\gamma} f(z) dz &= 2\pi i} \quad (n = -1) \end{aligned}$$

Conclusion:

$$\int_{\partial B_r(0)} z^n dz = \begin{cases} 0 & n \neq -1 \\ 2\pi i & n = -1 \end{cases}$$

5.1.2 Cauchy's Theorem

A boundary of a subset $\Omega \subset \mathbb{C}$ is $\overline{\Omega} \setminus \Omega^\circ$. Basically the boundary is the points in the closure of a set which is not in the interior of the set.

An important thing to note about substitutions: If $g(z)$ is holomorphic on an open neighborhood of γ

$$\int_{\gamma} f(g(z)) dz = \int_{g \circ \gamma} f(z) g'(z) dz.$$

Theorem 5.5: Cauchy's Theorem

If $U \subset \mathbb{C}$ is an open set and has a piecewise-smooth boundary, and $f(z)$ is holomorphic on a domain containing \overline{U} , then

$$\int_{\partial U} f(z) dz = 0.$$

This is the most fundamental and remarkable tool in complex analysis.

Contour manipulation is an important technique to note. This is because to calculate an integral on a curve, we can deform that curve through a domain where our function is holomorphic in order to calculate our desired result.

Theorem 5.6: Cauchy's formula

Suppose we have $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$ which is defined on an open set as the domain, and is holomorphic. Let $z_0 \in \Omega$ and $r > 0$ such that $B_r(z_0) \subset \Omega$. Then for all $z \in B_r(z_0)$, we have

$$f(z) = \frac{1}{2\pi i} \int_{\partial B_r(z_0)} \frac{f(w)}{w - z} dw.$$

Proof. Take a delta ball around some hole $f(z)$ and use it to approximate $f(w)$, then show the error goes to zero. □

Corollary 5.7: Infinite differentiability

As a result of Cauchy's formula we have that wherever f is holomorphic it is infinitely differentiable. We can differentiate Cauchy's formula to derive the following more general formula for higher order derivatives:

$$f^{(n)}(z) = \frac{n!}{2\pi i} \int_{\partial B_r(z_0)} \frac{f(w)}{(w - z)^{n+1}} dw$$

Lemma 5.8: Jordan's Lemma

Let $f(z) = e^{iaz}g(z)$ where $z \in \mathbb{C}$ and $a > 0$.
Then

$$\left| \int_{\gamma} f(z) dz \right| \leq \frac{\pi}{a} \sup_{\gamma} |g(z)|.$$

Note that the curve is a semicircle with R as the radius.

Theorem 5.9: Goursat's Theorem

If $f(z)$ is holomorphic on a neighborhood containing Δ then

$$\int_{\Delta} f(z) dz = 0.$$

Theorem 5.10

If f is holomorphic on a ball $B_r(z_0)$ where $z_0 \in \mathbb{C}$ and $r > 0$, then f has a primitive on $B_r(z_0)$.

Definition 5.11: Homotopic Curves

Say we are given two parametrized curves (where $U \subset \mathbb{C}$ is an open set)

$$\gamma_0 : [a, b] \rightarrow U$$

$$\gamma_1 : [a, b] \rightarrow U$$

where

$$\gamma_0(a) = \gamma_1(a) = z_a$$

$$\gamma_0(b) = \gamma_1(b) = z_b.$$

We say γ_0 and γ_1 are homotopic in U if there exists a jointly continuous $\gamma_s(t)$ where $s \in [0, 1]$ and $t \in [a, b]$ such that

$$\gamma_s(a) = z_a \quad \forall s$$

$$\gamma_s(b) = z_b \quad \forall s$$

$$\gamma_s(t) \big|_{s=0} = \gamma_0(t)$$

$$\gamma_s(t) \big|_{s=1} = \gamma_1(t)$$

Definition 5.12: Simply Connected

We say U is simply connected if any two curves γ_0 and γ_1 in U with the same end points are homotopic in U .

Theorem 5.13: Equivalence of Homotopic Curves

If f is holomorphic on a open subset $U \subset \mathbb{C}$ where γ_0 and γ_1 are homotopic in U , then

$$\int_{\gamma_0} f(z) dz = \int_{\gamma_1} f(z) dz.$$

Theorem 5.14: Existence of Primitives

Any holomorphic function f on a simply connected domain $U \subset \mathbb{C}$ has a primitive F on U .