
Top-1 accuracy by INTERMED complexity: four buckets (v2, repetition 1)

4 questions: strategies_272_v2_rep1_4it.eval.json, 8 questions: strategies_272_v2_rep1_8it.eval.json, 16 questions: strategies_272_v2_rep1_16it.eval.json. Top 1: the true diagnosis's ICD-11 code is among the strategy's first 1 diagnoses (exact code match). Brackets: 95% Wilson interval.



Strategy	Run	0–11 54 cases	12–17 74 cases	18–23 73 cases	24–60 71 cases
1. Hypothesis-driven	4 questions	11.1% (5–22)	17.6% (11–28)	11.0% (6–20)	19.7% (12–30)
1. Hypothesis-driven	8 questions	20.4% (12–33)	27.0% (18–38)	21.9% (14–33)	21.1% (13–32)
1. Hypothesis-driven	16 questions	27.8% (18–41)	36.5% (26–48)	32.9% (23–44)	21.1% (13–32)
2. Knowledge-driven	4 questions	20.4% (12–33)	24.3% (16–35)	12.3% (7–22)	18.3% (11–29)

Strategy	Run	0–11 54 cases	12–17 74 cases	18–23 73 cases	24–60 71 cases
2. Knowledge-driven	8 questions	16.7% (9–29)	27.0% (18–38)	23.3% (15–34)	26.8% (18–38)
2. Knowledge-driven	16 questions	33.3% (22–47)	36.5% (26–48)	37.0% (27–48)	28.2% (19–40)
3. Channel-based	4 questions	24.1% (15–37)	31.1% (22–42)	11.0% (6–20)	21.1% (13–32)
3. Channel-based	8 questions	20.4% (12–33)	32.4% (23–44)	17.8% (11–28)	16.9% (10–27)
3. Channel-based	16 questions	33.3% (22–47)	39.2% (29–51)	24.7% (16–36)	32.4% (23–44)
4. Sequence-based	4 questions	18.5% (10–31)	23.0% (15–34)	13.7% (8–23)	26.8% (18–38)
4. Sequence-based	8 questions	31.5% (21–45)	37.8% (28–49)	26.0% (17–37)	23.9% (16–35)
4. Sequence-based	16 questions	37.0% (25–50)	39.2% (29–51)	27.4% (18–39)	28.2% (19–40)