
Top-1 accuracy by INTERMED complexity: five buckets (v2, repetition 1)

4 questions: strategies_272_v2_rep1_4it.eval.json, 8 questions: strategies_272_v2_rep1_8it.eval.json, 16 questions: strategies_272_v2_rep1_16it.eval.json. Top 1: the true diagnosis's ICD-11 code is among the strategy's first 1 diagnoses (exact code match). Brackets: 95% Wilson interval.



Strategy	Run	0–11 54 cases	12–15 54 cases	16–20 62 cases	21–26 48 cases	27–60 54 cases
1. Hypothesis-driven	4 questions	11.1% (5–22)	18.5% (10–31)	16.1% (9–27)	4.2% (1–14)	24.1% (15–37)
1. Hypothesis-driven	8 questions	20.4% (12–33)	25.9% (16–39)	22.6% (14–34)	22.9% (13–37)	22.2% (13–35)

Strategy	Run	0–11 54 cases	12–15 54 cases	16–20 62 cases	21–26 48 cases	27–60 54 cases
1. Hypothesis-driven	16 questions	27.8% (18–41)	35.2% (24–49)	35.5% (25–48)	25.0% (15–39)	24.1% (15–37)
2. Knowledge-driven	4 questions	20.4% (12–33)	24.1% (15–37)	17.7% (10–29)	12.5% (6–25)	18.5% (10–31)
2. Knowledge-driven	8 questions	16.7% (9–29)	27.8% (18–41)	22.6% (14–34)	25.0% (15–39)	27.8% (18–41)
2. Knowledge-driven	16 questions	33.3% (22–47)	40.7% (29–54)	32.3% (22–45)	31.2% (20–45)	31.5% (21–45)
3. Channel-based	4 questions	24.1% (15–37)	29.6% (19–43)	19.4% (11–31)	12.5% (6–25)	22.2% (13–35)
3. Channel-based	8 questions	20.4% (12–33)	37.0% (25–50)	22.6% (14–34)	8.3% (3–20)	20.4% (12–33)
3. Channel-based	16 questions	33.3% (22–47)	40.7% (29–54)	32.3% (22–45)	16.7% (9–30)	37.0% (25–50)
4. Sequence-based	4 questions	18.5% (10–31)	24.1% (15–37)	17.7% (10–29)	14.6% (7–27)	27.8% (18–41)
4. Sequence-based	8 questions	31.5% (21–45)	35.2% (24–49)	32.3% (22–45)	25.0% (15–39)	24.1% (15–37)
4. Sequence-based	16 questions	37.0% (25–50)	38.9% (27–52)	32.3% (22–45)	27.1% (17–41)	27.8% (18–41)