



Your grade: **100%**

Your latest: 100% • Your highest: 100% • To pass you need at least 80%. We keep your highest score.

Next item →

1. Compute the length of

1 / 1 point

$$\mathbf{x} = \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix}$$

using the inner product defined

$$\langle \mathbf{a}, \mathbf{b} \rangle = \mathbf{a}^T \begin{bmatrix} 2 & 1 & 0 \\ 1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix} \mathbf{b}$$

Do the exercise using pen and paper.

- ☒ $\sqrt{26}$
- ☐ $\sqrt{29}$
- ☐ $\sqrt{11}$
- ☐ $\sqrt{31}$
- ☐ 26

✓ **Correct**
Good job.

2. Compute the squared distance between

$$\mathbf{x} = \begin{bmatrix} \frac{1}{2} \\ -1 \\ -\frac{1}{2} \end{bmatrix}$$

and

$$\mathbf{y} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

using the inner product defined as

$$\langle \mathbf{a}, \mathbf{b} \rangle = \mathbf{a}^T \begin{bmatrix} 2 & 1 & 0 \\ 1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix} \mathbf{b}$$

Do the exercise using pen and paper.

☐ $\frac{9}{2}$

☐ $\sqrt{5}$


☒ 5

☐ $\sqrt{\frac{9}{2}}$



Correct

Well done.



3. Compute the length of $\mathbf{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ using the inner product defined by

$$\langle \mathbf{a}, \mathbf{b} \rangle = \mathbf{a}^T \frac{1}{2} \begin{bmatrix} 5 & -1 \\ -1 & 5 \end{bmatrix} \mathbf{b}$$

Do the exercise using pen and paper.

- ☐ $\sqrt{12}$
- ☐ 6
- ☐ $\sqrt{2}$
- ☒ $\sqrt{6}$
- ☐ 12

✔ Correct
Good job!

4. Compute the distance (not squared) between

1 / 1 point

$$\mathbf{x} = \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix}$$

and

$$\mathbf{y} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

using the inner product defined as

$$\langle \mathbf{a}, \mathbf{b} \rangle = \mathbf{a}^T \begin{bmatrix} 2 & 1 & 0 \\ 1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix} \mathbf{b}$$

Do the exercise using pen and paper (and calculator if necessary). Please enter a decimal number.

6.48

✓ Correct

Well done!

5. Compute the length of $\mathbf{x} = \begin{bmatrix} -1 \\ -1 \\ -1 \end{bmatrix}$ using the inner product defined as $\langle \mathbf{a}, \mathbf{b} \rangle = \mathbf{a}^T \mathbf{I} \mathbf{b}$ where \mathbf{I} is the identity matrix.

1 / 1 point

Do the exercise using pen and paper.

☒ $\sqrt{3}$

☐ $-\sqrt{3}$

☐ 3

☐ -3

✓ Correct

Well done! Our inner product is the dot product.