

AUDIT-C Alcohol Use Disorders Identification Test — consumption items

Name _____

Date of birth _____

Date _____

A standard drink is 14 g of pure alcohol: a 12 oz beer, a 5 oz glass of wine, or 1.5 oz of spirits. Tick one answer for each question.

1. How often do you have a drink containing alcohol?

- ☐ Never **0** ☐ Monthly or less **1** ☐ 2–4 times a month **2** ☐ 2–3 times a week **3**
☐ 4 or more times a week **4**

2. How many standard drinks containing alcohol do you have on a typical day when you are drinking?

- ☐ 1 or 2 **0** ☐ 3 or 4 **1** ☐ 5 or 6 **2** ☐ 7 to 9 **3** ☐ 10 or more **4**

3. How often do you have six or more drinks on one occasion?

- ☐ Never **0** ☐ Less than monthly **1** ☐ Monthly **2** ☐ Weekly **3** ☐ Daily or almost daily **4**

For office coding _____ + _____ + _____ = Total (0–12)

INTERPRETING THE TOTAL

0–2	Negative screen	In women, a score of 3 or more is already positive.
3+	Positive in women	Discuss the pattern; assess for a use disorder.
4+	Positive in men	Discuss the pattern; assess for a use disorder.
8+	Hazardous drinking	A score of 8 or more indicates active hazardous or harmful use.

A positive screen is a conversation, not a diagnosis. Ask about the pattern, about withdrawal, and about what the patient wants to change.