


Class II Cavity Preparation for Resin Composite — Reference Pictures


Year 3 Preclinical Conservative Dentistry · Cairo University · Week 3 demonstration / Week 4 practical (teeth 36 & 16)




1. Proximal and occlusal views. Left: compound preparation (proximal box + occlusal extension following the central groove). Right: proximal box only (slot), when the occlusal surface is sound.




2. Occlusal outline on a typodont molar. Extension follows the central and buccal/lingual grooves with smooth curves; width not more than $\frac{1}{4}$ of the intercusp distance.




3. Breaking the contact. The proximal box extends just beyond the contact with the adjacent tooth (dotted line); the occlusal extension follows the grooves.




4. Dimensions — occlusal view (from the figure): proximal box 2 mm, isthmus 1.5 mm, occlusal extension 1 mm and 2.5 mm as marked.




5. Dimensions — proximal view (from the figure): box 2 mm wide occlusally, 2.5 mm at the gingival floor, 2 mm deep.



6. Checking the proximal box with a periodontal probe. (a) occlusal view of the box; (b–d) depth of the box and axial wall from the proximal side, probe parallel to the long axis; (e) gingival floor and axial wall clearance.



7. Bucco-lingual extent of the proximal box: the box walls end 1–1.5 mm from the cusp peaks.



8. Cavity nomenclature: walls, line angles (1st S / 2nd S) and point angles in Class I, Class II (d, e), Class III and Class V preparations.