



**SpO₂/FiO₂ ratio (SF ratio) as a predictor of
mortality in ICU patients: Retrospective study
using MIMIC Database.**

Ahmed Gobba

**Capstone Final Report for BSc (Honours) in
Mathematical, Computational and Statistical Sciences**

Supervised by: Willem Van Den Boom

AY 2019/2020

YaleNUSCollege

Contents

1	Background Information	1
1.1	Introduction	1
1.2	Aim and Objectives	3
	Bibliography	4
A	None	5

List of Tables

List of Figures

Chapter 1

Background Information

1.1 Introduction

Allocation of resources to critical care patients to minimize mortality is a priority for healthcare professionals. As such the search for connections between various physiological indicators and unfavourable outcomes such as extreme illnesses or even mortality is a constant priority of researchers. One such pulmonary physiological indicator is the $\text{PaO}_2/\text{FiO}_2$ ratio (PF ratio). PaO_2 refers to the partial pressure of oxygen in arterial blood and is measured in mmHg via drawing a sample of blood from an artery in the wrist, that then is tested in a laboratory. FiO_2 refers to the initial fraction of inspired oxygen and is approximately 21% in breathable atmosphere and can be controlled with the use of mechanical ventilation. The PF ratio is importantly used in the diagnosis of extreme illnesses such as Acute Respiratory Distress Syndrome (ARDS) and has been shown to be predictor of mortality in specific subsets of patients such as newborns with Meconium aspiration syndrome (MAS) (Narayanan et al., 2019).

However, there are challenges associated with measurement of PaO_2

specifically. Most importantly, the procedure is invasive and as such is not easy to measure for all patients and to track at frequent intervals. A more convenient value to measure however is SpO_2 or peripheral capillary oxygen saturation, an estimate of the amount of oxygen in the blood. It is measured using pulse oximetry, a noninvasive method for monitoring a person's oxygen saturation. Moreover, $\text{SpO}_2/\text{FiO}_2$ ratio has been shown to be a non-invasive surrogate for $\text{PaO}_2/\text{FiO}_2$ ratio to diagnose subsets of patients such as children with ALI or ARDS (Rice et al., 2007) and burned children with smoke inhalation injury (Cambiaso-Daniel et al., 2017).

A retrospective study found that the $\text{SpO}_2/\text{FiO}_2$ ratio Time-at-Risk (SF-TAR), defined as the total time spent with severe hypoxemia (SF ratio ≤ 145), is not only significantly correlated with hospital mortality for mechanically ventilated patients, but is as well or a better predictor of it than arterial gas-derived measurements of the PF ratio.

Moreover, there have been several studies that aim to link SpO_2 to mortality. In 2015, the Tromsø study concluded that an $\text{SpO}_2 \leq 95\%$ is associated by all-cause mortality and mortality caused by pulmonary diseases (over a 10-year follow-up period) after adjusting for sex, age, history of smoking, self-reported diseases and respiratory symptoms, BMI, and CRP concentration. When Forced Expiratory Volume (FEV1) was included as a covariate, the correlation remained significant for mortality due to pulmonary diseases but no longer significant for all-cause mortality (Vold et al., 2015).

However, a prospectively planned meta-analysis participant data from 5 randomized clinical trials (conducted from 2005-2014) of infants born

before 28 weeks' gestation period found no significant difference between a lower SpO₂ target range (85%-89%) and a higher SpO₂ target range (91%-95%) on mortality or major disability at a corrected age of 18 to 24 months (Askie et al., 2018). Therefore, it seems that the use of SpO₂ as a predictor of mortality might not be applicable to all patient phenotypes, with a possibility of further sub-phenotyping.

1.2 Aim and Objectives

The main goal of this Capstone can be summarized in the following statement:

Investigate whether SpO₂/FiO₂ ratio is a statistically significant predictor of mortality in general ICU patient population or subsets thereof using a retrospective analysis of data.

The use of SpO₂/FiO₂ ratio instead of solely SpO₂ allows us to account for the different levels of mechanical ventilation that an ICU patient receives. In essence, it allows us to account for the patient's ability to convert inspired oxygen to peripheral oxygen saturation at the tissue level.

Bibliography

- Askie, Lisa M et al. (2018). "Association between oxygen saturation targeting and death or disability in extremely preterm infants in the neonatal oxygenation prospective meta-analysis collaboration". In: *Jama* 319.21, pp. 2190–2201.
- Cambiaso-Daniel, Janos et al. (2017). "Correlation Between PaO₂/FIO₂ and Peripheral Capillary Oxygenation/FIO₂ in Burned Children With Smoke Inhalation Injury." In: *Pediatric critical care medicine: a journal of the Society of Critical Care Medicine and the World Federation of Pediatric Intensive and Critical Care Societies* 18.10, e472–e476.
- Narayanan, Anand et al. (2019). "PaO₂/FiO₂ Ratio as Predictor of Mortality in Neonates with Meconium Aspiration Syndrome". In: *American journal of perinatology* 36.06, pp. 609–614.
- Rice, Todd W et al. (2007). "Comparison of the SpO₂/FIO₂ ratio and the PaO₂/FIO₂ ratio in patients with acute lung injury or ARDS". In: *Chest* 132.2, pp. 410–417.
- Vold, Monica Linea et al. (2015). "Low oxygen saturation and mortality in an adult cohort: the Tromsø study". In: *BMC pulmonary medicine* 15.1, p. 9.

Appendix A

None