1. **Which of these external genitalia in the female is considered the most sensitive region?**
    A. The clitoris
    B. The labia majora
    C. The labia minora
    D. The vaginal opening
2. **Which of these internal female reproductive organs produce a mucous secretion to help nourish the spermatozoa?**
    A. The vagina
    B. The cervix
    C. The uterus
    D. The ovaries
3. **The triangular mound of fatty tissue that covers the pubic bone**
    A. Mons Pubis
    B. Labia Majora
    C. The perineum.
    D. Labia Minora
4. **. . . . . are comparable to the scrotum in males**
    A. Labia Minora
    B. Labia Majora
    C. The ovaries
    D. The uterus
5. **Which of the following is incorrect statement about Labia Majora**
    A. Fleshy folds of tissue that extended from the mons pubis **to** the perineum.
    B. Protect the other external reproductive organs.
    C. contain sweat and oil-secreting glands
    D. Hairless folds.
6. **Which of the following is a correct statement about uterine tubes?**
    A. They are directly connected to the ovaries.
    B. They are the site of implantation.
    C. They have no role in fertilization.

D. They consist of the infundibulum, ampulla, and isthmus.

7. . . . . .glands are found on each side of the vestibule.

A. penile bulb.

B. Vestibular bulb:

C. Bartholin's glands.

D. The lesser vestibular glands

8. . . . . glands are homologous to the male prostate.

A. Vestibular bulb

B. Bartholin's glands.

C. The greater vestibular

D. The lesser vestibular glands

9. The vagina communicates superiorly with the . . . . .

A. Rectum

B. The uterus

C. urinary bladder

D. cervical canal and opens

Postpartum Physiological Changes and its Management

10. During which period does the puerperium refer to?

A. The second trimester of pregnancy

B. The six week period after delivery

C. The first trimester of pregnancy

D. The third trimester of pregnancy

11. How is uterine involution assessed?

A. By counting the number of contractions

B. By measuring the fundus by fingerbreadths

C. By measuring the length of the umbilical cord

D. By estimating the weight of the baby

12. Which type of lochia should the nurse expect to find in a client 2 days PP?

A. Lochia delta

B. Lochia serosa

C. Lochia alba

D. Lochia rubra

13. The postpartum period is also referred to as _____ or the fourth stage of pregnancy.

A. Puerperium
B. Menarche
C. Lactation
D. Gestation

14. What is the postpartum period?

A. The first trimester
B. The time during pregnancy
C. The time before conception
D. The interval between birth and return of reproductive organs to their nonpregnant state

15. How long does the postpartum period traditionally last?

A. 2 weeks
B. 4 weeks
C. 6 weeks
D. 8 weeks

16. What is the involution process in the context of the uterus?

A. The involution process is the onset of labor contractions.
B. The involution process is the expansion of the uterus during pregnancy.
C. The involution process is the formation of the placenta after conception.
D. The involution process is the return of the uterus to a nonpregnant state following birth.

17. How rapidly does the involution process of the uterus progress?

A. Fundus descends 1 every 24 hours
B. Fundus does not descend after birth
C. Fundus descends 5 cm every 24 hours
D. Fundus descends 1 to 2 cm every week

18. How many weeks after childbirth is the uterus no longer palpable?

A. 2 weeks after childbirth the uterus is no longer palpable.
B. 1 week after childbirth the uterus is no longer palpable.
C. 4 weeks after childbirth the uterus is no longer palpable.
D. 6 weeks after childbirth the uterus is no longer palpable.

A. Critical period for monitoring complications such as, Postpartum hemorrhage (PPH), Uterine atony

A. Partum
B. Late Postpartum
C. Early Postpartum
D. Immediate Postpartum

19. Late Postpartum

A. First 24 hours
B. Day 2 to Day 7
C. Week 2 to 6
D. 6 or up to 12 weeks

20. Assisted reproductive technologies used to treat infertility include the following:

A. Induction of ovulation
B. In vitro fertilization (IVF)
C. Zygote intrafallopian transfer ZIFT)
D. Gamete intrafallopian transfer (GIFT)

B. **To be considered a PPH, what would the estimated blood loss have to be for a C-section?**

A. < 550 ML
B. > 600 ML
C. > 1000 ML
D. < 900 ML

21. **Fertility in women is impaired by**

(a) Hernia

(b) High blood pressure

(c) Obesity

(d) Gallstones

22.**Infertility is when a couple fails to conceive after**

(a) 4 years of unprotected sex

(b) 3years of unprotected sex

(c) 2 years of unprotected sex

(d) 1 year of unprotected sex

C. **In women, this sexually transmitted disease can result in infertility**

(a) Human papillomavirus (HPV)

(b) Pelvic inflammatory disease

(c) Genital herpes

(d) Hernia

1.	Sterility defines as failure of conception due to incurable cause.	**T**
2.	Failure of conception after a period of previous fertility is secondary infertility.	**T**
3.	PPH is defined as blood loss of > 500 ml following caesarean delivery within 24 hours of childbirth.	**F**
4.	Late Postpartum start 2 to 6 or up to 12 weeks	**T**
5.	Lochia rubra bright red, lasts 4-10 days	**F**
6.	Major obstetric hemorrhage is denied as blood loss in excess of 1,000 ml.	**T**
7.	Most mothers resume menstruation within 7 to 9 weeks after childbirth.	**T**
8.	In lactating mothers, menstruation usually 3-4 months, and sometimes as late as 24 months.	**T**

9.	Postpartum physiological bradycardia mean decreases to rate 60 to 70 b/m for 24-48 hrs	**T**
10.	**PPH** is defined as blood loss of more than 24 hours after childbirth but within 6 weeks postpartum.	**T**
11.	Failure to ovulate is the most common infertility problem.	**T**
12.	The main cause of male infertility is diseases of the hypothalamus	**F**
13.	Pelvic ultrasonograms most frequently used diagnostic tool to assess endometrial cavity	**F**