High Risk Pregnancy

https://hiremathhospital.in/wp-content/uploads/2024/06/Hiremath-Hopital-16-940x470.png

5} Assessing of Pitting Edema

Goal:

To assess degree, site and types of edema in pregnant woman suffering from common medical disorder



Objectives:

By the end of this procedure the students will be able to achieve the following:

1. Define the term of pitting edema during pregnancy.
2. Estimate causes of pitting edema during pregnancy.
3. Distinguishes between different degrees of pitting edema.
4. Apple to assess women with pitting edema
5. Demonstrate the assessment of pitting edema to the pregnant woman.
6. Describe signs & symptoms that woman with pitting edema has.

introduction:

Definition of edema:

- The swelling of body parts on account of fluid retention is referred to as edema.
- Edema is basically an indication of some other underlying problem or condition.

Types of edema:

1) Pitting edema

2) Non pitting edema.

Causes:

- Blood volume increases by 50 %. So, there is tendency of retaining more fluids than normal while pregnancy.
- This fluid retention leads to edema or swelling.
- Baby grows; there is a growing pressure on veins, especially on the large vein called the vena cava, which returns blood to the heart from limbs.
- Because of this growing pressure the blood flow slows down, leading to the swelling of feet and legs.

Steps:

- Observe general appearance of the woman (face, hands, and hand ring place in the finger and eye lids).
- Make pressure using three middle fingers in pendicular line on tibia shaft for 30 seconds then assess the degree of pitting occurred.
- Determine the degree of edema as the following :

- 1+ →Minimal edema of lower extremities (at pedal and pretibial sites.
- 2+ →Marked edema of lower extremities
- 3+ →Edema of lower extremities, face, hands, and sacral area.
- 4+ →Generalized massive edema that includes ascites (accumulation of fluid in peritoneal cavity.

Degree	Site	Depth	Rebound

+1	Minimal edema of lower limbs	2mm	Immediately
+2	Moderate edema of lower limbs	4mm	A few seconds to rebound
+3	Marked edema of lower limbs, face, hands	6mm	10-12 seconds
+4	Generalized edema	8mm	>20 seconds

- ☞ Reassure mother about result.
- ☞ Chart the time and result of test.

Nursing Action	Second Trial		First Trial	
	Done	Not done	Done	Not done
1. Greet the mother.				
2. Explain the procedure.				

Pitting Edema Assessment

3. Edema should be assessed routinely at each prenatal visit after 20 weeks' gestation.				
4. Ask the women and /or family members if the women's face or hands appear swollen.				
5. Position the mother flat in the bed.				
6. Wash hands.				
Edema Assessment Steps				
7. Observe for general appearance (skin texture, swelling), face, hands, and hand ring place in the finger and eye lids, extremities, and sacral area for signs of pitting edema.				
8. Use three middle fingers in pendicular line on tibia shaft for 30 seconds then assess the degree of pitting occurred.				
9. Evaluate: - Extensiveness of the edema. - Depth of the depression. - Length of time it takes to clear (pitting edema leaves a depression after finger pressure has been applied to an affected area).				

10.Grade the pitting edema according to the following scale: - 1+ = Minimal edema of the lower extremities (at pedal and pretibial sites). - 2+ = Marked edema of the lower extremities. - 3+= edema of the lower extremities, face, hands, and sacral area. - 4+ = Generalized, massive edema.				
Post Procedure Task				
13. Record and. compare the finding with those previously recorded.				
14. Cover the woman and keep her in comfortable position.				
15.Reassure mother about result.				
Total score:				
Date:				

Student Name **Instructor Name**

6} Deep Tendon Reflex Assessment

Goals:

Objectives: