

Numerical Analysis

Ahmed Mohamed

December 23, 2025

1 Introduction

Matrix: Rectangular array of numbers, symbols or expressions arranged in rows and column

Ex: 2×3 matrix would look like this

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$$

The number of elements in the matrix is $2 \times 3 = 6$

Remark: Matrices are used in different mathematical operations such as addition, multiplication, and finding determinants and inverses.

General form for the matrix $m \times n$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{bmatrix}$$

Matrix Applications:

1. Computer Graphics
 - Scaling and rotation images
 - 3D
2. Machine Learning:
 - Features of data points in algorithms, Such as linear regression, Neural networks
3. Economics
 - Building an economy system
 - Analyze the strategies
4. Single processing
 - Can be used in Image processing
5. Control systems
 - Finding the relationship between inputs/outputs in databases
6. Statistics
 - Regression analysis

2 Matrices Types

1. **Row Matrix:** Have only one row

EX: $A = 1 \times 3$

$$A = \begin{bmatrix} 4 & 0 & \frac{1}{3} \end{bmatrix}$$

2. **Column Matrix:** Have only one column

EX: $A = 3 \times 1$

$$A = \begin{bmatrix} 5 \\ 33 \\ 0.4 \end{bmatrix}$$

3. **Square Matrix:** Have an equal number of rows and columns

EX: $A = 3 \times 3$

$$A = \begin{bmatrix} 5 & -1 & 2 \\ 33 & 3 & 20 \\ 0.4 & -3 & 0 \end{bmatrix}$$

EX: $B = 1 \times 1$

$$B = [3.14]$$

4. **Diagonal Matrix:** All non-diagonal elements are zeros

EX: $A = 3 \times 3$

$$A = \begin{bmatrix} \boxed{3} & 0 & 0 \\ 0 & \boxed{3} & 0 \\ 0 & 0 & \boxed{8} \end{bmatrix}$$

5. **Scalar Matrix:** Square matrix with all non-diagonal elements are zero and all diagonal elements are equal

EX: $A = 3 \times 3$

$$A = \begin{bmatrix} \boxed{3} & 0 & 0 \\ 0 & \boxed{3} & 0 \\ 0 & 0 & \boxed{3} \end{bmatrix}$$

6. **Unit Matrix(Identity):** Type of diagonal matrix, its all diagonal elements equal to 1

EX: $A = 3 \times 3$

$$A = \begin{bmatrix} \boxed{1} & 0 & 0 \\ 0 & \boxed{1} & 0 \\ 0 & 0 & \boxed{1} \end{bmatrix}$$

7. **Null(Zero) Matrix:** All elements equal to 0

EX: $A = 3 \times 3$

$$A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

8. **Symmetric Matrix:** If the matrix A is equal to A^T

Where

A^T : is the transpose matrix from A , Generated by switching rows and columns

EX: Check if $A = A^T$

$$A = \begin{bmatrix} -1 & 2 & 10 \\ 30 & 0 & 1 \\ -3 & -2 & -5 \end{bmatrix} \rightarrow A^T = \begin{bmatrix} -1 & 30 & -3 \\ 2 & 0 & -2 \\ 10 & 1 & -5 \end{bmatrix}$$

$$\therefore A \neq A^T$$

$\therefore A$ Is not a symmetric matrix

3 Addition And Subtraction

Two matrices can be added or subtracted if and only if they have the same order which means the same numbers of rows and columns

Ex: Find $A + B$ and $A - B$ Where

$$A = \begin{bmatrix} 2 & 0 \\ -5 & 6 \end{bmatrix} \quad B = \begin{bmatrix} 3 & 6 \\ 4 & 1 \end{bmatrix}$$

Sol:

$$A + B = \begin{bmatrix} 2+3 & 0+6 \\ -5+4 & 6+1 \end{bmatrix} \rightarrow A + B = \begin{bmatrix} 5 & 6 \\ -1 & 7 \end{bmatrix}$$

$$A - B = \begin{bmatrix} 2-3 & 0-6 \\ -5-4 & 6-1 \end{bmatrix} \rightarrow A - B = \begin{bmatrix} -1 & -6 \\ -9 & 5 \end{bmatrix}$$

Ex: Find $A + B$ Where

$$A = \begin{bmatrix} 1 & 5 \\ 6 & 7 \\ 8 & 9 \end{bmatrix} \quad B = \begin{bmatrix} 3 & 6 \\ 4 & 1 \end{bmatrix}$$

Sol: $A + B$ Is not defined since they have different orders

Properties Of Addition:

1. $A + B = B + A$
2. $(A + B) + C = A + (B + C)$
3. $A + 0$ (zero matrix) $= A$
4. $A - B \neq B - A$
5. $(A - B) - C = A - (B - C)$
6. $A^{\top\top} = A$
7. $aA^{\top} = aA^{\top}$ Where a is a number
8. $(A + B)^{\top} = B^{\top} + A^{\top}$

H.W(1): Prove that all the properties of matrix addition are satisfied

$$A = \begin{bmatrix} -1 & 3 & -2 \\ 0 & 1 & -10 \\ 3 & 5 & 7 \end{bmatrix} \quad B = \begin{bmatrix} 6 & -5 & -3 \\ -2 & 1 & -0 \\ 4 & -7 & 10 \end{bmatrix} \quad C = \begin{bmatrix} 8 & 6 & -1 \\ 20 & 0 & 10 \\ -10 & 3 & -30 \end{bmatrix}$$

4 Matrix multiplication

To defined the multiplication of two matrices AB , If and only if the number of columns in A is equal to the number of rows in B

Ex: Find AB Where

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 0 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 1 & 4 \\ 3 & 0 & 5 \end{bmatrix}$$

Sol:

$$AB = \begin{bmatrix} 1 \times 2 + 2 \times 3 & 1 \times 1 + 2 \times 0 & 1 \times 4 + 2 \times 5 \\ 3 \times 2 + 4 \times 3 & 3 \times 1 + 4 \times 0 & 3 \times 4 + 4 \times 5 \\ 0 \times 2 + 1 \times 3 & 0 \times 1 + 1 \times 0 & 0 \times 4 + 1 \times 5 \end{bmatrix} \Rightarrow AB = \begin{bmatrix} 8 & 1 & 14 \\ 18 & 3 & 32 \\ 3 & 0 & 5 \end{bmatrix}$$

Properties of multiplication:

1. $A(B + C) = AB + AC$
2. $(B + C)A = BA + CA$
3. $AB \neq BA$
4. $AB(C) = A(BC) = ABC$
5. $AI = IA = A$
6. $AA^{-1} = I$
Where A^{-1} : is the inverse of A
7. $(AB)^{\top} = B^{\top}A^{\top}$

5 Determinant of matrix

Single number can be found only for 2×2 matrices, Will be denoted by $\det(A)$ or $|A|$

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

$$\det(A) = |A| = a_{11} \times a_{22} - a_{12} \times a_{21}$$

Ex: Find Determinant of A Where

$$A = \begin{bmatrix} 6 & 4 \\ 7 & 9 \end{bmatrix} \Rightarrow |A| = 6 \times 9 - 4 \times 7 \Rightarrow |A| = 26$$

For a 3×3 matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$A = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

$$|A| = a_{11}(a_{22}a_{33} - a_{23}a_{32}) - a_{12}(a_{21}a_{33} - a_{23}a_{31}) + a_{13}(a_{21}a_{32} - a_{22}a_{31})$$

Ex: Find $|A|$

Where

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & -1 & 0 \\ -2 & 0 & 3 \end{bmatrix}$$

Sol:

$$A = 1 \times \begin{bmatrix} -1 & 0 \\ 0 & 3 \end{bmatrix} - 2 \times \begin{bmatrix} 0 & 0 \\ -2 & 3 \end{bmatrix} + 4 \times \begin{bmatrix} 0 & -1 \\ -2 & 0 \end{bmatrix}$$

$$|A| = 1(-1 \times 3 - 0 \times 0) - 2(0 \times 3 - 0 \times -2) + 4(0 \times 0 - 1 \times -2) \rightarrow |A| = 5$$

6 Cofactor of a matrix

$$\text{let } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

We can find the Cofactor of A by

$$C_{ij} = (-1)^{i+j} A_{ij}$$

$$C_{11} = -1^2 \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} \Rightarrow C_{11} = a_{22}a_{33} - a_{23}a_{32}$$

$$C_{12} = (-1)^{1+2} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} \Rightarrow C_{12} = -(a_{21}a_{33} - a_{23}a_{31})$$

$$C_{13} = (-1)^{1+3} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \Rightarrow C_{13} = a_{21}a_{32} - a_{22}a_{31}$$

$$C_{21} = (-1)^{2+1} \begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix} \Rightarrow C_{21} = -(a_{12}a_{33} - a_{13}a_{32})$$

$$C_{22} = (-1)^{2+2} \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} \Rightarrow C_{22} = a_{11}a_{33} - a_{13}a_{31}$$

$$C_{23} = (-1)^{2+3} \begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} \Rightarrow C_{23} = -(a_{11}a_{32} - a_{12}a_{31})$$

$$C_{31} = (-1)^{3+1} \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix} \Rightarrow C_{31} = a_{12}a_{23} - a_{13}a_{22}$$

$$C_{32} = (-1)^{3+2} \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} \Rightarrow C_{32} = -(a_{11}a_{23} - a_{13}a_{21})$$

$$C_{33} = (-1)^{3+3} \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \Rightarrow C_{33} = a_{11}a_{22} - a_{12}a_{21}$$

Ex(2): Find the cofactor for the following

$$A = \begin{bmatrix} 3 & 4 & -1 \\ 1 & 0 & 3 \\ 2 & 5 & -4 \end{bmatrix}$$

$$C_{11} = (-1)^{1+1} \begin{vmatrix} 0 & 3 \\ 5 & -4 \end{vmatrix} \Rightarrow C_{11} = 0 \cdot (-4) - 3 \cdot 5 = -15$$

$$C_{12} = (-1)^{1+2} \begin{vmatrix} 1 & 3 \\ 2 & -4 \end{vmatrix} \Rightarrow C_{12} = -(1 \cdot (-4) - 3 \cdot 2) = -(-4 - 6) = 10$$

$$C_{13} = (-1)^{1+3} \begin{vmatrix} 1 & 0 \\ 2 & 5 \end{vmatrix} \Rightarrow C_{13} = 1 \cdot 5 - 0 \cdot 2 = 5$$

$$C_{21} = (-1)^{2+1} \begin{vmatrix} 4 & -1 \\ 5 & -4 \end{vmatrix} \Rightarrow C_{21} = -(4 \cdot (-4) - (-1) \cdot 5) = -(-16 + 5) = 11$$

$$C_{22} = (-1)^{2+2} \begin{vmatrix} 3 & -1 \\ 2 & -4 \end{vmatrix} \Rightarrow C_{22} = 3 \cdot (-4) - (-1) \cdot 2 = -12 + 2 = -10$$

$$C_{23} = (-1)^{2+3} \begin{vmatrix} 3 & 4 \\ 2 & 5 \end{vmatrix} \Rightarrow C_{23} = -(3 \cdot 5 - 4 \cdot 2) = -(15 - 8) = -7$$

$$C_{31} = (-1)^{3+1} \begin{vmatrix} 4 & -1 \\ 0 & 3 \end{vmatrix} \Rightarrow C_{31} = 4 \cdot 3 - (-1) \cdot 0 = 12$$

$$C_{32} = (-1)^{3+2} \begin{vmatrix} 3 & -1 \\ 1 & 3 \end{vmatrix} \Rightarrow C_{32} = -(3 \cdot 3 - (-1) \cdot 1) = -(9 + 1) = -10$$

$$C_{33} = (-1)^{3+3} \begin{vmatrix} 3 & 4 \\ 1 & 0 \end{vmatrix} \Rightarrow C_{33} = 3 \cdot 0 - 4 \cdot 1 = -4$$

$$C = \begin{bmatrix} -15 & 10 & 5 \\ 11 & -10 & -7 \\ 12 & -10 & -4 \end{bmatrix}$$

Remark: if A is a matrix of 2×2 you can find the cofactor like this

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \Rightarrow C = \begin{bmatrix} a_{22} & -a_{21} \\ -a_{12} & a_{11} \end{bmatrix}$$

Ex: Find cofactor for A Where

$$A = \begin{bmatrix} -2 & -4 \\ 10 & 5 \end{bmatrix}$$

Sol:

$$A = \begin{vmatrix} -2 & -4 \\ 10 & 5 \end{vmatrix} \Rightarrow C = \begin{vmatrix} 5 & -10 \\ 4 & -2 \end{vmatrix}$$

7 Adjoint Matrix

$\text{Adj}(A)$ Can be found $\text{adj}(A) = C^\top$ Where C is the cofactor of A

Ex(3): Find adjoint for A from the example number 2

$$C = \begin{bmatrix} 15 & 10 & 5 \\ 11 & -10 & -7 \\ 12 & -10 & -4 \end{bmatrix}$$

Sol:

$$\text{adj}(A) = C^\top = \begin{bmatrix} -15 & 11 & 12 \\ 10 & -10 & -10 \\ 5 & -7 & -4 \end{bmatrix}$$

8 Inverse Matrix (A^{-1})

You can find the inverse from this form $A^{-1} = \frac{\text{adj}(A)}{|A|}$, Where $|A| \neq 0$

If $|A| = 0$ Then A doesn't have inverse

Ex: Find A^{-1} from Ex(3)

$$\text{adj}(A) = C^\top = \begin{bmatrix} -15 & 11 & 12 \\ 10 & -10 & -10 \\ 5 & -7 & -4 \end{bmatrix}$$

Sol:

After Finding the cofactor it will equal $|A| = -10$

$$A^{-1} = \frac{\text{adj}(A)}{|A|} = \begin{bmatrix} \frac{15}{10} & -\frac{11}{10} & -\frac{12}{10} \\ -\frac{10}{10} & \frac{10}{10} & \frac{10}{10} \\ -\frac{5}{10} & \frac{7}{10} & \frac{4}{10} \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} 1.5 & -1.1 & -1.2 \\ -1 & 1 & 1 \\ -0.5 & 0.7 & 0.4 \end{bmatrix}$$

9 Lower & Upper Triangular Matrix

Lower Triangular Matrix All the elements above the main diagonal are zeros

Ex: 3×3 Lower Triangular matrix

$$A = \begin{bmatrix} a_{11} & \boxed{0} & \boxed{0} \\ a_{21} & a_{22} & \boxed{0} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Upper Triangular Matrix All the elements below the main diagonal are zeros

Ex: 3×3 Upper Triangular matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ \boxed{0} & a_{22} & a_{23} \\ \boxed{0} & \boxed{0} & a_{33} \end{bmatrix}$$

Ex: Transfer A to upper triangular matrix, Where

$$A = \begin{bmatrix} 0 & 4 & 2 \\ 2 & 3 & 5 \\ 3 & 1 & 1 \end{bmatrix} \begin{array}{l} \rightarrow R_1 \\ \rightarrow R_2 \\ \rightarrow R_3 \end{array}$$

Sol:

$R_1 = R_2$, Swapping two rows

$$A = \begin{bmatrix} 2 & 3 & 5 \\ 0 & 4 & 2 \\ 3 & 1 & 1 \end{bmatrix}$$

Apply $R_3 = 2R_3 - 3R_1$ to the above matrix

$$A = \begin{bmatrix} 2 & 3 & 5 \\ 0 & 4 & 2 \\ 0 & -7 & -13 \end{bmatrix}$$

Apply $R_3 = 7R_2 + 4R_3$ to the above matrix

$$A = \begin{bmatrix} 2 & 3 & 5 \\ 0 & 4 & 2 \\ 0 & 0 & -13 \end{bmatrix}$$

H.W(2): Make it lower triangular matrix

10 Systems of linear equations

Linear system of equations can be written as

$$\begin{aligned} a_{11}X_1 + a_{12}X_2 + \cdots + a_{1n}X_n &= b_1 \\ a_{21}X_1 + a_{22}X_2 + \cdots + a_{2n}X_n &= b_2 \\ &\vdots \\ a_{m1}X_1 + a_{m2}X_2 + \cdots + a_{mn}X_n &= b_m \end{aligned}$$

This system can be transfer into $AX = b$

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ x_n \end{bmatrix} \quad b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

To solve this system of linear equations, We can use the following methods

1. **Gauss Elimination Method** To use this method we need to transfer the matrix A into a lower or upper Triangular matrix using row operations

Ex: Solve the following system of the linear equations using Gauss method, Where

$$\begin{aligned} 4y + 2z &= 1 \\ 2x + 3y + 5z &= 0 \\ 3x + y + z &= 11 \end{aligned}$$

Sol:

$$AX = b, \quad AX = \begin{bmatrix} 0 & 4 & 2 \\ 2 & 3 & 5 \\ 3 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad b = \begin{bmatrix} 1 \\ 0 \\ 11 \end{bmatrix}$$

$$\left[\begin{array}{ccc|c} 0 & 4 & 2 & 1 \\ 2 & 3 & 5 & 0 \\ 3 & 1 & 1 & 11 \end{array} \right]$$

$$R_1 = R_2$$

\Downarrow

$$\left[\begin{array}{ccc|c} 2 & 3 & 5 & 0 \\ 0 & 4 & 2 & 1 \\ 3 & 1 & 1 & 11 \end{array} \right]$$

$$R_3 = 2R_3 - 3R_1$$

$$\Downarrow$$

$$\left[\begin{array}{ccc|c} 2 & 3 & 5 & 0 \\ 0 & 4 & 2 & 1 \\ 0 & -7 & -13 & 22 \end{array} \right]$$

$$R_3 = 4R_3 - 7R_2$$

$$\Downarrow$$

$$\left[\begin{array}{ccc|c} 2 & 3 & 5 & 0 \\ 0 & 4 & 2 & 1 \\ 0 & 0 & -38 & 95 \end{array} \right]$$

$$-38z = 95 \Rightarrow z = -\frac{95}{38} = -2.5$$

$$4y + 2z = 1 \Rightarrow 4y + 2(-2.5) = 1$$

$$\Rightarrow 4y - 5 = 1$$

$$\Rightarrow 4y = 6 \Rightarrow y = \frac{3}{2} = 1.5$$

$$2x + 3y + 5z = 0 \Rightarrow 2x + 3(1.5) + 5(-2.5) = 0$$

$$\Rightarrow 2x + 4.5 - 12.5 = 0$$

$$\Rightarrow 2x - 8 = 0$$

$$\Rightarrow x = 4$$

2. Gauss-Jordan Elimination Method

We have to transfer the system to $JX = T$

$$J = \begin{bmatrix} a_{11} & 0 & 0 & \dots \\ 0 & a_{22} & 0 & \dots \\ 0 & 0 & a_{33} & \vdots \\ 0 & 0 & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} t_1 \\ t_2 \\ \vdots \\ t_n \end{bmatrix}$$

The solution will be

$$t_1, t_2, t_3, \dots, t_n$$

Ex: Solve the following system of linear equations using Gauss-Jordan

$$x + 2y + 3z = 9$$

$$2x + 3y + z = 8$$

$$3x + y + 2z = 7$$

Sol:

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 9 \\ 2 & 3 & 1 & 8 \\ 3 & 1 & 2 & 7 \end{array} \right]$$

$$R_2 = R_2 - 2R_1, R_3 = R_3 - 3R_1$$

\Downarrow

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 9 \\ 0 & -1 & -5 & -10 \\ 0 & -5 & -7 & -20 \end{array} \right]$$

$$R_2 = -R_2$$

\Downarrow

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 9 \\ 0 & 1 & 5 & 10 \\ 0 & -5 & -7 & -20 \end{array} \right]$$

$$R_3 = R_3 + 5R_2$$

\Downarrow

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 9 \\ 0 & 1 & 5 & 10 \\ 0 & 0 & 18 & 30 \end{array} \right]$$

$$R_1 = R_1 - 2R_2$$

\Downarrow

$$\left[\begin{array}{ccc|c} 1 & 0 & -7 & -11 \\ 0 & 1 & 5 & 10 \\ 0 & 0 & 18 & 30 \end{array} \right]$$

$$R_1 = 18R_1 + 7R_3, R_2 = 18R_2 - 5R_3$$

\Downarrow

$$\left[\begin{array}{ccc|c} 18 & 0 & 0 & 12 \\ 0 & 18 & 0 & 30 \\ 0 & 0 & 18 & 30 \end{array} \right]$$

$$18x = 12 \Rightarrow x = \frac{12}{18} = 0.6$$

$$18y = 30 \Rightarrow y = \frac{30}{18} = 1.667$$

$$18z = 30 \Rightarrow z = \frac{30}{18} = 1.667$$

3. **General Form For Cramer's Rule** For a system with linear equations with n variables of the form

$$\begin{aligned}a_{11}X_1 + a_{12}X_2 + \cdots + a_{1n}X_n &= b_1 \\a_{21}X_1 + a_{22}X_2 + \cdots + a_{2n}X_n &= b_2 \\&\vdots \\a_{m1}X_1 + a_{m2}X_2 + \cdots + a_{mn}X_n &= b_m\end{aligned}$$

the solution will be

$$x_i = \frac{D_i}{D}$$

Where

D : is the Determinant of the matrix A generated from the system $AX = b$

D_i : Determinant of the matrix formed by replacing the i - th column of A with the constant vector b

Ex: Solve using Cramer's rule

$$\begin{aligned}x + 2y + z &= 4 \\2x + y + 3z &= 10 \\3x + 2y + 2z &= 12\end{aligned}$$

Sol:

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 3 \\ 3 & 2 & 2 \end{bmatrix} \quad b = \begin{bmatrix} 4 \\ 10 \\ 12 \end{bmatrix}$$

$$D = |A| = \det(A) = 1 \begin{vmatrix} 1 & 3 \\ 2 & 2 \end{vmatrix} - 2 \begin{vmatrix} 2 & 3 \\ 3 & 2 \end{vmatrix} + 1 \begin{vmatrix} 2 & 1 \\ 3 & 2 \end{vmatrix} = 7$$

Now Replacing first column with the constants vector b

$$\begin{aligned}D_1 &= \begin{vmatrix} 4 & 2 & 1 \\ 10 & 1 & 3 \\ 12 & 2 & 2 \end{vmatrix} = 4 \begin{vmatrix} 1 & 3 \\ 2 & 2 \end{vmatrix} - 2 \begin{vmatrix} 10 & 3 \\ 12 & 2 \end{vmatrix} + 1 \begin{vmatrix} 10 & 1 \\ 12 & 2 \end{vmatrix} \\&= 4(1 \times 2 - 3 \times 2) - 2(10 \times 2 - 3 \times 12) + 1(10 \times 2 - 1 \times 12) \\&= 24\end{aligned}$$

$$\begin{aligned}D_2 &= \begin{vmatrix} 1 & 4 & 1 \\ 2 & 10 & 3 \\ 3 & 12 & 2 \end{vmatrix} = 1 \begin{vmatrix} 10 & 3 \\ 12 & 2 \end{vmatrix} - 4 \begin{vmatrix} 2 & 3 \\ 3 & 2 \end{vmatrix} + 1 \begin{vmatrix} 2 & 10 \\ 3 & 12 \end{vmatrix} \\&= 1(10 \times 2 - 3 \times 12) - 4(2 \times 2 - 3 \times 3) + 1(2 \times 12 - 10 \times 3) \\&= -2\end{aligned}$$

$$\begin{aligned}
D_3 &= \begin{vmatrix} 1 & 2 & 4 \\ 2 & 1 & 10 \\ 3 & 2 & 12 \end{vmatrix} = 1 \begin{vmatrix} 1 & 10 \\ 2 & 12 \end{vmatrix} - 2 \begin{vmatrix} 2 & 10 \\ 3 & 12 \end{vmatrix} + 4 \begin{vmatrix} 2 & 1 \\ 3 & 2 \end{vmatrix} \\
&= 1(1 \times 12 - 10 \times 2) - 2(2 \times 12 - 10 \times 3) + 4(2 \times 2 - 1 \times 3) \\
&= 8
\end{aligned}$$

$$\begin{aligned}
x &= \frac{D_1}{D} \Rightarrow x = \frac{24}{7} \Rightarrow x \approx 3.4 \\
y &= \frac{D_2}{D} \Rightarrow y = -\frac{2}{7} \Rightarrow y \approx 0.2 \\
z &= \frac{D_3}{D} \Rightarrow z = \frac{8}{7} \Rightarrow z \approx 1.1
\end{aligned}$$

4. Inverse Matrix Method

For system of equation written in a matrix form $AX = b$

If $A \neq 0$, Then the solution of the system will be $X = A^{-1}b$, which comes from the following

$$AX = b \Rightarrow A^{-1}AX = A^{-1}b \Rightarrow IX = A^{-1}b \Rightarrow \boxed{X = A^{-1}b}$$

Where I is a matrix with all its diagonal elements are ones

$$I_{3 \times 3} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Ex(1): solve the system using inverse method

$$2x + 3y = 8$$

$$x + 4y = 7$$

$$A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \quad b = \begin{bmatrix} 8 \\ 7 \end{bmatrix}$$

$$AX = b \Rightarrow X = A^{-1}b$$

$$A^{-1} = \frac{adj(A)}{|A|}$$

$$|A| = \begin{vmatrix} 2 & 3 \\ 1 & 4 \end{vmatrix} = 5$$

$$Adj(A) = \begin{bmatrix} 4 & -3 \\ -1 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{adj(A)}{|A|} = \begin{bmatrix} \frac{4}{5} & -\frac{3}{5} \\ -\frac{1}{5} & \frac{2}{5} \end{bmatrix}$$

$$\begin{aligned}
x &= A^{-1}b \\
\begin{bmatrix} x \\ y \end{bmatrix} &= \begin{bmatrix} \frac{4}{5} & -\frac{3}{5} \\ -\frac{1}{5} & \frac{2}{5} \end{bmatrix} \times \begin{bmatrix} 8 \\ 7 \end{bmatrix} \\
x &= \frac{4}{5} \times 8 - \frac{3}{5} \times 7 \Rightarrow x = \frac{32}{5} - \frac{21}{5} \Rightarrow x = \frac{11}{5} = 2.2 \\
y &= -\frac{1}{5} \times 8 + \frac{2}{5} \times 7 \Rightarrow y = -\frac{8}{5} + \frac{14}{5} \Rightarrow y = \frac{6}{5} \Rightarrow y = 1.2
\end{aligned}$$

Ex(2):

$$\begin{aligned}
x + y + z &= 6 \\
2y + 5z &= -4 \\
2x + 5y - z &= 27
\end{aligned}$$

Sol:

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 5 \\ 2 & 5 & -1 \end{bmatrix} \quad b = \begin{bmatrix} 6 \\ -4 \\ 27 \end{bmatrix}$$

$$AX = b \Rightarrow X = A^{-1}b, \text{ Where } A^{-1} = \frac{adj(A)}{|A|}$$

$$adj(A) = C^T$$

$$C = \begin{vmatrix} 1 & 1 & 1 \\ 0 & 2 & 5 \\ 2 & 5 & -1 \end{vmatrix}$$

$$C_{ij} = (-1)^{i+j} M_{ij}$$

$$C_{11} = (-1)^{1+1} \begin{vmatrix} 2 & 5 \\ 5 & -1 \end{vmatrix} = 1[2 \times (-1) - 5 \times 5] = -27$$

$$C_{12} = (-1)^{1+2} \begin{vmatrix} 0 & 5 \\ 2 & -1 \end{vmatrix} = -1[0 \times (-1) - 5 \times 2] = 10$$

$$C_{13} = (-1)^{1+3} \begin{vmatrix} 0 & 2 \\ 2 & 5 \end{vmatrix} = 1[0 \times 5 - 2 \times 2] = -4$$

$$C_{21} = (-1)^{2+1} \begin{vmatrix} 1 & 1 \\ 5 & -1 \end{vmatrix} = -1[1 \times (-1) - 1 \times 5] = 6$$

$$C_{22} = (-1)^{2+2} \begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix} = 1[1 \times (-1) - 1 \times 2] = -3$$

$$C_{23} = (-1)^{2+3} \begin{vmatrix} 1 & 1 \\ 2 & 5 \end{vmatrix} = -1[1 \times 5 - 1 \times 2] = -3$$

$$C_{31} = (-1)^{3+1} \begin{vmatrix} 1 & 1 \\ 2 & 5 \end{vmatrix} = 1[1 \times 5 - 1 \times 2] = 3$$

$$C_{32} = (-1)^{3+2} \begin{vmatrix} 1 & 1 \\ 0 & 5 \end{vmatrix} = -1[1 \times 5 - 1 \times 0] = -5$$

$$C_{33} = (-1)^{3+3} \begin{vmatrix} 1 & 1 \\ 0 & 2 \end{vmatrix} = 1[1 \times 2 - 1 \times 0] = 2$$

$$C = \begin{bmatrix} -27 & 10 & -4 \\ 6 & -3 & -3 \\ 3 & -5 & 2 \end{bmatrix}$$

$$\text{adj}(A) = C^T = \begin{bmatrix} -27 & 6 & 3 \\ 10 & -3 & -5 \\ -4 & -3 & 2 \end{bmatrix}$$

Finding the cofactor

$$\begin{aligned} |A| &= 1 \begin{vmatrix} 2 & 5 \\ 5 & -1 \end{vmatrix} - 1 \begin{vmatrix} 0 & 5 \\ 2 & -1 \end{vmatrix} + 1 \begin{vmatrix} 0 & 2 \\ 2 & 5 \end{vmatrix} \\ |A| &= 1(2 \times -1 - 5 \times 5) - 1(0 \times -1 - 5 \times 2) \\ &\quad + 1(0 \times 5 - 2 \times 2) \\ |A| &= -21 \end{aligned}$$

$$A^{-1} = \frac{\text{adj}(A)}{|A|} = \begin{bmatrix} \frac{-27}{-21} & \frac{6}{-21} & \frac{3}{-21} \\ \frac{10}{-21} & \frac{-3}{-21} & \frac{-5}{-21} \\ \frac{-4}{-21} & \frac{-3}{-21} & \frac{2}{-21} \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} \frac{9}{7} & -\frac{2}{7} & -\frac{1}{7} \\ -\frac{10}{21} & \frac{1}{7} & \frac{5}{21} \\ \frac{4}{21} & \frac{1}{7} & -\frac{2}{21} \end{bmatrix}$$

$$X = A^{-1}b$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{9}{7} & -\frac{2}{7} & -\frac{1}{7} \\ -\frac{10}{21} & \frac{1}{7} & \frac{5}{21} \\ \frac{4}{21} & \frac{1}{7} & -\frac{2}{21} \end{bmatrix} \times \begin{bmatrix} 6 \\ -4 \\ 27 \end{bmatrix}$$

$$x = \frac{9}{7} \times 6 + \frac{2}{7} \times 4 - \frac{1}{7} \times 27 \Rightarrow x = 5$$

$$y = -\frac{10}{21} \times 6 - \frac{1}{7} \times 4 + \frac{5}{21} \times 27 \Rightarrow y = 3$$

$$z = \frac{4}{21} \times 6 - \frac{1}{7} \times 4 - \frac{2}{21} \times 27 \Rightarrow z = -2$$

5. Lower/Upper Decomposition Method

In this method we will write $A = LU$, Where

L : Lower triangular matrix with (1)s on the main diagonal

U : Upper triangular Matrix

Then

$$L = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \quad U = \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$

Which will produce the following simplification

$$A = LU$$

$$LU = \begin{bmatrix} 1 \times u_{11} + 0 \times 0 + 0 \times 0 & 1 \times u_{12} + 0 \times u_{22} + 0 \times 0 & 1 \times u_{13} + 0 \times u_{23} + 0 \times u_{33} \\ l_{21}u_{11} + 1 \times 0 + 0 \times 0 & l_{21}u_{12} + 1 \times u_{22} + 0 \times 0 & l_{21}u_{13} + 1 \times u_{23} + 0 \times u_{33} \\ l_{31}u_{11} + l_{32} \times 0 + 1 \times 0 & l_{31}u_{12} + l_{32}u_{22} + 1 \times 0 & l_{31}u_{13} + l_{32}u_{23} + 1 \times u_{33} \end{bmatrix}$$

Simplifies this

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = LU$$

$$A = \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ l_{21}u_{11} & l_{21}u_{12} + u_{22} & l_{21}u_{13} + u_{23} \\ l_{31}u_{11} & l_{31}u_{12} + l_{32}u_{22} & l_{31}u_{13} + l_{32}u_{23} + u_{33} \end{bmatrix}$$

This will produce this equations

$$u_{11} = a_{11}$$

$$u_{12} = a_{12}$$

$$u_{13} = a_{13}$$

$$l_{21}u_{11} = a_{21}$$

$$l_{21}u_{12} + u_{22} = a_{22}$$

$$l_{21}u_{13} + u_{23} = a_{23}$$

$$l_{31}u_{11} = a_{31}$$

$$l_{31}u_{12} + l_{32}u_{22} = a_{32}$$

$$l_{31}u_{13} + l_{32}u_{23} + u_{33} = a_{33}$$

Ex:

$$2x + 3y + z = 1$$

$$4x + 7y + 5z = 2$$

$$6x + 18y + 19z = 3$$

Sol:

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 4 & 7 & 5 \\ 6 & 18 & 19 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$u_{11} = a_{11} = 2$$

$$u_{12} = a_{12} = 3$$

$$u_{13} = a_{13} = 1$$

$$l_{21}u_{11} = a_{21}$$

$$l_{21} \times 2 = 4$$

$$l_{21} = 2$$

$$l_{21}u_{12} + u_{22} = a_{22}$$

$$2 \times 3 + u_{22} = 7$$

$$6 + u_{22} = 7$$

$$u_{22} = 7 - 6$$

$$u_{22} = 1$$

$$l_{21}u_{13} + u_{23} = a_{23}$$

$$2 \times 1 + u_{23} = 5$$

$$2 + u_{23} = 5$$

$$u_{23} = 5 - 2$$

$$u_{23} = 3$$

$$l_{31}u_{11} = a_{31}$$

$$l_{31}2 = 6$$

$$l_{31} = 3$$

$$l_{31}u_{12} + l_{32}u_{22} = a_{32}$$

$$3 \times 3 + l_{32}1 = 18$$

$$9 + l_{32} = 18$$

$$l_{32} = 18 - 9$$

$$l_{32} = 9$$

$$l_{31}u_{13} + l_{32}u_{23} + u_{33} = a_{33}$$

$$3 \times 1 + 9 \times 3 + u_{33} = 19$$

$$3 + 27 + u_{33} = 19$$

$$30 + u_{33} = 19$$

$$u_{33} = 19 - 30$$

$$u_{33} = -11$$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 9 & 1 \end{bmatrix}, \quad U = \begin{bmatrix} 2 & 3 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & -11 \end{bmatrix}$$

$$Ly = b$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 9 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$1 \times y_1 + 0 \times y_2 + 0 \times y_3 = 1$$

$$2 \times y_1 + 1 \times y_2 + 0 \times y_3 = 2$$

$$2 \times 1 + y_2 = 2$$

$$2 + y_2 = 2$$

$$y_2 = 2 - 2$$

$$y_2 = 0$$

$$3 \times y_1 + 9 \times y_2 + 1 \times y_3 = 3$$

$$3 \times 1 + 9 \times 0 + y_3 = 3$$

$$3 + 0 + y_3 = 3$$

$$y_3 = 3 - 3$$

$$y_3 = 0$$

$$y = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$ux = y$$

$$\begin{bmatrix} 2 & 3 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & -11 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

From the third row

$$\begin{aligned} 0 \times x + 0 \times y - 11 \times z &= 0 \\ -11z &= 0 \\ z &= 0 \end{aligned}$$

The second row

$$\begin{aligned} 0 \times x + 1 \times y + 3 \times z &= 0 \\ 0 + y + 3(0) &= 0 \\ y &= 0 \end{aligned}$$

The first row

$$\begin{aligned} 2 \times x + 3 \times y + 1 \times z &= 1 \\ 2x + 3 \times 0 + 1 \times 0 &= 1 \\ 2x &= 1 \\ x &= \frac{1}{2} \end{aligned}$$

The past 5 methods are called numerical methods to solve a system of linear equations

Direct Methods: these methods are called iterative methods

1. Jacobi Method

General form:

$$x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j \neq i}^n a_{ij} x_j^k \right)$$

used to solve $AX = b$, Where

A : square matrix

X : Vector of unknowns

b : Vector of knowns

k : is a counter starts from 0

Where $i = 1$ so $j = 2, 3$ because $j \neq i$

$$x_1^{(k+1)} = \frac{1}{a_{11}}(b_1 - [a_{12}x_2^k + a_{13}x_3^k])$$

$i = 2$, $j = 1, 3$

$$x_2^{(k+1)} = \frac{1}{a_{22}}(b_2 - [a_{21}x_1^k + a_{23}x_3^k])$$

$i = 3$, $j = 1, 2$

$$x_3^{(k+1)} = \frac{1}{a_{33}}(b_3 - [a_{31}x_1^k + a_{32}x_2^k])$$

Method Steps

- (a) Start with initial values: $x^0 = [x_1^0, x_2^0, \dots, x_n^0]$
- (b) Compute new values x_i : $x_i^{(k+1)} = (x_1^{k+1}, x_2^{k+1}, \dots, x_n^{k+1})$
- (c) Repeat until the values converge between iterations, This mean

$$|x_i^{(k+1)} - x_i^k| < \epsilon$$

Ex: solve the following equations

$$\begin{aligned} 10x_1 + 2x_2 + x_3 &= 9 \\ x_1 + 5x_2 + x_3 &= 7 \\ 2x_1 + 3x_2 + 10x_3 &= 15 \end{aligned}$$

Sol:

$$A = \begin{bmatrix} 10 & 2 & 1 \\ 1 & 5 & 1 \\ 2 & 3 & 10 \end{bmatrix}, \quad b = \begin{bmatrix} 9 \\ 7 \\ 15 \end{bmatrix}$$

$$x_1^{(k+1)} = \frac{1}{a_{11}}(b_1 - [a_{12}x_2^k + a_{13}x_3^k])$$

$$x_2^{(k+1)} = \frac{1}{a_{22}}(b_2 - [a_{21}x_1^k + a_{23}x_3^k])$$

$$x_3^{(k+1)} = \frac{1}{a_{33}}(b_3 - [a_{31}x_1^k + a_{32}x_2^k])$$

$$x_1^{(k+1)} = \frac{1}{10}(9 - 2x_2^k - x_3^k)$$

$$x_2^{(k+1)} = \frac{1}{5}(7 - x_1^k - x_3^k)$$

$$x_3^{(k+1)} = \frac{1}{10}(15 - 2x_1^k - 3x_2^k)$$

$$k = 0 \Rightarrow (x_1^0 = 0, \quad x_2^0 = 0, \quad x_3^0 = 0)$$

$$x_1^1 = \frac{1}{10}(9 - 2x_2^0 - x_3^0) \Rightarrow x_1^1 = \frac{1}{10}(9 - 2 \times 0 - 0) \Rightarrow x_1^1 = \frac{9}{10} = 0.9$$

$$x_2^1 = \frac{1}{5}(7 - x_1^0 - x_3^0) \Rightarrow x_2^1 = \frac{1}{5}(7 - 0 - 0) \Rightarrow x_2^1 = \frac{7}{5} = 1.4$$

$$x_3^1 = \frac{1}{10}(15 - 2x_1^0 - 3x_2^0) \Rightarrow x_3^1 = \frac{1}{10}(15 - 2 \times 0 - 3 \times 0) \Rightarrow x_3^1 = \frac{15}{10} = 1.5$$

$$|x_1^1 - x_1^0| \Rightarrow |0.9 - 0| = 0.9 > \epsilon$$

$$k = 1 \Rightarrow (x_1^1 = 0.9, \quad x_2^1 = 1.4, \quad x_3^1 = 1.5)$$

$$x_1^2 = \frac{1}{10}(9 - 2x_2^1 - x_3^1) \Rightarrow x_1^2 = \frac{1}{10}(9 - 2 \times 1.4 - 1.5) \Rightarrow x_1^2 = 0.47$$

$$x_2^2 = \frac{1}{5}(7 - x_1^1 - x_3^1) \Rightarrow x_2^2 = \frac{1}{5}(7 - 0.9 - 1.5) \Rightarrow x_2^2 = 0.92$$

$$x_3^2 = \frac{1}{10}(15 - 2x_1^1 - 3x_2^1) \Rightarrow x_3^2 = \frac{1}{10}(15 - 2 \times 0.9 - 3 \times 1.4) \Rightarrow x_3^2 = 0.9$$

$$|x_1^2 - x_1^1| \Rightarrow |0.47 - 0.9| = 0.43 > \epsilon$$

$$k = 2 \Rightarrow (x_1^2 = 0.47, \quad x_2^2 = 0.92, \quad x_3^2 = 0.9)$$

$$x_1^3 = \frac{1}{10}(9 - 2x_2^2 - x_3^2) \Rightarrow x_1^3 = \frac{1}{10}(9 - 2 \times 0.92 - 0.9) \Rightarrow x_1^3 = 0.626$$

$$x_2^3 = \frac{1}{5}(7 - x_1^2 - x_3^2) \Rightarrow x_2^3 = \frac{1}{5}(7 - 0.47 - 0.9) \Rightarrow x_2^3 = 1.126$$

$$x_3^3 = \frac{1}{10}(15 - 2x_1^2 - 3x_2^2) \Rightarrow x_3^3 = \frac{1}{10}(15 - 2 \times 0.47 - 3 \times 0.92) \Rightarrow x_3^3 = 1.13$$

$$|x_1^3 - x_1^2| \Rightarrow |0.626 - 0.47| = 0.156 < \epsilon$$

$$|x_2^3 - x_2^2| \Rightarrow |1.126 - 0.92| = 0.206 < \epsilon$$

$$|x_3^3 - x_3^2| \Rightarrow |1.13 - 0.9| = 0.23 < \epsilon$$

So the solution set: $x_1^3 = 0.626, \quad x_2^3 = 1.126, \quad x_3^3 = 1.13$

2. Gauss-Seidal Iteration Formula

Suppose you have a system of 3 equations

$$(a) \quad a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1$$

$$(b) \quad a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2$$

$$(c) \quad a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3$$

To solve this system, use this form which refers to the general form

$$x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij}x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij}x_j^{(k)} \right)$$

Where:

$i : 1, 2, 3, \dots$

$k : 0, 1, 2, \dots$

n : number of equations

It can be written as the following

$i = 1$

$$x_1^{(k+1)} = \frac{1}{a_{11}} (b_1 - a_{12}x_2^{(k)} - a_{13}x_3^{(k)})$$

$i = 2$

$$x_2^{(k+1)} = \frac{1}{a_{22}} (b_2 - a_{21}x_1^{(k+1)} - a_{23}x_3^{(k)})$$

$i = 3$

$$x_3^{(k+1)} = \frac{1}{a_{33}} (b_3 - a_{31}x_1^{(k+1)} - a_{32}x_2^{(k+1)})$$

Method Steps

- (a) Choose initial values $(x_1^{(0)}, x_2^{(0)}, x_3^{(0)})$
- (b) Apply the iteration using the Formula
- (c) Check if $\max(|x_i^{(k+1)} - x_i^{(k)}|) < \epsilon$

Ex:

$$4x_1 - x_2 + x_3 = 7$$

$$-2x_1 + 5x_2 - x_3 = -8$$

$$x_1 + x_2 + 3x_3 = 6$$

$$x_1^{(0)} = 0, x_2^{(0)} = 0, x_3^{(0)} = 0$$

$$\epsilon = 0.01$$

Sol:

$$\begin{bmatrix} 4 & -1 & 1 \\ -2 & 5 & -1 \\ 1 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ -8 \\ 6 \end{bmatrix}$$

$$x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right)$$

$$k = 0$$

$$i = 1$$

$$x_1^{(1)} = \frac{1}{a_{11}} (b_1 - a_{12}x_2^{(0)} - a_{13}x_3^{(0)})$$

$$x_1^{(1)} = \frac{1}{4} (7 - (-1 \times 0) - (1 \times 0)) = \frac{7}{4} \Rightarrow x_1^{(1)} = 1.75$$

$$i = 2$$

$$x_2^{(1)} = \frac{1}{a_{22}} (b_2 - a_{21}x_1^{(1)} - a_{23}x_3^{(0)})$$

$$x_2^{(1)} = \frac{1}{5} (-8 - (-2 \times 1.75) - (-1 \times 0)) = -\frac{9}{10} \Rightarrow x_2^{(1)} = -0.9$$

$$i = 3$$

$$x_3^{(1)} = \frac{1}{a_{33}} (b_3 - a_{31}x_1^{(1)} - a_{32}x_2^{(1)})$$

$$x_3^{(1)} = \frac{1}{3} (6 - (1 \times 1.75) - (1 \times -0.9)) = \frac{5.15}{3} \Rightarrow x_3^{(1)} = 1.71$$

$$x_1^{(1)} = 1.75, x_2^{(1)} = -0.9, x_3^{(1)} = 1.71$$

$$|x_1^{(1)} - x_1^{(0)}| = |1.75 - 0| = 1.75 > \epsilon$$

$$k = 1$$

$$i = 1$$

$$x_1^{(2)} = \frac{1}{a_{11}}(b_1 - a_{12}x_2^{(1)} - a_{13}x_3^{(1)})$$

$$x_1^{(2)} = \frac{1}{4}(7 - (-1 \times -0.9) - (1 \times 1.71)) \Rightarrow x_1^{(2)} = 1.0975$$

$$i = 2$$

$$x_2^{(2)} = \frac{1}{a_{22}}(b_2 - a_{21}x_1^{(2)} - a_{23}x_3^{(1)})$$

$$x_2^{(2)} = \frac{1}{5}(-8 - (-2 \times 1.0975) - (-1 \times 1.71)) \Rightarrow x_2^{(2)} = -0.819$$

$$i = 3$$

$$x_3^{(2)} = \frac{1}{a_{33}}(b_3 - a_{31}x_1^{(2)} - a_{32}x_2^{(2)})$$

$$x_3^{(2)} = \frac{1}{3}(6 - (1 \times 1.0975) - (1 \times -0.819)) \Rightarrow x_3^{(2)} = 1.9071$$

$$x_1^{(2)} = 1.0975, x_2^{(2)} = -0.819, x_3^{(2)} = 1.9071$$

$$\left| x_1^{(2)} - x_1^{(1)} \right| = |1.0975 - 1.75| = 0.6525 > \epsilon$$

$$k = 2$$

$$i = 1$$

$$x_1^{(3)} = \frac{1}{a_{11}}(b_1 - a_{12}x_2^{(2)} - a_{13}x_3^{(2)})$$

$$x_1^{(3)} = \frac{1}{4}(7 - (-1 \times -0.819) - (1 \times 1.9071)) \Rightarrow x_1^{(3)} = 1.0684$$

$$i = 2$$

$$x_2^{(3)} = \frac{1}{a_{22}}(b_2 - a_{21}x_1^{(3)} - a_{23}x_3^{(2)})$$

$$x_2^{(3)} = \frac{1}{5}(-8 - (-2 \times 1.0684) - (-1 \times 1.9071)) \Rightarrow x_2^{(3)} = -0.79122$$

$$i = 3$$

$$x_3^{(3)} = \frac{1}{a_{33}}(b_3 - a_{31}x_1^{(3)} - a_{32}x_2^{(3)})$$

$$x_3^{(3)} = \frac{1}{3}(6 - (1 \times 1.0684) - (1 \times -0.79122)) \Rightarrow x_3^{(3)} = 1.9076$$

$$x_1^{(3)} = 1.0684, x_2^{(3)} = -0.79122, x_3^{(3)} = 1.9076$$

$$\left| x_1^{(3)} - x_1^{(2)} \right| = |1.0684 - 1.0975| = 0.0291 > \epsilon$$

$$k = 3$$

$$i = 1$$

$$x_1^{(4)} = \frac{1}{a_{11}}(b_1 - a_{12}x_2^{(3)} - a_{13}x_3^{(3)})$$

$$x_1^{(4)} = \frac{1}{4}(7 - (-1 \times -0.79122) - (1 \times 1.9076)) \Rightarrow x_1^{(4)} = 1.0752$$

$$i = 2$$

$$x_2^{(4)} = \frac{1}{a_{22}}(b_2 - a_{21}x_1^{(4)} - a_{23}x_3^{(3)})$$

$$x_2^{(4)} = \frac{1}{5}(-8 - (-2 \times 1.0752) - (-1 \times 1.9076)) \Rightarrow x_2^{(4)} = -0.7884$$

$$i = 3$$

$$x_3^{(4)} = \frac{1}{a_{33}}(b_3 - a_{31}x_1^{(4)} - a_{32}x_2^{(4)})$$

$$x_3^{(4)} = \frac{1}{3}(6 - (1 \times 1.0752) - (1 \times -0.7884)) \Rightarrow x_3^{(4)} = 1.9044$$

$$x_1^{(4)} = 1.0752, x_2^{(4)} = -0.7884, x_3^{(4)} = 1.9044$$

$$|x_1^{(4)} - x_1^{(3)}| = |1.0752 - 1.0684| = 0.0068 < \epsilon$$

$$|x_2^{(4)} - x_2^{(3)}| = |-0.7884 - (-0.79122)| = 0.00282 < \epsilon$$

$$|x_3^{(4)} - x_3^{(3)}| = |1.9044 - 1.9076| = 0.0032 < \epsilon$$

So the solution set is: $x_1^{(4)} = 1.0752, x_2^{(4)} = -0.7884, x_3^{(4)} = 1.9044$

3. textbfNewton-Raphson method

The general form of the method is

$$x_{(n+1)} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Where:

$$n = 0, 1, 2, 3, \dots$$

$$f(x) = 0 \text{ at } [a, b]$$

x_0 : Defined the value $x_{(n+1)}$ is a root of the function $f(x) = 0$ if and only if it satisfies the following

$$|x_{(n+1)} - x_n| \leq \epsilon, \text{ Where } \epsilon \text{ is a small value}$$

Remark: If x_0 is not defined it can be calculated using the formula Below

$$x_0 = \frac{a+b}{2}$$

EX: using Newton-Raphson method to find the root of the following function within $[0, 1]$

$$x_0 = 0, x = e^{-x}, \epsilon = 0.005$$

Sol:

$$f(x) = 0, x = e^{-x} \xrightarrow{\text{move } e^{-x}} f(x) = x - e^{-x}$$

Now find derivative of $f(x)$

$$f'(x) = 1 + e^{-x}$$

$$x(n+1) = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$n = 0$$

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$x_1 = 0 - \frac{f(0)}{f'(0)} \Rightarrow x_1 = 0 - \frac{0 - e^0}{1 + e^0} \Rightarrow x_1 = \frac{1}{2} = 0.5$$

$$|x_{n+1} - x_n| \Rightarrow |x_1 - x_0| \Rightarrow |0.5 - 0| = 0.5 > \epsilon$$

$$n = 1$$

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$x_2 = 0.5 - \frac{f(0.5)}{f'(0.5)} \Rightarrow x_2 = 0.5 - \frac{0.5 - e^{-0.5}}{1 + e^{-0.5}} \Rightarrow x_2 = 0.566$$

$$|x_2 - x_1| \Rightarrow |0.566 - 0.5| = 0.066 > \epsilon$$

$$n = 2$$

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)}$$

$$x_3 = 0.566 - \frac{f(0.566)}{f'(0.566)} \Rightarrow x_3 = 0.566 - \frac{0.566 - e^{-0.566}}{1 + e^{-0.566}} \Rightarrow x_3 = 0.567$$

$$|x_3 - x_2| \Rightarrow |0.567 - 0.566| = 0.001 < \epsilon$$

$x_3 = 0.567$ is the root of $f(x) = x - e^{-x}$

Newton's method to find the n_{th} roots To find \sqrt{B} , we can follow the function below

$$(a) \ x = \sqrt{B} \Rightarrow x^2 = B \Rightarrow x^2 - B = 0 \Rightarrow \boxed{f(x) = x^2 - B}$$

$$(b) \ \text{Now } x = \pm\sqrt{B}$$

$$(c) \ f'(x) = 2x$$

$$(d) \ x = g(x) = x - \frac{f(x)}{f'(x)} \Rightarrow x - \frac{x^2 - B}{2x} = \frac{x^2 + B}{2x} \Rightarrow \frac{x + \frac{B}{x}}{2}$$

$$(e) \ \text{Now we can use the formula: } r_{(n+1)} = \frac{r_n + \frac{B}{r_n}}{2}$$

EX: Find $\sqrt{3}$ with $r_0 = 1$

Sol:

$$r_{(n+1)} = \frac{r_n + \frac{B}{r_n}}{2}, \quad B = 3$$

$$n = 0$$

$$r_1 = \frac{r_0 + \frac{B}{r_0}}{2}$$

$$r_1 = \frac{1 + \frac{3}{1}}{2} \Rightarrow r_1 = \frac{4}{2} \Rightarrow r_1 = 2$$

$$n = 1$$

$$r_2 = \frac{r_1 + \frac{B}{r_1}}{2}$$

$$r_2 = \frac{2 + \frac{3}{2}}{2} \Rightarrow r_2 = \frac{7}{4} \Rightarrow r_2 = 1.75$$

$$n = 2$$

$$r_3 = \frac{r_2 + \frac{B}{r_2}}{2}$$

$$r_3 = \frac{1.75 + \frac{3}{1.75}}{2} \Rightarrow r_3 = \frac{97}{56} \Rightarrow r_3 = 1.732142857$$

$$n = 3$$

$$r_4 = \frac{r_3 + \frac{B}{r_3}}{2}$$

$$r_4 = \frac{1.732142857 + \frac{3}{1.732142857}}{2} \Rightarrow r_4 = 1.73205$$

$$n = 4$$

$$r_5 = \frac{r_4 + \frac{B}{r_4}}{2}$$

$$r_5 = \frac{1.73205 + \frac{3}{1.73205}}{2} \Rightarrow r_5 = 1.73205$$

We will stop if our last two results matches

n	r_{n+1}
0	2
1	1.75
2	1.732142857
3	1.73205
4	1.73205

4. Fixed point method

To find the root of an equation using this method, we can follow the following steps:

- Putting the function $f(x)$ in the form of $x = g(x)$
- Make sure $|g'(x_0)| < 1$, If not we need to chose another form of the $g(x)$
- Find x_{n+1} , using the formula $x_{n+1} = g(x_n)$, $n = 0, 1, 2, \dots$
- The value x_{n+1} is a root of the function $f(x) = 0$ if and only if $|x_{n+1} - x_n| \leq \epsilon$, Where ϵ is a small value

EX: Use fixed point method to find the root of $f(x) = x^2 - x - 3 = 0$ within $[1, 2]$ and $\epsilon = 0.01$

Sol:

$$x_0 = \frac{a+b}{2} \Rightarrow x_0 = \frac{1+2}{2} \Rightarrow x_0 = \frac{3}{2} \Rightarrow x_0 = 1.5$$

$$f(x) = x^2 - x - 3 = 0$$

$$x^2 - x - 3 = 0 \xrightarrow{\text{move } x} x = x^2 - 3$$

$$x = g(x), \quad g'(x) = 2x$$

$$|g'(x_0)| \Rightarrow |g'(1.5)| \Rightarrow |2 \times 1.5| = 3 > 1$$

We need to find another form of $g(x)$

$$x^2 - x - 3 = 0 \Rightarrow x^2 = x + 3 \Rightarrow x = \sqrt{x+3} = g(x)$$

$$g'(x) = \frac{1}{2\sqrt{x+3}}$$

$$|g'(x_0)| \Rightarrow \left| \frac{1}{2\sqrt{1.5+3}} \right| = 0.23 < 1$$

$$x_{n+1} = g(x_n)$$

$$n = 0$$

$$x_1 = g(x_0)$$

$$x_1 = g(1.5) = \sqrt{1.5 + 3} = \sqrt{4.5} = 2.12 \Rightarrow x_1 = 2.12$$

$$|x_1 - x_0| = |2.12 - 1.5| = 0.62 > \epsilon$$

$$n = 1$$

$$x_2 = g(x_1)$$

$$x_2 = g(2.12) = \sqrt{2.12 + 3} = \sqrt{5.12} = 2.26 \Rightarrow x_2 = 2.26$$

$$|x_2 - x_1| = |2.26 - 2.12| = 0.14 > \epsilon$$

$$n = 2$$

$$x_3 = g(x_2)$$

$$x_3 = g(2.26) = \sqrt{2.26 + 3} = \sqrt{5.26} = 2.29 \Rightarrow x_3 = 2.29$$

$$|x_3 - x_2| = |2.29 - 2.26| = 0.03 > \epsilon$$

$$n = 3$$

$$x_4 = g(x_3)$$

$$x_4 = g(2.29) = \sqrt{2.29 + 3} = \sqrt{5.29} = 2.3 \Rightarrow x_4 = 2.3$$

$$|x_4 - x_3| = |2.3 - 2.29| = 0.01 = \epsilon$$

So $x_4 = 2.3$ is the root of $f(x)$

Interpolation theory: Used to find the missing data (values) of the existing data based on the relationships between the data (values)

Lagrang interpolation method: In a series of x-values, This method can be used to find the value of $f(x)$, this is one of the simplest numerical methods that can be show by the following data

x	x_0	x_1	\dots	x_n
$f(x)$	$f(x_0)$	$f(x_1)$	\dots	$f(x_n)$

The Lagrang method for the above data will be

$$f(x^*) = f(x_0)L_0x^* + f(x_1)L_1x^* + f(x_2)L_2x^*$$

Where:

x^* : The value we want to find

$$L_0x^* = \frac{(x^*-x_1)(x^*-x_2)}{(x_0-x_1)(x_0-x_2)} \times \frac{(x^*-x_0)(x^*-x_2)}{(x_1-x_0)(x_1-x_2)} \times \frac{(x^*-x_0)(x^*-x_1)}{(x_2-x_0)(x_2-x_1)}$$

So the general form is

$$f(x^*) = \sum_{j=0}^n f(x_j) \prod_{\substack{i=0 \\ i \neq j}}^n \frac{x^* - x_i}{x_j - x_i}$$

EX: By Lagrang formula, find $f(3)$ and $f(5)$ using the following set

x	0	1	2	4
$f(x)$	1	1	2	5

Sol

$$f(x^*) = \sum_{j=0}^n f(x_j) \prod_{\substack{i=0 \\ i \neq j}}^n \frac{x^* - x_i}{x_j - x_i}$$

$n = 3$ because we have 4 values in the table

$$\begin{aligned} f(x^*) = & f(x_0) \frac{(x^* - x_1)(x^* - x_2)(x^* - x_3)}{(x_0 - x_1)(x_0 - x_2)(x_0 - x_3)} \quad j = 0, i = 1, 2, 3 \\ & + f(x_1) \frac{(x^* - x_0)(x^* - x_2)(x^* - x_3)}{(x_1 - x_0)(x_1 - x_2)(x_1 - x_3)} \quad j = 1, i = 0, 2, 3 \\ & + f(x_2) \frac{(x^* - x_0)(x^* - x_1)(x^* - x_3)}{(x_2 - x_0)(x_2 - x_1)(x_2 - x_3)} \quad j = 2, i = 0, 1, 3 \\ & + f(x_3) \frac{(x^* - x_0)(x^* - x_1)(x^* - x_2)}{(x_3 - x_0)(x_3 - x_1)(x_3 - x_2)} \quad j = 3, i = 0, 1, 2 \end{aligned}$$

$$\begin{aligned}
f(3) &= f(0) \frac{(3-1)(3-2)(3-4)}{(0-1)(0-2)(0-4)} \\
&\quad + f(1) \frac{(3-0)(3-2)(3-4)}{(1-0)(1-2)(1-4)} \\
&\quad + f(2) \frac{(3-0)(3-1)(3-4)}{(2-0)(2-1)(2-4)} \\
&\quad + f(4) \frac{(3-0)(3-1)(3-2)}{(4-0)(4-1)(4-2)}
\end{aligned}$$

$$f(3) = 1 \times \frac{-2}{-8} + 1 \times \frac{-3}{3} + 2 \times \frac{-6}{-4} + 5 \times \frac{6}{24} = 3 \Rightarrow f(3) = 3$$

$$\begin{aligned}
f(5) &= f(0) \frac{(5-1)(5-2)(5-4)}{(0-1)(0-2)(0-4)} \\
&\quad + f(1) \frac{(5-0)(5-2)(5-4)}{(1-0)(1-2)(1-4)} \\
&\quad + f(2) \frac{(5-0)(5-1)(5-4)}{(2-0)(2-1)(2-4)} \\
&\quad + f(4) \frac{(5-0)(5-1)(5-2)}{(4-0)(4-1)(4-2)}
\end{aligned}$$

$$f(5) = 1 \times \frac{12}{-8} + 1 \times \frac{15}{3} + 2 \times \frac{20}{-4} + 5 \times \frac{60}{24} = 6 \Rightarrow f(5) = 6$$

Inverse Lagrang: used to find x from y^* as follows

$$x^* = \sum_{j=0}^n x_j \prod_{\substack{i=0 \\ i \neq j}}^n$$

Lets say $n = 2$

$$x_* = x_0 \frac{(y^* - y_1)(y^* - y_2)}{(y_0 - y_1)(y_0 - y_2)} + x_1 \frac{(y^* - y_0)(y^* - y_2)}{(y_1 - y_0)(y_1 - y_2)} + x_2 \frac{(y^* - y_0)(y^* - y_1)}{(y_2 - y_0)(y_2 - y_1)}$$

EX: Find the value of x^* when $y^* = 0.2703$ for the following data

x	1.6	2.9	4.8
y	0.625	0.3448	0.2083

Sol:

$$x^* = \sum_{j=0}^n x_j \prod_{\substack{i=0 \\ i \neq j}}^n$$

$n = 2$

$$x_* = x_0 \frac{(y^* - y_1)(y^* - y_2)}{(y_0 - y_1)(y_0 - y_2)} + x_1 \frac{(y^* - y_0)(y^* - y_2)}{(y_1 - y_0)(y_1 - y_2)} + x_2 \frac{(y^* - y_0)(y^* - y_1)}{(y_2 - y_0)(y_2 - y_1)}$$

$$\begin{aligned}
x_* &= 1.6 \frac{(0.2703 - 0.3448)(0.2703 - 0.2083)}{(0.625 - 0.3448)(0.625 - 0.2083)} \\
&\quad + 2.9 \frac{(0.2703 - 0.625)(0.2703 - 0.2083)}{(0.3448 - 0.625)(0.3448 - 0.2083)} \\
&\quad + 4.8 \frac{(0.2703 - 0.625)(0.2703 - 0.3448)}{(0.2083 - 0.625)(0.2083 - 0.3448)} \\
x_* &= 1.6 \frac{-0.004619}{0.11675934} + 2.9 \frac{-0.0219914}{-0.0382473} + 4.8 \frac{0.02642515}{0.05687955} \Rightarrow x^* = 3.834131631
\end{aligned}$$