

A READER'S COMPANION · PHARMACOLOGY & EVERYDAY SAFETY

THE MEDICINE CABINET

An illustrated guide to what medicines are, how they
work inside the body, and how to use them wisely &
safely at home — expanded second edition.

EST. GUIDE NO. 07 — EXPANDED GENERAL EDITION FOR EDUCATIONAL USE

What Is a Medicine?

INDICATION — Understanding the basic idea behind every tablet, syrup and injection.

At its simplest, a medicine is a substance used to prevent, treat, or manage a disease, or to relieve a symptom — anything from a headache to a serious infection.

Every medicine is built around one or more active ingredients: the chemical compounds that actually produce the effect on the body. Surrounding the active ingredient are excipients — inactive substances such as fillers, coatings, flavourings, and preservatives. Excipients do not treat the illness, but they hold the tablet together, help it dissolve at the right time, make a syrup taste pleasant, or keep the product stable on a shelf for years.

Medicine, Food, and Supplements

People sometimes confuse medicines with food supplements or vitamins. The difference lies in purpose and evidence: a medicine is developed to produce a specific, measurable effect on the body and must prove this through clinical testing before it can be sold. A food supplement, by contrast, is meant to add to the diet — such as extra vitamins — and is not required to prove it treats a disease.

IN SHORT

A true medicine has three things: a known active ingredient, a proven effect, and clear instructions for how, when, and in what amount it should be used.

Why Medicines Are Regulated

Because medicines act on the body in powerful ways, nearly every country has a regulatory authority — such as a Ministry of Health or a national medicines agency — that reviews evidence before a medicine can reach a pharmacy shelf. Regulators check that a medicine works for its intended purpose, that its benefits outweigh its risks, and that it is manufactured to a consistent, safe quality every time.

The Anatomy of a Medicine

Every packet, bottle, or blister strip is really a small system of parts working together, each with its own job. Understanding these parts makes any label easier to read.

Component	Role
Active ingredient	The molecule responsible for the medicine's effect on the body.
Excipient / filler	Bulks out a tablet so it is a practical, handle-able size.
Coating	Protects the stomach, masks taste, or controls where the tablet dissolves.
Preservative	Stops bacteria or mould from growing in liquids over time.
Binder	Holds powder together so a tablet does not crumble in the packet.

How a Medicine Is Named

Most medicines carry two names. The generic name identifies the active ingredient itself, written in plain scientific language and used everywhere in the world. The brand name is a trademarked, marketed title chosen by the company that sells it. Paracetamol, for example, is sold worldwide under dozens of different brand names, even though the active ingredient — and its effect on the body — is identical.

From Idea to Pharmacy Shelf

Before a new medicine ever reaches a patient, it typically passes through laboratory research, animal studies, and then several phases of human clinical trials — first checking that it is safe in a small group, then that it works as intended in a larger one, and finally comparing it against existing treatments. This process commonly takes a decade or more and represents one of the most heavily regulated activities in modern industry.

DID YOU KNOW?

Only a small fraction of chemical compounds that begin laboratory testing ever reach a pharmacy shelf — most are set aside for lacking effect, or for side effects that outweigh their benefit.

A Short History of Medicine

INDICATION — From garden herbs to the modern pharmacy.

Long before laboratories existed, healing came from plants, minerals, and careful observation passed down through generations.

Ancient Egyptian, Chinese, and Greek healers catalogued hundreds of plant remedies — willow bark for pain, moulds for wound care, poppy for sleep. Much of this knowledge was recorded by early physicians such as Hippocrates, whose emphasis on observing patients rather than relying on superstition laid a foundation for medicine as we know it.

Turning Plants into Pure Chemistry

The nineteenth century brought a turning point: chemists learned to isolate the exact active compound inside a plant. Willow bark's pain-relieving ingredient was purified and refined into aspirin — one of the first modern manufactured medicines, and still one of the most widely used in the world today.

The Age of Antibiotics and Vaccines

In 1928, the accidental discovery of a mould that killed bacteria in a laboratory dish led to penicillin, the first true antibiotic — a discovery that transformed infections that once killed millions into treatable conditions. Around the same period, vaccination — introduced a century earlier against smallpox — expanded to protect against diseases such as polio, measles, and tetanus, preventing illness before it ever begins.

MILESTONE TIMELINE

1800s — Isolation of pure active compounds from plants (morphine, quinine, aspirin).

1920s — Discovery of penicillin, the first antibiotic.

1950s — Mass production of vaccines and the first oral contraceptives.

Today — Precision medicines, biologics, and treatments designed for a person's own biology.

Modern pharmaceutical science now designs medicines deliberately, molecule by molecule, testing them for years in laboratories and clinical trials before they ever reach a patient — a far cry from the trial-and-error herbal remedies of the past, yet still built on the same basic goal: to relieve suffering and extend healthy life.

Medicine Around the World

Healing traditions developed independently across many regions long before modern trade or translation connected them — and several of these traditions still inform treatments used today, either directly or as inspiration for modern research.

- **Traditional Chinese Medicine** — centuries of herbal formulas and acupuncture; artemisinin, a modern antimalarial, was isolated after study of a traditional Chinese remedy.
- **Ayurveda (South Asia)** — a system built around diet, herbs, and lifestyle balance, still practised widely alongside modern medicine.
- **Islamic Golden Age medicine** — scholars such as Ibn Sina compiled vast medical encyclopaedias and pioneered clinical trial-like observation and hospital care.
- **African traditional medicine** — extensive plant-based pharmacopoeias, some of which continue to be studied for active compounds today.

Why This History Still Matters

Roughly half of all modern medicines can trace some part of their origin to a natural source — a plant, fungus, or micro-organism — later purified, modified, or used as a blueprint for a synthetic version. Understanding this history helps explain why dose still matters so much: traditional preparations varied widely in strength, which is exactly the inconsistency that modern manufacturing and regulation were built to remove.

A USEFUL DISTINCTION

"Natural" and "safe" are not the same idea. Many of the most powerful and dangerous substances known are entirely natural in origin — which is precisely why purification, testing, and correct dosing matter regardless of where a compound came from.

How Medicines Work in the Body

INDICATION — The journey from swallowing a tablet to feeling relief.

A medicine's journey through the body follows two parallel stories: what the body does to the medicine, and what the medicine does to the body.

What the Body Does to the Medicine

Scientists describe this journey with four stages, often remembered by the letters A-D-M-E:

- **Absorption** — the medicine enters the bloodstream, commonly through the stomach and intestine walls after swallowing.
- **Distribution** — blood carries the medicine throughout the body, reaching the organ or tissue it is meant to affect.
- **Metabolism** — the liver gradually breaks the medicine down into simpler substances.
- **Excretion** — the kidneys remove the leftover substances from the body, mainly through urine.

What the Medicine Does to the Body

This second story is about targets. Most medicines work by attaching to specific receptors — tiny docking points on the surface of cells — much like a key fitting a particular lock. When the medicine locks into place, it can switch a process on, slow it down, or block it entirely: a painkiller may quiet pain signals travelling to the brain, while an antihistamine blocks the receptor responsible for an allergic reaction.

WHY TIMING MATTERS

Because the body is constantly absorbing and clearing a medicine, levels in the blood rise and fall throughout the day. Taking a dose too early, too late, or skipping one can let levels fall below what is needed to work — which is exactly why directions specify "every 8 hours" or "twice daily" rather than "whenever convenient."

Dose, Response & Generics

Dose and Response

Too little of a medicine may have no effect at all; too much can turn a helpful substance into a harmful one. This relationship is called the dose-response curve, and it is the reason every medicine is tested extensively to find the smallest dose that reliably works with the fewest unwanted effects — the basis of the phrase used since the sixteenth century: "the dose makes the poison."

This is also why children are rarely given a simple fraction of an adult dose. Their bodies absorb, break down, and clear medicines differently, so paediatric doses are calculated and tested separately, often based on body weight.

"The right dose differentiates a poison from a remedy."

— PARACELSUS, 16TH-CENTURY PHYSICIAN

Generic and Brand-Name Medicines

When a company first develops a new medicine, it is sold under a protected brand name while patent laws give the manufacturer exclusive rights to sell it. Once that patent expires, other manufacturers may produce the same active ingredient as a generic medicine. Regulators require generics to match the original in strength, quality, and the way the body absorbs them — so a generic is expected to work the same way, usually at a lower price.

Feature	Brand-Name	Generic
Active ingredient	Original, patented	Identical
Appearance	Company's own design	May differ in shape/colour
Price	Higher	Typically lower
Effectiveness	Proven in trials	Proven equivalent

Forms of Medicine

INDICATION — Why the same medicine can come as a pill, a syrup, or a patch.

The active ingredient is only half the story — the form it is delivered in decides how quickly it works, where in the body it acts, and how easy it is to use.

Taken by Mouth

- **Tablets** — compressed powder, often coated to protect the stomach or delay release.
- **Capsules** — a soft or hard shell holding powder, liquid, or granules.
- **Syrups & suspensions** — liquid forms, useful for children or anyone who cannot swallow tablets.
- **Chewables & dissolvables** — break down in the mouth for faster, easier dosing.

Beyond the Mouth

- **Injections** — delivered directly into a vein, muscle, or under the skin for fast or precise action.
- **Topical creams & patches** — absorbed through the skin, often for local or slow, steady effect.
- **Inhalers** — deliver medicine straight to the lungs, ideal for asthma treatment.
- **Drops & suppositories** — used for the eyes, ears, nose, or when swallowing is not possible.

WHY FORM MATTERS

An injection can act within seconds because it bypasses digestion entirely, while a tablet may take thirty to sixty minutes to be absorbed. An inhaler delivers a tiny, targeted dose straight to the lungs, avoiding the larger dose that would be needed if the same medicine were swallowed. Choosing the right form is part of how a treatment is designed to work safely and effectively.

A Word on Route and Speed

Recognising a form at a glance helps you follow directions correctly — a "drop" and a "spoonful" are never interchangeable.

Route	Typical Onset	Common Use
Intravenous injection	Seconds to minutes	Emergencies, hospital care
Intramuscular injection	10–20 minutes	Vaccines, some antibiotics
Oral (tablet/capsule)	30–60 minutes	Everyday, long-term treatment
Inhaled	Minutes	Asthma, respiratory relief
Topical / transdermal	Hours	Local pain, steady slow release

Choosing Between Forms

When a medicine is available in more than one form — for example, a pain reliever sold as both a tablet and a fast-dissolving powder — the choice often comes down to how quickly relief is needed, whether the person can swallow easily, and personal preference. A pharmacist can advise when one form offers a genuine advantage over another for a particular situation, such as a child, an older adult, or someone recovering from surgery.

A NOTE ON CRUSHING TABLETS

Not every tablet can be safely crushed or opened, even to make swallowing easier. Some coatings are designed to control exactly where or how fast a medicine releases, and breaking that coating can release the full dose too quickly. Always ask a pharmacist before crushing, splitting, or opening a capsule.

Classes of Medicine

INDICATION — A field guide to the major families of medicine.

Rather than memorising thousands of individual products, it helps to think in families — groups of medicines that share a purpose and a way of working.

Class	General Purpose
Analgesics	Relieve pain, from mild headaches to post-surgical discomfort.
Antibiotics	Treat infections caused by bacteria; ineffective against viruses.
Antivirals	Slow or stop viruses from multiplying in the body.
Antihistamines	Ease allergy symptoms such as sneezing, itching, and rashes.
Antihypertensives	Help manage and lower high blood pressure.
Antidiabetic agents	Help the body regulate blood sugar levels.
Antidepressants	Support the brain's chemical balance in mood disorders.
Vaccines	Train the immune system to recognise a disease before exposure.
Vitamins & minerals	Correct or prevent a nutritional shortfall in the diet.

One Symptom, Many Possible Causes

A single symptom, such as a cough, may call for entirely different classes of medicine depending on its cause — an antihistamine for an allergy, an antibiotic for a bacterial chest infection, or simply a soothing syrup for a dry throat. This is precisely why self-diagnosis can mislead: the right class of medicine depends on identifying the right underlying cause, which is a task for a qualified professional.

A COMMON MISUNDERSTANDING

Antibiotics treat bacteria, not viruses. Taking an antibiotic for a cold or flu will not speed recovery, and unnecessary use contributes to a serious global problem known as antibiotic resistance, where bacteria evolve to survive medicines that once controlled them.

Working Together, Working Against

Medicines from different classes are often prescribed together deliberately — for example, pairing a pain reliever with an anti-inflammatory after an injury. But combinations can also work against each other or amplify side effects, which is why a doctor or pharmacist reviews the full list of medicines a person takes before adding something new.

Biologics and Precision Medicines

A newer family of medicines, called biologics, is made from living cells rather than pure chemistry — used for conditions such as certain cancers, autoimmune diseases, and some forms of diabetes (insulin itself is a biologic). These medicines are often more complex to manufacture, store, and administer, frequently requiring refrigeration and injection rather than a simple tablet.

Combination Products

Some everyday medicines already combine more than one active ingredient in a single tablet — a cold remedy, for instance, may pair a decongestant with a pain reliever. This is convenient, but it also means it is easy to accidentally double up on an ingredient by taking two different combination products at once. Checking the active ingredient list, not just the brand name, is the safest habit.

REMEMBER

The class of a medicine tells you its general purpose — but only a healthcare professional can match the right class, and the right specific medicine within it, to an individual's diagnosis, age, and health history.

Prescription vs. Over-the-Counter

INDICATION — Why some medicines sit behind the counter, and some don't.

Every medicine sold legally falls into one of two broad categories, decided by how much oversight it needs to be used safely.

Prescription (Rx)

Requires a licensed prescriber's authorisation. Typically reserved for medicines that are more potent, have higher risk of side effects,

Over-the-Counter (OTC)

Why the Line Sometimes Moves

A medicine can shift from prescription-only to over-the-counter once years of real-world use confirm it is safe enough for people to manage without direct supervision — often at a lower strength than the original prescription version. This is a formal regulatory decision, not simply a matter of a shop deciding to stock it.

A Third Category in Some Countries

Several regulators also recognise a middle tier — sometimes called "pharmacist-only" or "behind-the-counter" medicines — that do not require a prescription but can only be sold after a short conversation with a pharmacist, allowing a trained professional to check for obvious risks before it leaves the shop.

THE PHARMACIST'S ROLE

Pharmacists are trained medicine experts, not just dispensers. Even for over-the-counter products, asking a pharmacist a question — about a possible interaction, a pregnancy, or a child's age — takes a moment and can prevent a real problem.

Reading the Label

INDICATION — Five lines of small print that matter more than they look.

A medicine label is a compressed instruction manual. Learning to read it properly is one of the most useful habits in everyday health.

- **Active ingredient & strength** — what the medicine actually is, and how concentrated each dose is (for example, 500 mg).
- **Directions for use** — how much to take, how often, and for how long; sometimes with or without food.
- **Warnings & precautions** — conditions, ages, or combinations (such as alcohol) to avoid.
- **Expiry date & batch number** — when the medicine should no longer be used, and its manufacturing trace for quality control.
- **Storage instructions** — temperature and light conditions the medicine needs to remain effective.

A SIMPLE HABIT

Read the label every time, even for a medicine you have used before — manufacturers occasionally update directions, strengths, or warnings between purchases.

Following Directions Precisely

Directions exist because they reflect how the medicine was actually tested. Stopping a course of antibiotics early because symptoms have improved, for instance, can leave surviving bacteria behind — some of which may be more resistant to treatment next time. Doubling a dose to "catch up" after a missed one can push levels beyond what the body was tested to handle safely.

Measuring Doses Correctly

Liquid medicines should always be measured with the dosing device provided — an oral syringe, cup, or spoon calibrated for that specific product. An ordinary kitchen teaspoon can vary in volume by nearly a third, which is more than enough to under- or over-dose a young child.

IF IN DOUBT

If a label's directions seem unclear, or conflict with what a doctor said aloud, call the pharmacy before taking the medicine — not after.

Understanding Missed-Dose Instructions

Many labels include a short line for missed doses — commonly something like "take as soon as you remember, unless it is nearly time for the next dose." This wording exists to prevent two doses being taken too close together. If a label does not address missed doses at all, a pharmacist can explain what to do for that specific medicine, since the right answer varies between products.

When a Label Is Hard to Read

Small print, glare, or a language barrier can all make a label difficult to read safely. Most pharmacies can provide large-print labels, translated patient information leaflets, or simply explain the directions aloud — it is always worth asking rather than guessing.

Side Effects, Interactions & Allergies

INDICATION — Understanding the body's unwanted responses.

No medicine acts on only one target in the body — which is why almost every medicine carries a list of possible side effects alongside its intended benefit.

Common vs. Serious Side Effects

Most side effects are mild and temporary — drowsiness, a dry mouth, mild nausea — and often fade as the body adjusts. Serious side effects are rarer but require immediate attention: difficulty breathing, swelling of the face or throat, severe rash, or an irregular heartbeat.

Interactions: Medicine Meeting Medicine

An interaction occurs when one substance changes how another behaves in the body — sometimes strengthening its effect, sometimes weakening it, and sometimes creating an entirely new risk. Interactions are not limited to other medicines: grapefruit juice, alcohol, herbal remedies, and even certain foods can all interact with specific medicines.

TELL YOUR PROVIDER EVERYTHING

Every doctor or pharmacist should know the complete list of what you take — prescription medicines, over-the-counter products, vitamins, and herbal supplements — before adding anything new.

Recognising an Allergic Reaction

An allergy is different from a side effect: it is the immune system mistakenly treating a harmless ingredient as a threat. Mild signs include a rash, itching, or hives. A severe reaction, called anaphylaxis, can involve swelling of the throat, difficulty breathing, or a sudden drop in blood pressure, and is a medical emergency requiring immediate care.

SEEK EMERGENCY HELP IMMEDIATELY IF YOU NOTICE

- Swelling of the lips, tongue, or throat
- Sudden difficulty breathing or wheezing
- Widespread hives with dizziness or fainting

Keeping a Personal Record

Once an allergy or serious reaction to a medicine is identified, it is worth recording clearly — on a card, a phone note, or a medical alert bracelet — so that any future doctor, dentist, or pharmacist can see it immediately, even in an emergency when you cannot explain it yourself.

Side Effect or Coincidence?

Not every symptom that appears after starting a medicine is caused by it — illness itself, stress, or an unrelated condition can happen to occur around the same time. Keeping a simple note of what changed and when helps a doctor tell the difference, rather than guessing after the fact.

Safe Storage & Disposal

INDICATION — Protecting a medicine's potency, and protecting curious hands.

A medicine's usefulness does not end when it leaves the pharmacy — how it is stored, and eventually discarded, matters just as much as how it is taken.

Storing Medicines Correctly

- **Cool and dry** — heat and humidity (such as a bathroom cabinet) can quietly break down active ingredients.
- **Original packaging** — keeps the label, batch number, and expiry date attached to the product.
- **Out of sight and reach of children** — a locked cabinet or high shelf, never a bedside table or open bag.
- **Refrigeration when specified** — certain liquids, insulin, and biologics must stay refrigerated to remain effective.

ON EXPIRY DATES

An expired medicine is not necessarily instantly dangerous, but its strength can no longer be guaranteed — meaning it may simply stop working as intended, which can be just as risky for a serious condition. When in doubt, replace it.

Disposing of Unused Medicine

Unused or expired medicines should never simply go in a household bin unprotected, nor be flushed down a toilet or sink unless a label specifically instructs it — both can allow the medicine to be recovered by others or to enter the water supply. Many pharmacies and health clinics offer a take-back collection point designed specifically for safe disposal.

A SIMPLE RULE

If your local area has no take-back programme, mixing the medicine with an unappealing substance (such as used coffee grounds), sealing it in a bag, and placing it in household waste is generally safer than leaving it loose or flushing it — but a pharmacist's advice is always the best first step.

Never Share Medicine

A medicine prescribed for one person is matched to that person's diagnosis, weight, age, and health history. Sharing it with a friend or family member with similar symptoms can mean the wrong treatment, the wrong dose, or a dangerous interaction with something else they are taking.

Cleaning Out a Cabinet Safely

A useful habit is a twice-yearly cabinet check: gather every bottle and box, check expiry dates, and set aside anything unused, unlabelled, or expired for proper disposal. Never pour leftover pills into a single unlabelled container — if a mistake ever needs untangling, mixed, unlabelled medicine is far harder to identify safely.

CHAPTER 10

Everyday Safety for Families

INDICATION — Small habits that quietly prevent big mistakes.

Most medicine mishaps at home are not dramatic — they are small, avoidable slips in a busy routine. A few simple habits close nearly all of them.

- Keep one written list of every medicine, supplement, and dose taken by each family member, updated after every doctor's visit.
- Use the device provided for liquids — never a kitchen spoon — and double-check units (mL versus teaspoons).
- Store up and away, ideally in a cabinet with a child-resistant latch, and relock it immediately after use.
- Never call medicine "candy" to encourage a child to take it — it can encourage them to seek it out unsupervised.
- Check with a provider before mixing a new medicine with alcohol, herbal products, or an existing prescription.
- Ask before, not after — during pregnancy, breastfeeding, or for a young child, confirm suitability with a doctor or pharmacist first.
- Travel prepared — carry medicines in original, labelled packaging, especially across borders.

IN CASE OF ACCIDENTAL OVERDOSE

Contact your local poison control centre or emergency services immediately, and have the medicine packaging on hand to describe exactly what was taken.

Medicine Across Life Stages

INDICATION — Why the same medicine can act differently depending on who takes it.

A medicine's effect is never just about the substance itself — it is about the body receiving it. Age, body size, pregnancy, and other health conditions can all change how a medicine behaves, which is why guidance differs so much between life stages.

Infants and Young Children

Children are not simply small adults. Their livers and kidneys — the organs responsible for breaking down and clearing medicines — mature gradually over the first years of life, so doses are calculated specifically for a child's age and weight rather than scaled down from an adult amount. Many medicines also come in child-specific forms, such as flavoured syrups, precisely because swallowing a tablet safely is a skill that develops with age.

Older Adults

As the body ages, the liver and kidneys often process medicines more slowly, and older adults are more likely to take several medicines at once for different conditions — increasing the chance of an interaction. For this reason, doctors frequently review an older person's full medicine list together, checking whether every item is still needed and at the right strength.

Pregnancy and Breastfeeding

During pregnancy, a medicine taken by the mother can reach the developing baby, and during breastfeeding, small amounts can pass through breast milk. Some medicines are well established as safe at these times, others are avoided, and many simply have not been studied enough to say for certain — which is why checking with a doctor or pharmacist before starting, stopping, or continuing any medicine during pregnancy or breastfeeding is particularly important.

Chronic Conditions and Multiple Medicines

People managing a long-term condition — such as high blood pressure, diabetes, or a heart condition — often take several medicines every day, sometimes for years. Living well with this kind of routine is less about memory alone and more about good systems.

- A pill organiser sorted by day and time reduces the chance of a missed or doubled dose.
- A single, current medicine list — shared with every doctor involved in care — prevents conflicting prescriptions from different specialists.
- Regular medicine reviews catch medicines that may no longer be needed as a condition changes over time.
- Refilling a prescription a few days before it runs out avoids an unplanned gap in treatment.

People with Reduced Kidney or Liver Function

Because the liver and kidneys are responsible for clearing most medicines from the body, someone with reduced function in either organ may need a lower dose, a longer interval between doses, or an entirely different medicine — since a standard dose could otherwise build up in the body to unsafe levels. This is a decision made by a treating doctor, often with blood tests to guide it, not something to estimate at home.

THE THEME THROUGHOUT THIS CHAPTER

There is no such thing as a medicine that is simply "safe" or "unsafe" in isolation — safety always depends on who is taking it, and that judgement belongs to a healthcare professional who knows the full picture.

Traveling with Medicine

INDICATION — Crossing time zones, borders, and climates without losing track of a routine.

Travel can disrupt even a well-established medicine routine — new time zones, unfamiliar climates, and border rules all add small complications that are easy to plan around in advance.

Before You Leave

- Pack enough medicine for the full trip, plus a small buffer in case of delays, split across two bags in case one is lost.
- Keep medicines in their original, labelled packaging — an unlabelled pill in a travel case can be difficult to identify, and may raise questions at a border.
- Carry a copy of the prescription, or a doctor's letter, especially for injectable medicines, controlled substances, or large quantities.
- Research the destination country's rules in advance — a medicine that is routine at home can occasionally be restricted or unfamiliar elsewhere.

Crossing Time Zones

For a medicine taken at a fixed time each day, crossing several time zones raises a simple question: adjust the clock time immediately, or keep counting hours from the last dose? For most medicines a gradual shift over the first day or two is fine, but for time-sensitive medicines — insulin is a common example — a doctor or pharmacist can suggest a specific adjustment plan before departure.

KEEP IT WITH YOU

Essential medicine should travel in hand luggage, not checked baggage — checked bags can be delayed, lost, or exposed to extreme temperatures in an aircraft hold.

Heat, Cold, and Storage on the Move

A hot car, a beach bag, or a cold mountain trip can all push a medicine outside the temperature range it was tested to withstand. A small insulated pouch, kept out of direct sun and away from a car's dashboard or boot, protects both refrigerated medicines and ordinary tablets on a long journey.

Common Myths About Medicine

INDICATION — Separating everyday folk wisdom from what the evidence actually shows.

Some ideas about medicine are repeated so often that they start to feel like fact. A few of the most common are worth examining directly.

"Natural remedies can't cause interactions."

Plant-based and herbal products contain active chemical compounds just like manufactured medicines, and can interact with prescriptions in exactly the same way. St John's Wort, for example, is well documented to reduce the effectiveness of several common prescription medicines.

"If a little helps, more will help faster."

This ignores the dose-response relationship covered in Chapter 3 — beyond a certain point, additional medicine adds risk rather than additional benefit.

"Old medicine is fine as long as it looks normal."

Appearance is not a reliable guide to potency. Many medicines lose strength well before any visible change occurs, and some can form new, harmful breakdown products past their expiry date.

More Myths, Examined

"Antibiotics work on any infection."

As covered in Chapter 5, antibiotics act only on bacteria. Colds, flu, and most sore throats are caused by viruses, which antibiotics cannot touch.

"Generic medicines are lower quality."

Regulators require a generic to match the original in active ingredient, strength, and how the body absorbs it. The price difference reflects the absence of development and marketing costs, not a difference in the medicine itself.

"You can build tolerance to any medicine to make it work better."

Tolerance is real for some medicines — but it usually means a reduced response over time, the opposite of a benefit, and often signals that a dose needs to be reviewed by a doctor rather than simply increased.

"Stopping a medicine once you feel better is always fine."

For some short courses this is true, but for many — especially antibiotics, and medicines for chronic conditions like blood pressure — stopping early can allow the original problem to return, sometimes worse than before.

THE SIMPLEST TEST

If a claim about a medicine sounds surprising, simple, and universal all at once, it is worth checking with a pharmacist before acting on it — real pharmacology is rarely that tidy.

Building a Home Medicine Cabinet

INDICATION — A practical checklist for what belongs on the shelf — and what doesn't.

A well-organised medicine cabinet is less about having everything, and more about having the right basics, stored correctly, and checked regularly.

A Sensible Starting List

- A digital thermometer
- A basic pain and fever reliever appropriate for each family member's age
- Adhesive bandages, sterile gauze, and adhesive tape
- Antiseptic wipes or solution for cleaning minor wounds
- Tweezers and small first-aid scissors
- Oral rehydration sachets, useful during vomiting or diarrhoea
- Any regularly prescribed medicine, plus a small emergency supply where appropriate
- A written, up-to-date list of every medicine each family member currently takes

WHAT NOT TO STOCKPILE

Avoid keeping large quantities of any single medicine beyond what is realistically needed — it increases both the chance it expires unused and the risk if a child or visitor finds it.

Where to Keep It

Despite the name, a bathroom cabinet is often one of the worst places to store medicine — the heat and humidity from showers and baths can quietly degrade active ingredients faster than in almost any other room.

Location	Verdict
Bathroom cabinet	Avoid — heat and humidity reduce potency over time.
Bedside table / open bag	Avoid — accessible to children and easy to lose track of.
High kitchen cupboard, away from the stove	Good — cool, dry, and out of reach.
Locked cabinet or box	Best — combines correct climate with restricted access.
Refrigerator (only if specified)	Correct only for medicines labelled for refrigeration.

A Seasonal Check-In

Tying a medicine cabinet check to an existing yearly habit — such as changing smoke alarm batteries, or a change of season — makes it far more likely to actually happen. Five minutes twice a year is usually enough to catch expired items, confirm the family medicine list is current, and make sure child-resistant latches still work properly.

Understanding Label Abbreviations

INDICATION — The shorthand pharmacists and doctors use, decoded.

Prescription labels often carry short abbreviations, many rooted in Latin, that compress detailed instructions into very little space. Recognising the common ones makes any label far easier to follow — though the printed instruction on the label, in full words, should always be the final word on how to take a medicine.

Abbreviation	Meaning
od	Once daily
bd / bid	Twice daily
td / tid	Three times daily
qds / qid	Four times daily
prn	As needed
ac	Before meals
pc	After meals
stat	Immediately, as a single dose
po	By mouth
top.	Applied to the skin

WHEN IN DOUBT, ASK

Abbreviations are a convenience for professionals, not a puzzle for patients to solve. Every pharmacy label is required to also print the full, plain-language instruction — if the shorthand and the plain text ever seem to disagree, always follow up with the pharmacy.

Frequently Asked Questions

INDICATION — Quick, honest answers to the questions people ask most.

Can I take two different pain relievers at the same time?

Sometimes, and doctors do occasionally recommend alternating two different classes of pain reliever for stronger relief. But this should be confirmed with a pharmacist first, since some combinations are unsafe and it is easy to accidentally double up if either medicine is part of a combination product.

Is it safe to take medicine on an empty stomach?

It depends entirely on the medicine. Some are absorbed better on an empty stomach; others can irritate the stomach lining unless taken with food. The label's directions reflect exactly how that particular medicine was tested.

Why does my doctor want to review all my medicines every year?

A yearly review, sometimes called a medicine reconciliation, checks whether every medicine is still needed, still at the right dose, and still free of new interactions — especially important as new prescriptions are added over time by different doctors.

What should I do if I'm not sure whether a symptom is a side effect?

Note when it started relative to starting the medicine, describe it clearly, and raise it with a doctor or pharmacist rather than stopping the medicine abruptly on your own, unless the symptom is severe — in which case, seek care immediately.

More Questions, Answered

Can children take adult medicine in a smaller amount?

Generally no — as covered in Chapter 11, children's bodies process medicines differently, and paediatric formulations and doses are specifically designed and tested for them, not simply scaled down from an adult version.

Does alcohol really interact with that many medicines?

Yes — alcohol can intensify drowsiness, irritate the stomach alongside certain pain relievers, and interfere with how the liver processes many other medicines. Checking the label's warnings section is the quickest way to know for a specific product.

Is it ever okay to use someone else's leftover prescription?

No. As explained in Chapter 9, a prescription is matched to one person's diagnosis, weight, and health history — using someone else's can mean the wrong treatment entirely, even if symptoms look similar.

How do I know if a supplement is regulated the same way as a medicine?

In most countries it isn't. As explained in Chapter 1, supplements are not required to prove they treat a disease the way medicines must, so it is worth discussing any supplement with a doctor or pharmacist, especially alongside other medicines.

REFERENCE

Glossary of Terms

INDICATION — Plain-language definitions for words used throughout this guide.

Active ingredient — the substance in a medicine responsible for its therapeutic effect.

Anaphylaxis — a severe, whole-body allergic reaction requiring emergency care.

Biologic — a medicine manufactured from living cells rather than pure chemistry.

Contraindication — a specific situation, condition, or combination in which a medicine should not be used.

Dosage — the amount of a medicine, and how often it is taken.

Dose-response curve — the relationship between how much of a medicine is taken and the size of its effect.

Excipient — an inactive ingredient that helps deliver, stabilise, or flavour a medicine.

Generic medicine — a version of a medicine sold after patent protection ends, matching the original in strength and quality.

Interaction — a change in how a medicine behaves caused by another medicine, food, or substance.

Pharmacodynamics — the study of how a medicine affects the body.

Pharmacokinetics — the study of how the body absorbs, moves, breaks down, and removes a medicine.

Prescription — formal authorisation from a licensed provider to use a specific medicine.

Side effect — an unintended effect of a medicine, separate from its main purpose.

Tolerance — a reduced response to a medicine after repeated use, sometimes requiring dose adjustment.

Where to Find Each Topic

A short index for readers looking to revisit a specific idea without paging through the whole guide.

Topic	Where to Look
Allergic reactions & anaphylaxis	Chapter 8
Antibiotic resistance	Chapter 5
Children's dosing	Chapters 3, 11
Dose-response relationship	Chapter 3
Expiry dates	Chapter 9
Generic vs. brand-name	Chapter 3
Interactions with alcohol & food	Chapter 8
Label abbreviations	Chapter 15
Pregnancy & breastfeeding	Chapter 11
Prescription vs. OTC	Chapter 6
Safe disposal	Chapter 9
Storage temperature & light	Chapters 7, 9
Travel & time zones	Chapter 12

CLOSING NOTE

A medicine is never just a chemical — it is trust, translated into a tablet, a syrup, or a drop. Use that trust wisely: read the label, ask the pharmacist, and let a professional guide the way.

THE MEDICINE CABINET · A GENERAL EDUCATIONAL GUIDE · EXPANDED EDITION