

Online Supplement For

"Count Data Models with Social Interactions under Rational Expectations"

Elysée Aristide Houndetoungan*

S.1 Proof of the convergence of the infinite summations

Many infinite summations appear in the paper (e.g., the expected choice, the infinite summations in Proposition 2.2, Assumption 2.4, and several others used throughout the proofs). In this section, I state and prove a general lemma on the convergence of these infinite sums.

Lemma S.1. *Let h be a continuous function on \mathbb{R} and f_γ be a function defined for any $u \in \mathbb{R}$ as $f_\gamma(u) = \sum_{r=0}^{+\infty} r^\gamma h(u - b_r)$, where $\gamma \geq 0$ and $(b_k)_{k \in \mathbb{N}}$ is an increasing positive sequence, such that $\lim_{r \rightarrow \infty} r^{-\rho}(b_{r+1} - b_r) > 0$, where $\rho \geq 0$. The following statements hold.*

- (i) *For any $u \in \mathbb{R}$, if $h(x) = o(|x|^{-\kappa})$ at $-\infty$, where $(1 + \rho)\kappa > 1 + \gamma$, then $f_\gamma(u) < \infty$.*
- (ii) *If $h(x) = o(|x|^{-\kappa})$ at both $-\infty$ and $+\infty$, where $(1 + \rho)\kappa > 1$, then f_0 is bounded on \mathbb{R} .*

Statement (ii) and Assumption 2.3 ensure that B_c defined in Assumption 2.4 is finite. Statement (i) and Assumption 2.3 also imply that the other infinite summations in the paper are finite.

Proof of Lemma S.1

The proof is done in several steps.

Step 0: I show that if $h(x) = o(|x|^{-\kappa})$ at both $-\infty$ and $+\infty$, then $\exists M \geq 1$, such that $|h(u - b_r)| \leq M(|u - b_r| + 1)^{-\kappa}$. Moreover, this is also true for large r even if $h(x) = o(|x|^{-\kappa})$ only at $-\infty$.

The condition $h(x) = o(|x|^{-\kappa})$ at both $-\infty$ and $+\infty$ is also equivalent to $|h(x)| = o((|x| + 1)^{-\kappa})$. Thus, $\exists x_0 \in \mathbb{R}_+ / \forall x < -x_0$ or $x > x_0$, $|h(x)| < (|x| + 1)^{-\kappa}$. As h is continuous, this implies that there exists $M \geq 1$, such that $\forall x \in \mathbb{R}$, $|h(x)| \leq M(|x| + 1)^{-\kappa}$. As a result, $|h(u - b_r)| \leq M(|u - b_r| + 1)^{-\kappa}$.

Step 1: I prove Statement (i).

Let f^* be the real-valued function defined as $f^*(u) = \sum_{r=0}^{\infty} (|u - b_r| + 1)^{-\kappa}$, $\forall u \in \mathbb{R}$.

The condition $\lim_{r \rightarrow \infty} r^{-\rho}(b_{r+1} - b_r) > 0$ implies that there exists $k_0 \in \mathbb{N}$ and $b > 0$, such that $\forall r \geq k_0$, $r^{-\rho}(b_{r+1} - b_r) \geq b$, i.e., $b_{r+1} \geq b \sum_{s=k_0}^r s^\rho + b_{k_0}$. As $\lim_{r \rightarrow \infty} b_r = \infty$, $\forall u \in \mathbb{R}$, it is possible to choose k_0 sufficiently large, such that $b_{k_0} > u$. It follows that $\forall r > k_0$, $|u - b_r| = b_r - u \geq b \sum_{s=k_0}^{r-1} s^\rho + b_{k_0} - u \geq 0$, which implies $(|u - b_r| + 1)^{-\kappa} \leq \left(b \sum_{s=k_0}^{r-1} s^\rho + b_{k_0} - u\right)^{-\kappa}$, and thus $(|u - b_r| + 1)^{-\kappa} \leq O(r^{-(1+\rho)\kappa})$

*Thema, Cy Cergy Paris Université; Email: aristide.houndetoungan@cyu.fr.

since $\sum_{s=k_0}^{r-1} s^\rho = O(r^{1+\rho})$. Therefore, $f^*(u) < \infty$, $\forall u \in \mathbb{R}$. Using the result of the step 0, it follows that $\forall u \in \mathbb{R}$, $\gamma \geq 0$, $r^\gamma h(u - b_r) = O(r^{-(1+\rho)\kappa+\gamma})$. Hence, $f_\gamma(u) < \infty$ if $(1 + \rho)\kappa > 1 + \gamma$.

Step 2: I prove Statement (ii).

As $|h(u - b_r)| \leq M(|u - b_r| + 1)^{-\kappa}$, it is sufficient to prove that f^* is bounded. Moreover, since f^* is a continuous function, this also amounts to proving $\lim_{u \rightarrow -\infty} f^*(u)$ and $\lim_{u \rightarrow +\infty} f^*(u)$ are finite.

For any $u \leq 0$, I have $(|u - b_r| + 1)^{-\kappa} = (b_r - u + 1)^{-\kappa} \leq (b_r + 1)^{-\kappa}$. Thus, $f^*(u) \leq f^*(0)$.

Since f^* is a positive function, this implies that $\lim_{u \rightarrow -\infty} f^*(u)$ is finite.

Let $k_0 \in \mathbb{N}^*$, such that $\forall r, r' \geq k_0$ with $r > r'$, $b_r - b_{r'} \geq b \sum_{s=r'}^{r-1} s^\rho$, for some $b > 0$.

For u positive and sufficiently large, $\exists k^* \in \mathbb{N}$ (with k^* depending on u), where $k^* > k_0$ and $\forall r < k^*$, $u > b_r$, and $\forall r \geq k^*$, $u \leq b_r$. Thus, $f^*(u)$ can be decomposed as

$$\begin{aligned} f^*(u) &= \sum_{r=0}^{k_0-1} (|u - b_r| + 1)^{-\kappa} + \sum_{r=k_0}^{k^*-1} (|u - b_r| + 1)^{-\kappa} + \sum_{r=k^*}^{\infty} (|u - b_r| + 1)^{-\kappa}, \\ f^*(u) &\leq k_0 + \sum_{r=k_0}^{k^*-1} (u - b_r + 1)^{-\kappa} + \sum_{r=k^*}^{\infty} (b_r - u + 1)^{-\kappa}, \\ f^*(u) &\leq k_0 + \sum_{r=k_0}^{k^*-1} (b_{k^*-1} - b_r + 1)^{-\kappa} + \sum_{r=k^*}^{\infty} (b_r - b_{k^*} + 1)^{-\kappa}, \\ f^*(u) &\leq 2 + k_0 + \sum_{r=k_0}^{k^*-2} (b_{k^*-1} - b_r)^{-\kappa} + \sum_{r=k^*+1}^{\infty} (b_r - b_{k^*})^{-\kappa}. \end{aligned}$$

If $k_0 \leq r \leq k^* - 1$, then $b_{k^*-1} - b_r \geq b \sum_{s=r}^{k^*-2} s^\rho$. Thus, $(b_{k^*-1} - b_r)^{-\kappa} \leq \left(b \sum_{s=r}^{k^*-2} s^\rho\right)^{-\kappa}$.

Analogously, if $k^* \leq r$, then $b_r - b_{k^*} \geq b \sum_{s=k^*}^{r-1} s^\rho$. Thus, $(b_r - b_{k^*})^{-\kappa} \leq \left(b \sum_{s=k^*}^{r-1} s^\rho\right)^{-\kappa}$. Therefore,

$$\begin{aligned} f^*(u) &\leq 2 + k_0 + \sum_{r=k_0}^{k^*-2} \left(b \sum_{s=r}^{k^*-2} s^\rho\right)^{-\kappa} + \sum_{r=k^*+1}^{\infty} \left(b \sum_{s=k^*}^{r-1} s^\rho\right)^{-\kappa}, \\ f^*(u) &\leq 2 + k_0 + \sum_{r=k_0}^{k^*-2} \left(b \sum_{s=k_0}^r s^\rho\right)^{-\kappa} + \sum_{r=k^*+1}^{\infty} \left(b \sum_{s=k^*}^{r-1} s^\rho\right)^{-\kappa}, \\ f^*(u) &\leq 2 + k_0 + 2 \sum_{r=k_0}^{\infty} \left(\sum_{s=k_0}^r s^\rho\right)^{-\kappa}. \end{aligned}$$

As $2 + k_0 + 2 \sum_{r=k_0}^{\infty} \left(\sum_{s=k_0}^r s^\rho\right)^{-\kappa}$ does not depend on u and $\left(\sum_{s=k_0}^r s^\rho\right)^{-\kappa} = O(r^{-(1+\rho)\kappa})$, $\lim_{u \rightarrow +\infty} f^*(u)$ is finite if $(1 + \rho)\kappa > 1$. As a result, f_0 is bounded.

S.2 Variance of the NPL estimator

I assume that $\hat{R} \geq \bar{R}^0$. By Proposition 3.2, this implies that $\hat{\theta}(\hat{R}) = \theta^0$ and $\hat{\mathbf{y}}_n^e(\hat{R}) = \mathbf{y}_\chi^{e0}$. I recall that $\theta = \left(\log(\lambda), \Gamma', \log(\tilde{\delta}'), \log(\bar{\delta}), \log(\rho)\right)'$. Let $\phi_{i,r} = \phi(\lambda \mathbf{g}_i \mathbf{y}^e + \mathbf{z}_i' \Gamma - a_r)$, $\Phi_{i,r} = \Phi(\lambda \mathbf{g}_i \mathbf{y}^e + \mathbf{z}_i' \Gamma - a_r)$, $\Delta \phi_{i,r} = \phi_{i,r} - \phi_{i,r-1}$, and $\Delta \Phi_{i,r} = \Phi_{i,r} - \Phi_{i,r-1}$ for any $r \geq 1$, where $\bar{R} = \bar{R}^0$, $\theta = \theta^0$, and $\mathbf{y}^e = \mathbf{y}_\chi^{e0}$. I have

$$\nabla_{\log(\lambda)} \mathcal{L}_n(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) = \lambda \sum_{i=1}^n \sum_{r=0}^{\infty} d_{ir} \frac{\Delta(\phi_{i,r+1} \tilde{z}_{i,r})}{\Delta \Phi_{i,r+1}}, \text{ where } \tilde{z}_{i,r} = \mathbf{g}_i \mathbf{y}^e - r,$$

$$\nabla_{\Gamma} \mathcal{L}_n(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) = \sum_{i=1}^n \sum_{r=0}^{\infty} d_{ir} \frac{\Delta \phi_{i,r+1}}{\Delta \Phi_{i,r+1}} \mathbf{z}_i,$$

$$\nabla_{\log(\tilde{\delta}_k)} \mathcal{L}_{n,i}(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) = -\tilde{\delta}_k \sum_{i=1}^n \sum_{r=k-1}^{\infty} d_{ir} \frac{\phi_{i,r+1}}{\Delta \Phi_{i,r+1}} + \tilde{\delta}_k \sum_{i=1}^n \sum_{r=k}^{\infty} d_{ir} \frac{\phi_{i,r}}{\Delta \Phi_{i,r+1}} \text{ for } 2 \leq k \leq \bar{R},$$

$$\nabla_{\log(\bar{\delta})} \mathcal{L}_{n,i}(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) = -\bar{\delta} \sum_{i=1}^n \sum_{r=\bar{R}}^{\infty} d_{ir} \frac{\dot{a}_{\delta,r+1} \phi_{i,r+1}}{\Delta \Phi_{i,r+1}} + \bar{\delta} \sum_{i=1}^n \sum_{r=\bar{R}+1}^{\infty} d_{ir} \frac{\dot{a}_{\delta,r} \phi_{i,r}}{\Delta \Phi_{i,r+1}},$$

$$\nabla_{\log(\rho)} \mathcal{L}_n(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) = -\rho \sum_{i=1}^n \sum_{r=\bar{R}}^{\infty} d_{ir} \frac{\dot{a}_{\rho,r+1} \phi_{i,r+1}}{\Delta \Phi_{i,r+1}} + \rho \sum_{i=1}^n \sum_{r=\bar{R}+1}^{\infty} d_{ir} \frac{\dot{a}_{\rho,r} \phi_{i,r}}{\Delta \Phi_{i,r+1}},$$

where $\dot{a}_{\delta,r} = \sum_{k=\bar{R}+1}^r (k-1)^\rho$ and $\dot{a}_{\rho,r} = \bar{\delta} \sum_{k=\bar{R}+1}^r (k-1)^\rho \log(k-1)$ for $r \geq \bar{R}+1$.

I define the following notations: $\mathbf{A}_i^{\lambda\lambda} = \lambda^2 \sum_{r=0}^{\infty} \frac{\phi_{i,r+1}^2 \tilde{z}_{i,r}^2 - 2\phi_{i,r} \phi_{i,r+1} \tilde{z}_{i,r} \tilde{z}_{i,r-1} + \phi_{i,r}^2 \tilde{z}_{i,r-1}^2}{\Delta \Phi_{i,r+1}},$

$$\mathbf{A}_i^{\Gamma\Gamma} = \sum_{r=0}^{\infty} \frac{\phi_{i,r+1}^2 - 2\phi_{i,r} \phi_{i,r+1} + \phi_{i,r}^2}{\Delta \Phi_{i,r+1}},$$

$$\mathbf{A}_i^{\delta_k \delta_k} = \tilde{\delta}_k^2 \left(\sum_{r=k-1}^{\infty} \frac{\phi_{i,r+1}^2}{\Delta \Phi_{i,r+1}} - 2 \sum_{r=k}^{\infty} \frac{\phi_{i,r} \phi_{i,r+1}}{\Delta \Phi_{i,r+1}} + \sum_{r=k}^{\infty} \frac{\phi_{i,r}^2}{\Delta \Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\bar{\delta}\bar{\delta}} = \bar{\delta}^2 \left(\sum_{r=\bar{R}}^{\infty} \frac{\dot{a}_{\delta,r+1} \phi_{i,r+1}^2}{\Delta \Phi_{i,r+1}} - 2 \sum_{r=\bar{R}+1}^{\infty} \frac{a_{\delta,r} a_{\delta,r+1} \phi_{i,r} \phi_{i,r+1}}{\Delta \Phi_{i,r+1}} + \sum_{r=\bar{R}+1}^{\infty} \frac{a_{\delta,r}^2 \phi_{i,r}^2}{\Delta \Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\rho\rho} = \rho^2 \left(\sum_{r=\bar{R}}^{\infty} \frac{\dot{a}_{\rho,r+1} \phi_{i,r+1}^2}{\Delta \Phi_{i,r+1}} - 2 \sum_{r=\bar{R}+1}^{\infty} \frac{a_{\rho,r} a_{\rho,r+1} \phi_{i,r} \phi_{i,r+1}}{\Delta \Phi_{i,r+1}} + \sum_{r=\bar{R}+1}^{\infty} \frac{a_{\rho,r}^2 \phi_{i,r}^2}{\Delta \Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\lambda\Gamma} = \lambda \sum_{r=0}^{\infty} \frac{\phi_{i,r+1}^2 \tilde{z}_{i,r} - \phi_{i,r+1} \phi_{i,r} \tilde{z}_{i,r-1} - \phi_{i,r} \phi_{i,r+1} \tilde{z}_{i,r} + \phi_{i,r}^2 \tilde{z}_{i,r-1}}{\Delta \Phi_{i,r+1}},$$

$$\mathbf{A}_i^{\lambda\delta_k} = \lambda \tilde{\delta}_k \left(\sum_{r=k-1}^{\infty} \frac{\phi_{i,r} \phi_{i,r+1} \tilde{z}_{i,r-1} - \phi_{i,r+1}^2 \tilde{z}_{i,r}}{\Delta \Phi_{i,r+1}} + \sum_{r=k}^{\infty} \frac{\phi_{i,r} \phi_{i,r+1} \tilde{z}_{i,r} - \phi_{i,r}^2 \tilde{z}_{i,r-1}}{\Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\lambda\bar{\delta}} = \lambda \bar{\delta} \left(\sum_{r=\bar{R}}^{\infty} \frac{\dot{a}_{\delta,r+1} \phi_{i,r} \phi_{i,r+1} \tilde{z}_{i,r-1} - \dot{a}_{\delta,r+1} \phi_{i,r+1}^2 \tilde{z}_{i,r}}{\Delta \Phi_{i,r+1}} + \sum_{r=\bar{R}+1}^{\infty} \frac{\dot{a}_{\delta,r} \phi_{i,r} \phi_{i,r+1} \tilde{z}_{i,r} - \dot{a}_{\delta,r} \phi_{i,r}^2 \tilde{z}_{i,r-1}}{\Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\lambda\rho} = \lambda \rho \left(\sum_{r=\bar{R}}^{\infty} \frac{\dot{a}_{\rho,r+1} \phi_{i,r} \phi_{i,r+1} \tilde{z}_{i,r-1} - \dot{a}_{\rho,r+1} \phi_{i,r+1}^2 \tilde{z}_{i,r}}{\Delta \Phi_{i,r+1}} + \sum_{r=\bar{R}+1}^{\infty} \frac{\dot{a}_{\rho,r} \phi_{i,r} \phi_{i,r+1} \tilde{z}_{i,r} - \dot{a}_{\rho,r} \phi_{i,r}^2 \tilde{z}_{i,r-1}}{\Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\Gamma\delta_k} = \tilde{\delta}_k \left(\sum_{r=k-1}^{\infty} \frac{\phi_{i,r} \phi_{i,r+1} - \phi_{i,r+1}^2}{\Delta \Phi_{i,r+1}} + \sum_{r=k}^{\infty} \frac{\phi_{i,r} \phi_{i,r+1} - \phi_{i,r}^2}{\Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\Gamma\bar{\delta}} = \bar{\delta} \left(\sum_{r=\bar{R}}^{\infty} \frac{\dot{a}_{\delta,r+1} \phi_{i,r} \phi_{i,r+1} - \dot{a}_{\delta,r+1} \phi_{i,r+1}^2}{\Delta \Phi_{i,r+1}} + \sum_{r=\bar{R}+1}^{\infty} \frac{\dot{a}_{\delta,r} \phi_{i,r} \phi_{i,r+1} - \dot{a}_{\delta,r} \phi_{i,r}^2}{\Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\Gamma\rho} = \rho \left(\sum_{r=\bar{R}}^{\infty} \frac{\dot{a}_{\rho,r+1} \phi_{i,r} \phi_{i,r+1} - \dot{a}_{\rho,r+1} \phi_{i,r+1}^2}{\Delta \Phi_{i,r+1}} + \sum_{r=\bar{R}+1}^{\infty} \frac{\dot{a}_{\rho,r} \phi_{i,r} \phi_{i,r+1} - \dot{a}_{\rho,r} \phi_{i,r}^2}{\Phi_{i,r+1}} \right),$$

$$\mathbf{A}_i^{\delta_k \delta_{k'}} = -\tilde{\delta}_k \mathbf{A}_i^{\Gamma\delta_{k'}} \text{ for } 2 \leq k < k' \leq \bar{R}, \quad \mathbf{A}_i^{\delta_k \bar{\delta}} = -\tilde{\delta}_k \mathbf{A}_i^{\Gamma\bar{\delta}}, \quad \mathbf{A}_i^{\delta_k \rho} = -\tilde{\delta}_k \mathbf{A}_i^{\Gamma\rho},$$

$$\mathbf{A}_i^{\bar{\delta}\rho} = \bar{\delta} \rho \left(\sum_{r=\bar{R}}^{\infty} \frac{\dot{a}_{\delta,r+1} \dot{a}_{\rho,r+1} \phi_{i,r+1}^2}{\Delta \Phi_{i,r+1}} - \sum_{r=\bar{R}+1}^{\infty} \frac{\dot{a}_{\delta,r} \dot{a}_{\rho,r+1} \phi_{i,r} \phi_{i,r+1} + \dot{a}_{\delta,r+1} \dot{a}_{\rho,r} \phi_{i,r} \phi_{i,r+1} - \dot{a}_{\delta,r} \dot{a}_{\rho,r} \phi_{i,r}^2}{\Delta \Phi_{i,r+1}} \right).$$

Let $\Sigma_{n,i} := \mathbb{V}(\nabla_{\boldsymbol{\theta}} \mathcal{L}_{n,i}(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) | \chi_n)$. It follows that

$$\Sigma_{n,i} = - \begin{pmatrix} \mathbf{A}_i^{\lambda\lambda} & \mathbf{A}_i^{\lambda\Gamma} \mathbf{z}'_i & \mathbf{A}_i^{\lambda\delta_2} & \dots & \mathbf{A}_i^{\lambda\rho} \\ \mathbf{A}_i^{\lambda\Gamma} \mathbf{z}_i & \mathbf{A}_i^{\Gamma\Gamma} \mathbf{z}_i \mathbf{z}'_i & \mathbf{A}_i^{\Gamma\delta_2} \mathbf{z}_i & \dots & \mathbf{A}_i^{\Gamma\rho} \mathbf{z}_i \\ \mathbf{A}_i^{\lambda\delta_2} & \mathbf{A}_i^{\Gamma\delta_2} \mathbf{z}'_i & \mathbf{A}_i^{\delta_2\delta_2} & \dots & \mathbf{A}_i^{\delta_2\rho} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{A}_i^{\lambda\rho} & \mathbf{A}_i^{\Gamma\rho} \mathbf{z}'_i & \mathbf{A}_i^{\delta_2\rho} & \dots & \mathbf{A}_i^{\rho\rho} \end{pmatrix}$$

By the law of large numbers (LLN), Σ_0 is the limit of $(1/n) \sum_{i=1}^n \Sigma_{n,i}$ as n grows to infinity.

On the other hand, I have

$$\begin{aligned} \mathbf{H}_{1,n} &:= \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}'} \mathcal{L}_n(\dot{\boldsymbol{\theta}}_n(\hat{R}), \dot{\mathbf{y}}_n^e(\hat{R})) \text{ and } \mathbf{H}_{2,n} := \nabla_{\boldsymbol{\theta}\mathbf{y}^{e'}} \mathcal{L}_n(\dot{\boldsymbol{\theta}}_n(\hat{R}), \dot{\mathbf{y}}_n^e(\hat{R})) \nabla_{\boldsymbol{\theta}'\mathbf{y}^{e'}}(\hat{R}) \text{ for some point } \dot{\boldsymbol{\theta}}_n(\hat{R}) \\ &\text{between } \hat{\boldsymbol{\theta}}_n(\hat{R}) \text{ and } \boldsymbol{\theta}^0, \text{ such that } \dot{\mathbf{y}}_n^e(\hat{R}) = \mathbf{L}(\dot{\boldsymbol{\theta}}_n(\hat{R}), \dot{\mathbf{y}}_n^e(\hat{R})). \text{ As } \hat{\boldsymbol{\theta}}_n(\hat{R}) \text{ converges to } \boldsymbol{\theta}^0, \text{ by the LLN,} \\ \mathbf{H}_{1,n} &\rightarrow \mathbf{H}_{1,0} := \lim_{n \rightarrow \infty} (1/n) \sum_{i=1}^n \mathbb{E}(\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}'} \mathcal{L}_{i,n}(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) | \chi_n) \text{ and} \\ \mathbf{H}_{2,n} &\rightarrow \mathbf{H}_{2,0} := \lim_{n \rightarrow \infty} (1/n) \sum_{i=1}^n \mathbb{E}(\nabla_{\boldsymbol{\theta}\mathbf{y}^{e'}} \mathcal{L}_{i,n}(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) | \chi_n) \nabla_{\boldsymbol{\theta}'\mathbf{y}^{e0}}. \end{aligned}$$

On can verify that $\mathbb{E}(\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}'} \mathcal{L}_{i,n}(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) | \chi_n) = -\Sigma_{n,i}$. Thus, $\mathbf{H}_{1,0} = -\Sigma_0$.

Besides, $\mathbb{E}(\nabla_{\boldsymbol{\theta}\mathbf{y}^{e'}} \mathcal{L}_{i,n}(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0}) | \chi_n) = \lambda(\mathbf{A}_i^{\lambda\Gamma}, \mathbf{A}_i^{\Gamma\Gamma} \mathbf{z}_i', \mathbf{A}_i^{\Gamma\delta_2}, \dots, \mathbf{A}_i^{\Gamma\rho})' \mathbf{g}_i$.

$\nabla_{\boldsymbol{\theta}\mathbf{y}_{\chi}^{e0}}$ can be computed using the implicit definition of \mathbf{y}_{χ}^{e0} ; that is $\mathbf{y}_{\chi}^{e0} = \mathbf{L}(\boldsymbol{\theta}^0, \mathbf{y}_{\chi}^{e0})$. This implies that $\nabla_{\boldsymbol{\theta}\mathbf{y}_{\chi}^{e0}} = \mathbf{S}^{-1}\mathbf{B}$, where $\mathbf{S} = \mathbf{I}_n - \lambda\mathbf{D}\mathbf{G}$, \mathbf{I}_n is the identity matrix of dimension n , $\mathbf{D} = \text{diag}(\sum_{r=1}^{\infty} \phi_{1,r}, \dots, \sum_{r=1}^{\infty} \phi_{n,r})$, and $\mathbf{B} = (\mathbf{B}^1, \mathbf{D}\mathbf{Z}, \mathbf{B}^2)$. The component \mathbf{B}^1 is an n -vector whose i -th element is $\lambda \sum_{r=1}^{\infty} \phi_{i,r} \tilde{z}_{i,r-1}$. The component \mathbf{B}^2 is a matrix of n rows, where the i -th row is $\mathbf{B}_i^2 = \left(-\tilde{\delta}_2 \sum_{r=2}^{\infty} \phi_{i,r}, \dots, -\tilde{\delta}_{\bar{R}} \sum_{r=\bar{R}}^{\infty} \phi_{i,r}, -\tilde{\delta} \sum_{r=\bar{R}+1}^{\infty} \phi_{i,r} \dot{a}_{\delta,r}, -\rho \sum_{r=\bar{R}+1}^{\infty} \phi_{i,r} \dot{a}_{\rho,r}\right)$.

I assume that $\lim_{n \rightarrow \infty} \frac{\lambda}{n} \sum_{i=1}^n (\mathbf{A}_i^{\lambda\Gamma}, \mathbf{A}_i^{\Gamma\Gamma} \mathbf{z}_i', \mathbf{A}_i^{\Gamma\delta_2}, \dots, \mathbf{A}_i^{\Gamma\rho})' \mathbf{g}_i \mathbf{S}^{-1} \mathbf{B}$ exists and is equal to $\boldsymbol{\Omega}_0$. As a result, $\mathbf{H}_{1,0} = -\Sigma_0$, $\mathbf{H}_{2,0} = \boldsymbol{\Omega}_0$, where Σ_0 is the limit of $(1/n) \sum_i \Sigma_{n,i}$ as n grows to infinity.

S.3 Asymptotics in the case of endogenous networks

Let $\hat{\boldsymbol{\mu}}_n = (\hat{\mu}_1, \dots, \hat{\mu}_n)'$ and $\hat{\boldsymbol{\nu}}_n = (\hat{\nu}_1, \dots, \hat{\nu}_n)'$, where $\hat{\mu}_i$ and $\hat{\nu}_i$ are the consistent estimators of μ_i and ν_i , respectively. Let also $\tilde{\mu}_i = \sum_{j=1}^n \mathbf{g}_{ij} \hat{\mu}_j$, $\tilde{\nu}_i = \sum_{j=1}^n \mathbf{g}_{ij} \hat{\nu}_j$, and $\hat{\chi}_n = (\hat{\boldsymbol{\mu}}_n', \hat{\boldsymbol{\nu}}_n')'$. As I have new regressors that are estimated, I define the following notations.

For any \bar{R} , $\boldsymbol{\theta}^*(\bar{R})$ is the vector of new parameters to be estimated. $\boldsymbol{\Theta}^*(\bar{R})$ is the space of $\boldsymbol{\theta}^*(\bar{R})$. The mapping \mathbf{L} is redefined as $\mathbf{L}^*(\boldsymbol{\theta}^*, \mathbf{y}^e) = \sum_{r=0}^{\infty} \Phi(\lambda \mathbf{g}_i \mathbf{y}^e + \mathbf{z}_i' \boldsymbol{\Gamma} - \hat{h}_{\varepsilon,i} - a_r)$, where $\hat{h}_{\varepsilon,i} = \sum_k^T (\theta_{1,k} \hat{\mu}_i^k + \theta_{2,k} \hat{\nu}_i^k + \theta_{3,k} \tilde{\mu}_i^k + \theta_{4,k} \tilde{\nu}_i^k)$ is the consistent approximation of $h_{\varepsilon}(\mu_i, \nu_i, \bar{\mu}_i, \bar{\nu}_i)$.

$\mathcal{L}_{n,i}^*(\boldsymbol{\theta}, \mathbf{y}^e) = \sum_{r=0}^{\infty} d_{ir} \log(\Phi(\lambda \mathbf{g}_i \mathbf{y}^e + \mathbf{z}_i' \boldsymbol{\Gamma} - \hat{h}_{\varepsilon,i} - a_r) - \Phi(\lambda \mathbf{g}_i \mathbf{y}^e + \mathbf{z}_i' \boldsymbol{\Gamma} - \hat{h}_{\varepsilon,i} - a_{r+1}))$. $\mathcal{L}_0^*(\boldsymbol{\theta}, \mathbf{y}^e) \equiv \mathbb{E}(\mathcal{L}_n^*(\boldsymbol{\theta}, \mathbf{y}^e) | \chi_n, \hat{\chi}_n)$, $\tilde{\boldsymbol{\theta}}^*(\mathbf{y}^e, \bar{R}) \equiv \arg \max_{\boldsymbol{\theta} \in \boldsymbol{\Theta}(\bar{R})} \mathcal{L}_0^*(\boldsymbol{\theta}, \mathbf{y}^e)$, $\phi_0^*(\mathbf{y}^e, \bar{R}) \equiv \mathbf{L}^*(\tilde{\boldsymbol{\theta}}^*(\mathbf{y}^e, \bar{R}), \mathbf{y}^e)$, and $\mathcal{A}_0^*(\bar{R}) \equiv \{(\boldsymbol{\theta}^*, \mathbf{y}^e) \in \boldsymbol{\Theta}^*(\bar{R}) \times [0, \bar{y}]^n, \text{ such that } \boldsymbol{\theta}^* = \tilde{\boldsymbol{\theta}}_0^*(\mathbf{y}^e, \bar{R}) \text{ and } \mathbf{y}^e = \phi_0^*(\mathbf{y}^e, \bar{R})\}$. Let also $\boldsymbol{\theta}^{*0}$ be the true value of $\boldsymbol{\theta}^*$ and $\mathbf{y}_{\chi}^{e*0} \in \mathbb{R}^n$, such that $\mathbf{y}_{\chi}^{e*0} = \mathbf{L}^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\chi}^{e*0})$.

S.3.1 Consistency

Under Assumptions A.1–A.2 adapted to the new regressors and $\boldsymbol{\Theta}^*(\bar{R})$, Results A.1–A.2 can be extended to the new pseudo-likelihood. Thus, \mathcal{L}_n^* uniformly converges to \mathcal{L}_0^* . Moreover, \mathcal{L}_0^* has a unique maximizer $(\check{\boldsymbol{\theta}}_n^*(\bar{R}), \check{\mathbf{y}}_n^{e*}(\bar{R}))$, such that $\check{\mathbf{y}}_n^{e*}(\bar{R}) = \mathbf{L}(\check{\boldsymbol{\theta}}_n^*(\bar{R}), \check{\mathbf{y}}_n^{e*}(\bar{R}))$. As for the case of exogenous network, under Assumptions A.1–A.2 and Assumptions A.3–A.5 adapted to the new maximizer

$(\check{\boldsymbol{\theta}}_n^*(\bar{R}), \check{\mathbf{y}}_n^{e*}(\bar{R}))$, the new NPL estimator $\hat{\boldsymbol{\theta}}_n^*(\hat{\bar{R}})$ converges in probability to $\check{\boldsymbol{\theta}}_n^*(\hat{\bar{R}})$. However, Gibbs' inequality cannot be applied because $h_\varepsilon(\mu_i, \nu_i, \bar{\mu}_i, \bar{\nu}_i)$ is replaced by its estimator. However, as the estimator is assumed to be consistent, Gibbs' inequality can be applied as n grows to infinity. It then follows that $\lim_{n \rightarrow \infty} \check{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}) = \boldsymbol{\theta}^{*0}$ if $\hat{\bar{R}} \geq \bar{R}^0$. As a result, the NPL estimator converges to $\boldsymbol{\theta}^{*0}$ if $\bar{R} \geq \bar{R}^0$.

S.3.2 Asymptotic normality

I assume for simplicity that $\hat{\bar{R}} \geq \bar{R}^0$.

By applying the mean value theorem (MVT) to $\nabla_{\boldsymbol{\theta}^*} \mathcal{L}_n^*(\hat{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}), \hat{\mathbf{y}}_n^{e*}(\hat{\bar{R}}))$ between $\hat{\boldsymbol{\theta}}_n^*(\hat{\bar{R}})$ and $\boldsymbol{\theta}^{*0}$, I get

$$\sqrt{n}(\hat{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}) - \boldsymbol{\theta}^{*0}) = -(\mathbf{H}_{1,n}^* + \mathbf{H}_{2,n}^*)^{-1} \sqrt{n} \nabla_{\boldsymbol{\theta}^*} \mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0}) \quad (\text{S.1})$$

for some point $\dot{\boldsymbol{\theta}}_n^*(\hat{\bar{R}})$ between $\hat{\boldsymbol{\theta}}_n^*(\hat{\bar{R}})$ and $\check{\boldsymbol{\theta}}_n^*(\hat{\bar{R}})$, where $\mathbf{H}_{1,n}^* := \nabla_{\boldsymbol{\theta}^* \boldsymbol{\theta}^{*0}} \mathcal{L}_n(\dot{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}), \dot{\mathbf{y}}_n^{e*}(\hat{\bar{R}}))$, $\mathbf{H}_{2,n}^* := \nabla_{\boldsymbol{\theta}^* \mathbf{y}^{e*}} \mathcal{L}_n(\dot{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}), \dot{\mathbf{y}}_n^{e*}(\hat{\bar{R}})) \nabla_{\boldsymbol{\theta}^* \mathbf{y}^{e*}} \mathcal{L}_n(\dot{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}), \dot{\mathbf{y}}_n^{e*}(\hat{\bar{R}})) = \mathbf{L}^*(\dot{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}))$, and $\dot{\mathbf{y}}_n^{e*}(\hat{\bar{R}})$.

I apply the MVT a second time to $\nabla_{\boldsymbol{\theta}^*} \mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0})$ between $\hat{\boldsymbol{\chi}}_n$ and $\hat{\boldsymbol{\chi}}_0 = (\mu_1^0, \dots, \mu_n^0, \nu_1^0, \dots, \nu_n^0)$, where μ_i^0 and ν_i^0 are the true values of μ_i and ν_i , respectively. I have

$$\sqrt{n}(\hat{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}) - \boldsymbol{\theta}^{*0}) = -(\mathbf{H}_{1,n}^* + \mathbf{H}_{2,n}^*)^{-1} (\sqrt{n} \nabla_{\boldsymbol{\theta}^*} \mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0}) + n^{-1/2} \sum_{i=1}^n \mathcal{E}_{i,n} \Delta \hat{\boldsymbol{\chi}}_n), \quad (\text{S.2})$$

where $\nabla_{\boldsymbol{\theta}^*} \mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0})$ is the value of $\nabla_{\boldsymbol{\theta}^*} \mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0})$ when $\hat{\boldsymbol{\mu}}_n$ and $\hat{\boldsymbol{\nu}}_n$ are equal to their true values, $\Delta \hat{\boldsymbol{\chi}}_n = \hat{\boldsymbol{\chi}}_n - \hat{\boldsymbol{\chi}}_0$, and $\mathcal{E}_{i,n}$ is the derivative of $\nabla_{\boldsymbol{\theta}^*} \mathcal{L}_{n,i}^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0})$ with respect to $\hat{\boldsymbol{\chi}}_n$, applied to some point between $\hat{\boldsymbol{\chi}}_n$ and $\hat{\boldsymbol{\chi}}_0$. I set the following regulatory condition.

Assumption S.1. $n^{-1/2} \sum_{i=1}^n \mathcal{E}_{i,n} \Delta \hat{\boldsymbol{\chi}}_n$ is $o_p(1)$.

A similar assumption is also set in the case of the control function approach.² Assumption S.1 requires $\Delta \hat{\boldsymbol{\chi}}_n$ to converge to zero at some rate. As $\|\Delta \hat{\boldsymbol{\chi}}_n\|_\infty = O_p\left((\log(n)/n)^{1/2}\right)$, a sufficient condition for this assumption to hold is that $(1/n) \|\sum_{i=1}^n \mathcal{E}_{i,n}\|_\infty = o_p(\zeta_n^{1/2})$, where $\zeta_n \log(n)$ converges to zero as n grows to infinity. This condition is realistic because $\zeta_n = n^{-\zeta}$ for any $\zeta \in (0, 1)$, if $\mathcal{E}_{i,n}$'s were independent across i . Therefore, if the dependence is not too strong, Assumption S.1 will be verified.

Under Assumption S.1, Equation (S.2) implies that

$$\sqrt{n}(\hat{\boldsymbol{\theta}}_n^*(\hat{\bar{R}}) - \boldsymbol{\theta}^{*0}) = -(\mathbf{H}_{1,n}^* + \mathbf{H}_{2,n}^*)^{-1} \sqrt{n} \nabla_{\boldsymbol{\theta}^*} \mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0}) + o_p(1).$$

The CLT can be applied to $\sqrt{n} \nabla_{\boldsymbol{\theta}^*} \mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0})$ since $\mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\boldsymbol{\chi}}^{e*0})$ is a sum of n independent

²See the Lipschitz condition set in Assumption 8 of Johnsson, I. and H. R. Moon (2021): "Estimation of peer effects in endogenous social networks: control function approach," *Review of Economics and Statistics*, 103, 328–345.

bounded variables conditionally on χ_n . It then follows that

$$\sqrt{n}(\hat{\boldsymbol{\theta}}_n^*(\hat{R}) - \boldsymbol{\theta}^{*0}) \xrightarrow{d} \mathcal{N}(0, (\mathbf{H}_{1,0}^* + \mathbf{H}_{2,0}^*)^{-1} \boldsymbol{\Sigma}_0^* (\mathbf{H}_{1,0}^{*'} + \mathbf{H}_{2,0}^{*'})^{-1}), \quad (\text{S.3})$$

where $\mathbf{H}_{1,0}^*$ and $\mathbf{H}_{2,0}^*$ are the limits of $\mathbf{H}_{1,n}^*$ and $\mathbf{H}_{2,n}^*$ as n grows to infinity, and $\boldsymbol{\Sigma}_0^*$ is a consistent estimator of the variance of $\sqrt{n} \nabla_{\boldsymbol{\theta}} \mathcal{L}_n^*(\boldsymbol{\theta}^{*0}, \mathbf{y}_{\chi}^{e*0})$.

S.4 Marginal effects

The parameters of the counting variable model cannot be interpreted directly. Policymakers are interested in the marginal effect of the explanatory variables on the expected outcome.

Let us recall that $\boldsymbol{\theta} = (\log(\lambda), \boldsymbol{\Gamma}', \log(\tilde{\boldsymbol{\delta}}'), \log(\bar{\delta}), \log(\rho))'$, where $\tilde{\boldsymbol{\delta}} = (\tilde{\delta}_2, \dots, \tilde{\delta}_{\bar{R}})$, $\tilde{\delta}_r = \delta_r - \lambda$, $a_0 = -\infty$, $a_1 = 0$, $a_r = \sum_{k=2}^r \delta_k$ for any $r \geq 2$, and $\delta_r = (r-1)\bar{\delta} + \lambda \forall r > \bar{R}$.

Let $\tilde{\mathbf{z}}'_i = (\mathbf{g}_i \mathbf{y}^e, \mathbf{z}'_i)$ and $\boldsymbol{\Lambda} = (\lambda, \boldsymbol{\Gamma}')'$. For any $k = 1, \dots, \dim(\boldsymbol{\Lambda})$, let λ_k and \tilde{z}_{ik} be the k -th component in $\boldsymbol{\Lambda}$ and $\tilde{\mathbf{z}}_i$, respectively. The marginal effect of the explanatory variable \tilde{z}_{ik} on y_i^e is given by

$$\delta_{ik}(\boldsymbol{\theta}) = \frac{\partial y_i^e}{\partial \tilde{z}_{ik}} = \lambda_k \sum_{r=1}^{\infty} \phi(\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r). \quad (\text{S.4})$$

The standard error of $\delta_{ik}(\boldsymbol{\theta})$ can be computed using the Delta method.

By the mean value theorem, I have

$$\frac{1}{n} \sum_{i=1}^n \delta_{ik}(\hat{\boldsymbol{\theta}}_n(\hat{R})) \stackrel{a}{\sim} \mathcal{N}\left(\delta_{ik}(\boldsymbol{\theta}_0), \mathbf{Q}_0^* \mathbb{V}(\hat{\boldsymbol{\theta}}_n(\hat{R}) | \chi_n) \mathbf{Q}_0^{*'}\right),$$

where $\mathbf{Q}_0^* = (1/n) \sum_{i=1}^n \nabla_{\boldsymbol{\theta}} \delta_{ik}(\boldsymbol{\theta}_0)$,

$$\nabla_{\log(\lambda)} \delta_{ik}(\boldsymbol{\theta}) = \mathbf{1}(k=1) \lambda \sum_{r=1}^{\infty} \phi(\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r) - \lambda \lambda_k \sum_{r=1}^{\infty} (\mathbf{g}_i \mathbf{y}^e - r) (\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r) \phi(\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r),$$

$\nabla_{\boldsymbol{\Gamma}'} \delta_{ik}(\boldsymbol{\theta}) = \mathbf{e}_k \sum_{r=1}^{\infty} \phi(\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r) - \lambda_k \mathbf{z}'_i \sum_{r=1}^{\infty} (\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r) \phi(\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r)$, where \mathbf{e}_k is a $\dim(\boldsymbol{\Gamma})$ -dimensional row vector with zero everywhere except the $(k-1)$ -th term which equals one if $k \geq 2$,

$$\nabla_{\log(\delta_l)} \delta_{ik}(\boldsymbol{\theta}) = \tilde{\delta}_l \lambda_k \sum_{r=l}^{\infty} (\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r) \phi(\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r) \quad \text{for } 2 \leq l < \bar{R},$$

$$\nabla_{\log(\bar{\delta})} \delta_{ik}(\boldsymbol{\theta}) = \bar{\delta} \lambda_k \sum_{r=\bar{R}+1}^{\infty} \dot{a}_{\delta,r} (\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r) \phi(\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r),$$

$$\nabla_{\log(\rho)} \delta_{ik}(\boldsymbol{\theta}) = \rho \lambda_k \sum_{r=l}^{\infty} \dot{a}_{\rho,r} (\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r) \phi(\tilde{\mathbf{z}}'_i \boldsymbol{\Lambda} - a_r).$$

S.5 Empirical results controlling for network endogeneity

Table S.2: Empirical results controlling for network endogeneity (NE). Unobserved attributes are treated as random effects and estimated using a Bayesian approach

Parameters	Count data model			Tobit model		
	Coef.	Marginal effects		Coef.	Marginal effects	
λ	0.046	0.084	(0.020)	0.307	0.249	(0.019)
Own effects						
Age	-0.044	-0.080	(0.008)	-0.045	-0.037	(0.005)
Male	-0.163	-0.298	(0.016)	-0.265	-0.215	(0.011)
Hispanic	-0.006	-0.012	(0.025)	0.116	0.094	(0.017)
Race						
Black	0.208	0.381	(0.031)	0.444	0.360	(0.02)
Asian	0.209	0.383	(0.034)	0.660	0.535	(0.023)
Other	0.027	0.050	(0.028)	0.137	0.111	(0.018)
Years at school	0.030	0.054	(0.007)	0.089	0.072	(0.005)
With both par.	0.073	0.134	(0.019)	0.146	0.118	(0.012)
Mother educ.						
<High	-0.052	-0.094	(0.023)	-0.033	-0.027	(0.015)
>High	0.202	0.369	(0.019)	0.382	0.310	(0.013)
Missing	0.028	0.052	(0.031)	0.211	0.172	(0.021)
Mother job						
Professional	0.134	0.245	(0.024)	0.235	0.191	(0.016)
Other	0.035	0.064	(0.020)	0.053	0.043	(0.013)
Missing	-0.041	-0.074	(0.028)	-0.069	-0.056	(0.019)
μ^1	0.480	0.879	(0.110)	0.917	0.744	(0.073)
μ^2	-1.065	-1.949	(0.428)	-2.122	-1.721	(0.286)
μ^3	-0.705	-1.290	(1.608)	1.050	0.851	(1.080)
μ^4	1.121	2.052	(2.088)	0.615	0.499	(1.404)
ν^1	0.286	0.524	(0.083)	0.338	0.274	(0.056)
ν^2	0.668	1.224	(0.348)	1.810	1.468	(0.232)
ν^3	-0.551	-1.009	(1.504)	0.606	0.492	(1.009)
ν^4	0.575	1.053	(1.736)	0.684	0.555	(1.170)
Contextual effects						
Age	-0.021	-0.039	(0.004)	-0.075	-0.061	(0.003)
Male	-0.050	-0.092	(0.030)	-0.044	-0.035	(0.019)
Hispanic	-0.067	-0.123	(0.042)	-0.077	-0.062	(0.027)
Race						
Black	0.059	0.109	(0.039)	0.016	0.013	(0.026)
Asian	-0.011	-0.020	(0.052)	-0.238	-0.193	(0.035)
Other	-0.101	-0.185	(0.052)	-0.254	-0.206	(0.035)
Years at school	0.016	0.030	(0.011)	-0.007	-0.006	(0.007)
With both par.	0.134	0.246	(0.036)	0.164	0.133	(0.024)
Mother educ.						
<High	-0.116	-0.213	(0.043)	-0.160	-0.130	(0.028)
>High	0.168	0.307	(0.038)	0.197	0.160	(0.025)
Missing	-0.056	-0.102	(0.060)	-0.132	-0.107	(0.040)
Mother job						
Professional	0.165	0.302	(0.048)	0.221	0.180	(0.031)
Other	0.029	0.053	(0.038)	0.019	0.016	(0.025)
Missing	-0.014	-0.026	(0.054)	0.025	0.020	(0.036)
μ^1	0.022	0.041	(0.203)	-0.200	-0.162	(0.135)
μ^2	-0.638	-1.169	(0.894)	-0.889	-0.721	(0.597)
μ^3	-1.894	-3.468	(4.116)	-2.241	-1.818	(2.752)
μ^4	0.967	1.771	(5.839)	0.871	0.706	(3.906)
ν^1	0.597	1.092	(0.171)	0.906	0.735	(0.114)
ν^2	0.705	1.290	(1.099)	0.881	0.715	(0.733)
ν^3	-0.739	-1.354	(4.220)	-1.183	-0.960	(2.821)
ν^4	-0.270	-0.494	(4.465)	-1.191	-0.966	(2.986)
σ				2.420		

For the count data model, $\hat{R} = 12$. The estimates of $\delta_2, \dots, \delta_{\hat{R}}$ are 1.555, 0.523, 0.452, 0.385, 0.320, 0.264, 0.218, 0.174, 0.130, 0.100, 0.086. The estimate of $\bar{\delta}$ is $1.2e^{-5}$.

Table S.3: Empirical results controlling for NE. Unobserved attributes are treated as fixed effects

Parameters	Count data model			Tobit model		
	Coef.	Marginal effects		Coef.	Marginal effects	
λ	0.046	0.084	(0.024)	0.304	0.246	(0.025)
Own effects						
Age	-0.044	-0.081	(0.008)	-0.050	-0.041	(0.005)
Male	-0.160	-0.293	(0.017)	-0.259	-0.210	(0.011)
Hispanic	-0.013	-0.023	(0.025)	0.108	0.088	(0.017)
Race						
Black	0.224	0.411	(0.032)	0.524	0.425	(0.021)
Asian	0.206	0.378	(0.034)	0.641	0.520	(0.023)
Other	0.027	0.050	(0.028)	0.127	0.103	(0.018)
Years at school	0.034	0.062	(0.007)	0.101	0.082	(0.005)
With both par.	0.074	0.137	(0.019)	0.150	0.122	(0.012)
Mother educ.						
<High	-0.055	-0.102	(0.023)	-0.040	-0.033	(0.015)
>High	0.206	0.378	(0.020)	0.390	0.317	(0.013)
Missing						
Mother job	0.027	0.049	(0.031)	0.205	0.166	(0.021)
Professional	0.135	0.249	(0.024)	0.242	0.196	(0.016)
Other	0.037	0.069	(0.020)	0.060	0.049	(0.013)
Missing	-0.039	-0.072	(0.028)	-0.066	-0.053	(0.019)
μ^1	0.126	0.232	(0.137)	0.300	0.244	(0.154)
μ^2	-0.082	-0.151	(0.751)	-0.339	-0.275	(1.440)
μ^3	-0.589	-1.081	(1.851)	-2.132	-1.729	(7.031)
μ^4	-0.452	-0.829	(2.074)	-1.424	-1.156	(18.735)
μ^5	-0.006	-0.010	(0.857)	1.021	0.828	(27.358)
μ^6				0.418	0.339	(20.495)
μ^7				-0.449	-0.364	(6.145)
ν^1	0.093	0.170	(0.014)	0.167	0.135	(0.010)
ν^2	0.061	0.113	(0.055)	0.233	0.189	(0.058)
ν^3	0.048	0.088	(0.074)	-0.356	-0.289	(0.119)
ν^4	-0.109	-0.200	(0.188)	-0.063	-0.051	(0.377)
ν^5	-0.144	-0.265	(0.163)	1.914	1.553	(0.312)
ν^6				0.070	0.056	(0.782)
ν^7				-1.499	-1.216	(0.535)
Contextual effects						
Age	-0.011	-0.021	(0.010)	-0.061	-0.049	(0.007)
Male	-0.044	-0.081	(0.030)	-0.030	-0.024	(0.020)
Hispanic	-0.048	-0.087	(0.041)	-0.059	-0.048	(0.027)
Race						
Black	0.123	0.226	(0.042)	0.074	0.060	(0.029)
Asian	-0.017	-0.032	(0.053)	-0.243	-0.197	(0.036)
Other	-0.105	-0.193	(0.052)	-0.259	-0.210	(0.035)
Years at school	0.003	0.005	(0.012)	-0.021	-0.017	(0.008)
With both par.	0.134	0.246	(0.037)	0.173	0.141	(0.024)
Mother educ.						
<High	-0.126	-0.231	(0.043)	-0.179	-0.145	(0.029)
>High	0.181	0.333	(0.040)	0.216	0.175	(0.026)
Missing	-0.065	-0.119	(0.060)	-0.143	-0.116	(0.041)
Mother job						
Professional	0.176	0.322	(0.048)	0.241	0.195	(0.032)
Other	0.036	0.066	(0.038)	0.034	0.028	(0.025)
Missing	-0.011	-0.019	(0.055)	0.036	0.029	(0.036)
μ^1	0.164	0.301	(0.241)	0.633	0.514	(0.213)
μ^2	0.138	0.253	(2.465)	2.213	1.795	(3.868)
μ^3	-0.019	-0.035	(11.064)	2.720	2.207	(36.684)
μ^4	-0.187	-0.344	(22.434)	-0.265	-0.215	(184.271)
μ^5	0.017	0.031	(16.807)	-0.539	-0.438	(498.076)
μ^6				0.787	0.639	(684.420)
μ^7				-0.370	-0.301	(374.754)
ν^1	0.032	0.060	(0.018)	0.113	0.092	(0.0160)
ν^2	0.021	0.038	(0.077)	-0.025	-0.021	(0.088)
ν^3	-0.034	-0.063	(0.109)	-0.651	-0.528	(0.168)
ν^4	0.075	0.138	(0.301)	0.609	0.494	(0.635)
ν^5	0.110	0.202	(0.280)	1.872	1.518	(0.476)
ν^6				-0.555	-0.450	(1.439)
ν^7				-1.341	-1.088	(0.962)
σ				2.424		

For the count data model, $\hat{R} = 12$. The estimates of $\delta_2, \dots, \delta_{\hat{R}}$ are 1.551, 0.521, 0.450, 0.384, 0.319, 0.263, 0.218, 0.174, 0.130, 0.100, 0.086. The estimate of $\bar{\delta}$ is $1.2e^{-5}$.