



Contents lists available at ScienceDirect

Clinical Microbiology and Infection

journal homepage: www.clinicalmicrobiologyandinfection.com

Original article

Norwegian patients and retail chicken meat share cephalosporin-resistant *Escherichia coli* and IncK/*bla*_{CMY-2} resistance plasmidsE.S. Berg^{1,*}, A.L. Wester¹, J. Ahrenfeldt², S.S. Mo³, J.S. Slettemeås³, M. Steinbakk¹, Ø. Samuelsen^{4,5}, N. Grude⁶, G.S. Simonsen^{4,7}, I.H. Løhr⁸, S.B. Jørgensen⁹, S. Tofteland¹⁰, O. Lund², U.R. Dahle¹, M. Sunde^{1,3}¹ Domain of Infection Control and Environmental Health, Norwegian Institute of Public Health, Oslo, Norway² Centre for Biological Sequence Analysis, Department of Systems Biology, Technical University of Denmark, Kongens Lyngby, Denmark³ Department of Diagnostic Services, Norwegian Veterinary Institute, Oslo, Norway⁴ Norwegian National Advisory Unit on Detection of Antimicrobial Resistance, Department of Microbiology and Infection Control, University Hospital of North Norway, Tromsø, Norway⁵ Research Group for Microbial Pharmacology and Population Biology, Department of Pharmacy, University of Tromsø – The Arctic University of Norway, Tromsø, Norway⁶ Department of Clinical Microbiology, Vestfold Hospital Trust, Tønsberg, Norway⁷ Research Group for Host-Microbe Interactions, Faculty of Health Sciences, University of Tromsø – The Arctic University of Norway, Tromsø, Norway⁸ Department of Medical Microbiology, Stavanger University Hospital, Stavanger, Norway⁹ Department of Clinical Microbiology and Infection Control, Akershus University Hospital, Lørenskog, Norway¹⁰ Department of Clinical Microbiology, Sørlandet Hospital, Kristiansand, Norway

ARTICLE INFO

Article history:

Received 11 August 2016

Received in revised form

11 November 2016

Accepted 28 December 2016

Available online xxx

Editor: G. Lina

Keywords:

Chicken meat

Plasmid-mediated AmpC transmission

Urinary tract infections

Zoonotic transfer of cephalosporin-resistant

Escherichia coli

ABSTRACT

Objectives: In 2012 and 2014 the Norwegian monitoring programme for antimicrobial resistance in the veterinary and food production sectors (NORM-VET) showed that 124 of a total of 406 samples (31%) of Norwegian retail chicken meat were contaminated with extended-spectrum cephalosporin-resistant *Escherichia coli*. The aim of this study was to compare selected cephalosporin-resistant *E. coli* from humans and poultry to determine their genetic relatedness based on whole genome sequencing (WGS). **Methods:** *Escherichia coli* representing three prevalent cephalosporin-resistant multi-locus sequence types (STs) isolated from poultry ($n = 17$) were selected from the NORM-VET strain collections. All strains carried an IncK plasmid with a *bla*_{CMY-2} gene. Clinical *E. coli* isolates ($n = 284$) with AmpC-mediated resistance were collected at Norwegian microbiology laboratories from 2010 to 2014. PCR screening showed that 29 of the clinical isolates harboured both IncK and *bla*_{CMY-2}. All IncK/*bla*_{CMY-2}-positive isolates were analysed with WGS-based bioinformatics tools.

Results: Analysis of single nucleotide polymorphisms (SNP) in 2.5 Mbp of shared genome sequences showed close relationship, with fewer than 15 SNP differences between five clinical isolates from urinary tract infections (UTIs) and the ST38 isolates from poultry. Furthermore, all of the 29 clinical isolates harboured IncK/*bla*_{CMY-2} plasmid variants highly similar to the IncK/*bla*_{CMY-2} plasmid present in the poultry isolates.

Conclusions: Our results provide support for the hypothesis that clonal transfer of cephalosporin-resistant *E. coli* from chicken meat to humans may occur, and may cause difficult-to-treat infections. Furthermore, these *E. coli* can be a source of AmpC-resistance plasmids for opportunistic pathogens in the human microbiota. **E.S. Berg, CMI 2017;■:1**

© 2017 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

* Corresponding author. E.S. Berg, Domain of Infection Control and Environmental Health, Norwegian Institute of Public Health, PO Box 4404 Nydalen, 0403 Oslo, Norway.
E-mail address: einarsverre.berg@fhi.no (E.S. Berg).

Introduction

Use of antimicrobials in human and veterinary medicine and in the livestock industry drives the emergence, selection and spread of bacterial resistance [1]. Food contaminated with antibiotic-resistant bacteria can be a source of resistance for human pathogens either by direct colonization of zoonotic bacteria, or by transfer of mobile genetic elements, e.g. plasmids, to other gut-colonizing bacteria [2].

The World Health Organization defines extended-spectrum cephalosporins as critically important antimicrobials for human medicine [3]. Many studies have shown that β -lactamase-producing *Escherichia coli* occurs in chicken meat [3,4]. Widespread use of cephalosporins in the broiler production pyramid and the international trade of breeding animals and hatching eggs enhance this into a global problem [5]. However, to what extent consumption of chicken meat contaminated by resistant bacteria affects human health has not been established [2,6].

With the exception of coccidiostats (not classified as an antibiotic in Europe), Norwegian broiler production has a very low usage of antimicrobials [7–10]. In spite of this, cephalosporin-resistant *E. coli* were detected in approximately one-third of the broilers and retail chicken fillets produced in Norway in the period from 2011 to 2014 [7–10]. A recent study showed that cephalosporin-resistant *E. coli* originating from Norwegian chicken fillets had low genetic diversity, mainly limited to a few multi-locus sequence types (STs), and typically carried the plasmid-mediated AmpC (pAmpC) *bla*_{CMY-2} gene on an IncK plasmid [11]. The poultry-associated *E. coli* had many of the same virulence factors as human extraintestinal pathogenic *E. coli* (ExPEC), which could facilitate human colonization and extraintestinal infections [11].

To investigate if poultry-associated strains existed among human clinical *E. coli* isolates with pAmpC production, we screened isolates from microbiology laboratory and national strain collections. Isolates that were PCR positive for both IncK and *bla*_{CMY-2} were selected for whole genome sequencing (WGS). The aim of the study was to compare human- and poultry-associated cephalosporin-resistant *E. coli* by use of single nucleotide polymorphism (SNP) analyses to determine the genetic relatedness of selected AmpC-producing strains from the two reservoirs with the highest possible resolution.

Materials and methods

Bacterial isolates

The scheme for selection and analysis of the isolates in the present study is illustrated in Fig. 1.

Escherichia coli isolates from broilers. The poultry-associated *E. coli* strains were a subset of isolates originally collected from retail chicken meat (fillets) in 2012 ($n = 4$) and 2014 ($n = 10$) and from chicken faecal samples in 2011 ($n = 3$) as a part of the Norwegian monitoring programme for antimicrobial resistance in the veterinary and food production sectors (NORM/NORM-VET) [7–11]. The monitoring program reported the following prevalence of pAmpC-positive *E. coli* isolates from broilers: 43% (108/252) from intestinal flora of healthy broilers in 2011, and from chicken file samples 32% (66/205) and 29% (58/201) in 2012 and 2014, respectively. Briefly, the 17 isolates selected for this study belonged to phylogroup D and carried similar conjugative IncK plasmids encoding *bla*_{CMY-2}. The strains were typed to ST38 ($n = 10$), ST1158 ($n = 5$) and ST115 ($n = 2$). ST38 and ST1158 were the two most common STs among the




Fig. 1. Pipeline for selection of isolates for whole genome-based comparison of *Escherichia coli* from poultry and human infections. Abbreviations: AMR, antimicrobial resistance; MDR, multidrug resistance; ST, multi-locus sequence type.

cephalosporin-resistant *E. coli* isolated from Norwegian chicken meat in 2012 and 2014 [11]. Furthermore, molecular characterization of the chicken isolates identified virulence factor genes (e.g. *iroN*, *cma* and *iss*) similar to those of ExPEC [11].

Human clinical *E. coli* isolates. The human clinical *E. coli* consisted of two collections of AmpC-producing ExPEC isolates. The first collection (A) included 158 isolates with a *bla*_{CMY-2}-positive genotype tested as described by Brolund *et al.* [12]. These isolates were from Norwegian clinical microbiology laboratories sampled during 2010–2012 and originally submitted for verification of the presence of pAmpC at the Norwegian National Advisory Unit on Detection of Antimicrobial Resistance. Submission criteria were resistance to cefotaxime (MIC >2 mg/L) and ceftazidime (MIC >4 mg/L) in combination with multi-drug resistance (MDR) (defined as resistance to at least two of the following groups of antibiotics: aminoglycosides, fluoroquinolones, trimethoprim-sulfamethoxazole or nitrofurantoin). The second collection (B) included 126 *E. coli* isolates displaying an AmpC phenotype identified at nine Norwegian clinical microbiology laboratories during 2013–2014. The phenotypic AmpC profile was determined according to EUCAST [13]. None of the isolates displayed classical serine extended-spectrum β -lactamase production. The original sample materials for the 284 human clinical *E. coli* isolates (of which 84% were from UTIs) are presented in Fig. 1.

Ethical considerations

This study was approved by the Norwegian Regional Committees for Medical and Health Research Ethics (REC) (ref. 2014/419/REK sør-øst).

Molecular characterization

PCR screening. All human clinical isolates were screened for the presence of IncK plasmids and for the *bla*_{CMY-2} gene. Twenty-nine *E. coli* isolates harboured both targets. The real-time PCR targeting the IncK replicon was based on previously published IncK primers [14]. The *bla*_{CMY-2} target was amplified in triplex-PCR format similar to a previously published real-time pAmpC PCR [12]. Further technical details for the PCR assays are presented in the Supplementary material (Appendix S1).

Whole genome sequencing (WGS). Human clinical (*n* = 29) and poultry (*n* = 17) isolates PCR-positive for *bla*_{CMY-2} and IncK were subjected to WGS. Paired-end genomic libraries with insert size 500 bp were made from approximately 100 ng purified DNA and sequenced on an Illumina HiSeq 2500 system (BGI Tech Solutions Co., Ltd., Hong Kong). Sequencing was performed twice for two poultry isolates (2011-01-2112 and 2012-01-1292) and one clinical isolate (E4-14) to provide WGS reproducibility controls.

Bioinformatic analysis of WGS data

Genome assembly and in silico genotyping. The raw sequence data were initially trimmed and cleaned for adaptors (BGI Tech Solutions). The paired-end reads were further cleaned for errors and *de novo* assembled into contigs (draft genomes) by the SPAdes-3.6.1 pipeline using default settings [15]. The contigs were submitted to the Centre for Genomic Epidemiology (CGE), Technical University of Denmark (DTU), for bioinformatics analysis (<https://cge.cbs.dtu.dk/services/>).

Identification of SNPs and phylogenetic analysis. The paired-end read sets, error-corrected by SPAdes, were uploaded to CSI Phylogeny 1.1 [16] on the DTU server by use of the default minimum settings (<https://cge.cbs.dtu.dk/services/>). The assembled scaffold of one of the poultry isolates (2012-01-1292) was used as the

reference sequence by the mapping of the read sets. A selection of completed *E. coli* reference genomes downloaded from National Center for Biotechnology Information (NCBI) (<http://www.ncbi.nlm.nih.gov/genome/167>) was included in the analysis to limit the SNP calling to the phylogenetic informative genomic sequences assumed to be representative for the *E. coli* core genome [17]. The rationale was that only the evolutionarily conserved sequences would remain as shared core genome sequences after the sequence alignment of highly different *E. coli* strains [18]. A final maximum-likelihood tree was constructed using FigTREE (<http://tree.bio.ed.ac.uk/software/figtree>).

IncK plasmid assembly and plasmid SNP analysis. Draft sequences of the IncK plasmid were made by mapping of the trimmed read sets against several IncK plasmids available from GenBank using BioNUMERICS, version 7.5 (Applied Maths, Sint-Martens-Latem, Belgium). The best match was obtained using the poultry-associated strain 53C unnamed 3 IncK plasmid with *bla*_{CMY-2} assembled by de Been *et al.* as reference (Accession no. NZ_JXMX01000007.1, (https://www.ncbi.nlm.nih.gov/nuccore/NZ_JXMX01000007.1?report=GenBank) [19]. Finally, paired-end read sets from each isolate were uploaded to CSI phylogeny for SNP-based plasmid similarity analysis using the 53C IncK plasmid as reference.

BEAST simulation. To estimate a timeline for the diversification of the most closely related isolates of poultry and human origin, Bayesian Evolutionary Analysis Sampling Trees, BEAST version 1.7 software was used with input from CSI Phylogeny analysis of the 15 most closely related isolates [20]. A basic assumption for the time estimation was that any SNPs located in recombination sites of the *E. coli* genomes had been removed in the CSI Phylogeny analysis, i.e. by pruning the SNPs. Among several models of evolution, the chosen model with the best fit assumed an uncorrelated relaxed molecular clock, expansion growth in population size and a lognormal distribution of the mutation rate. All the BEAST Monte Carlo Markov Chain simulations were run for 150 million steps, and samples were saved every 10 000 steps. A single maximum clade credibility tree was produced using the TREEANNOTATOR tool in the BEAST pipeline, where 10% of the Monte Carlo Markov Chain steps were removed as burn-in. A final tree was constructed using FigTREE.

Results

In total, 29 of the 284 clinical *E. coli* isolates were PCR positive for both *bla*_{CMY-2} and IncK. In collection A, consisting of 158 genotypic *bla*_{CMY-2}-positive MDR clinical isolates, 16 carried IncK plasmids (14 isolates from UTIs, one isolate from bloodstream infection, and one isolate from an unknown body site). In collection B with 126 phenotypic AmpC-positive clinical isolates, 53 isolates were *bla*_{CMY-2}-positive and 13 of these were also IncK positive (11 isolates from UTIs, one isolate from a perianal abscess and one from an unknown body site) (see Supplementary material, Table S1, Appendix S1).

WGS-based *in silico* genotyping of the 29 IncK/*bla*_{CMY-2}-positive isolates revealed a diversity of other plasmid replicons (e.g. IncFIB, IncFII and IncI1), acquired antibiotic resistance genes (e.g. *bla*_{TEM-1B}, *sul1/2* and *strA/B*), STs (e.g. ST38, ST69 and ST131), serotypes (e.g. O7:H18, O17/O77:H18 and O25:H4) and virulence factors (e.g. *iroN*, *cma*, *iss*, *eilA* and *iha*) (see Supplementary material, Appendix S1, Table S2 and Fig. S2). A subset of the human isolates (*n* = 7), all from UTIs, showed genotypic features similar to the most prevalent poultry-associated *E. coli* strains (ST38, phylogroup D, serotype O7:H18, carriage of IncK, IncFIB-, IncFII- and p0111 plasmids, and *iroN*, *cma*, *iss* and *eilA* virulence factor genes) [11] (see Supplementary material, Appendix S1, Table S2 and Fig. S2).

The CSI Phylogeny pipeline identified 56 610 valid SNP positions in the shared core genome of 2.51 Mbp of the 29 clinical isolates, the 17 poultry strains and the 23 NCBI reference strains. The phylogenetic analysis revealed that the majority of the 29 clinical isolates were genetically diverse and not closely related to the poultry isolates (Fig. 2). However, from the subset of the seven ST38 UTI isolates, five clustered together with the ten ST38 poultry

isolates (Fig. 3). The number of SNP differences between these UTI and poultry isolates was only 1–13 (Fig. 2 and see [Supplementary material, Appendix S1 and Fig. S2](#)). The remaining ST38 clinical isolates, E2-21 and E4-13 had 34 and 529 SNP differences, respectively, compared with the ST38 reference poultry isolate (Fig. 2 and see [Supplementary material, Appendix S1 and Fig. S2](#)). In contrast, pairwise comparison of any of the ST38 poultry isolates to the non-




Fig. 2. Genome single nucleotide polymorphism (SNP) differences in human clinical *Escherichia coli* isolates compared with one AmpC-producing poultry *E. coli* (ST38) strain collected from Norwegian retail chicken fillets. Multiple sequence alignment resulted in 56,610 valid SNPs in the core genome of 2.51 Mbp positions shared by all strains. The data labels show the span in SNP differences within the group of isolates.




Fig. 3. Dendrogram after whole genome comparison with single nucleotide polymorphism (SNP) calling and phylogenetic analysis of *Escherichia coli* isolates from human clinical infections, poultry samples and a selection of GenBank *E. coli* reference genomes retrieved from GenBank. The colours of the boxes reflect the origin of the isolates as indicated. The multilocus sequence types (STs) of the included poultry isolates and of isolates of clinical relevance are indicated on the figure. STs of all investigated isolates can be found in supplementary material. The length of the scale bar represents the average number of substitutions per site at the variable sites.

ST38 human clinical isolates as well as to the NCBI reference *E. coli* genomes, resulted in 4000–10 600 SNP differences. The poultry strains clustered in three genetically diverse groups according to their STs and showed few SNP differences within each group (ST38 group up to eight SNP differences, ST1158 group up to five SNP differences, and ST115 group one SNP difference) (Figs. 2 and 3). For the three isolates that were sequenced twice as WGS reproducibility controls, no SNP differences were observed between each pair (see Supplementary material, Appendix S1 and Fig. S2).

Mapping of the paired-end read sets to the 53C unnamed 3 IncK plasmid sequence re-constructed by de Been *et al.* [19] showed that the 29 clinical isolates and the poultry strains had IncK plasmid variants highly similar to the 53C reference plasmid (85.9 kb) albeit with a size range of 80–85 kb. The closely related human clinical and chicken isolates as well as the rest of the clinical and poultry isolates shared an IncK plasmid backbone with a 79-kb core sequence. The CSI phylogeny analysis clustered the IncK variants from most of the isolates into one large group whereas the five ST1158 poultry isolates formed a separate group (see Supplementary material, Appendix S1 and Fig. S3).

The genome sequences of the closely related clinical ST38 UTI isolates ($n = 5$) and the poultry strains ($n = 10$) were subjected to a new CSI Phylogeny analysis to produce data input to the BEAST simulation. The SNP alignments were used as input to the BEAST analyses that estimated the point in time for the diversification of the ancestor of the 15 closely related isolates to year 2005 with a mutation rate of 0.0098 mutations/genome/year (Fig. 4). The final tree illustrates that the five clinical UTI *E. coli* isolates belonged to three different clusters (Fig. 4). Apparently, the genetic diversity within this group of UTI isolates was within the same order of magnitude as the genetic diversity between the UTI and poultry isolates.

Discussion

The aim of this study was to use WGS-based SNP analysis to achieve the best discriminatory power by the determination of the genetic relatedness of a selected group of AmpC-producing human clinical ExPEC isolates compared with cephalosporin-resistant *E. coli* found in retail chicken fillets. To focus the investigation on isolates that might have received their AmpC-resistance from poultry-associated *E. coli*, the presence of IncK and *bla*_{CMY-2} was used as a selection criterion for the human clinical isolates [11]. The PCR screening showed that 29 of the 284 AmpC-producing human clinical isolates were IncK/*bla*_{CMY-2}-positive. This indicates that the majority ($n = 182$) of our *bla*_{CMY-2}-positive clinical ExPECs have acquired the pAmpC from other sources than the cephalosporin-resistant *E. coli* found in chicken fillets and were therefore unrelated to the chicken meat *E. coli*. Sequence analysis after WGS confirmed that all of the 29 clinical isolates selected in the screening carried a plasmid backbone with high similarity to the IncK/*bla*_{CMY-2} plasmid in the poultry-associated *E. coli*.

The WGS-based *in silico* low-resolution *E. coli* genotyping of the 29 clinical strains showed that most of these ExPEC isolates were distantly related to the poultry *E. coli*. However, seven of them, all from UTIs, belonged to ST38. The dominating cephalosporin-resistant *E. coli* isolated from the Norwegian retail chicken fillets also belonged to this ST [11]. Several European studies have shown that ST38 is associated with human UTIs, and in addition, may occur as a contaminant of chicken meat [21–25]. The phylogenetic SNP-based analysis provided higher resolution than multilocus sequence typing and showed that only five of the seven clinical ST38 ExPEC isolates were nearly identical to the dominating ST38 *E. coli* isolates collected from poultry ($n = 10$). The high-resolution comparisons of informative SNPs within the multiple sequence




Fig. 4. Bayesian Evolutionary Analysis Sampling Trees (BEAST) analysis of the closely related human and chicken *Escherichia coli* isolates. Yellow boxes; poultry isolates, pink boxes; urinary tract isolates.

alignment-defined common 'core' genome (2.5 Mbp) of the reference strains, the poultry and human isolates revealed very few SNP differences (< 15) between these 15 ST38 isolates. Hereby, the SNP-based phylogenetic result strongly suggests a link between these isolates. Extrapolation and comparison of our findings to similar studies using *E. coli* core genomes derived from proteome analyses indicate that our ten poultry and five UTI isolates were as closely related as would be expected for clones in a foodborne *E. coli* outbreak [6,19]. With the assumption that the isolates were clonal, the BEAST analyses determined the time when these 15 human and poultry *E. coli* isolates diverged from a common ancestor to year 2005. The BEAST output showed that the five clinical *E. coli* isolates were positioned among the poultry isolates in three different clusters. Furthermore, the diversity within these human ExPEC isolates was in the same low order of magnitude as the diversity between the human and poultry *E. coli* isolates.

A limitation of our study is the lack of epidemiological data of patient exposure to chicken meat to strengthen the hypothesis of resistance transmission from poultry to human. Alternatively, data of exposure to other potential sources of antimicrobial resistance could weaken the hypothesis. In addition, the BEAST analysis that was based on only 15 isolates collected over a short time period could not show the evolution of the strains in the two compartments over time. Another limitation of our study is the narrow focus on comparison of SNPs in *E. coli* from the two reservoirs without a view to identify the genes harbouring the SNPs to see a potential biological significance or to study the evolution of ST38 ExPEC. Consumption of chicken meat contaminated with cephalosporin-resistant *E. coli* can be a driver for increasing occurrence of pAmpC-positive pathogens causing human infections. Previous findings have demonstrated the same plasmid-mediated resistance genes (e.g. *bla*_{CTX-M}, *bla*_{SHV}, *bla*_{TEM} and *bla*_{CMY-2}), the same mobile genetic element, and highly similar genotypes of *E. coli* in humans and retail chicken meat. These studies have indicated a common link between these reservoirs [21,25–30]. The WGS-based genome comparisons enabled us to investigate if transmission of resistance may occur by whole bacterium clonal transfer between poultry and humans. Furthermore, the high-resolution sequence comparison also enabled us to investigate if the resistance transmissions may have occurred by horizontal dissemination of plasmids between bacteria. The results of this study provide support for the hypothesis that clonal transmission of AmpC-producing *E. coli* between poultry and humans may occur. Our results based on the SNP comparisons and the BEAST analysis showed that certain ST38 strains from the two reservoirs are very closely related and that they may have a common ancestor. Since the strains are highly similar, we cannot firmly decide in which direction the transmission between the two reservoirs occurred. However, contaminated poultry meat is a recognized source for infections with other zoonotic bacteria such as *Campylobacter*. This strengthens the hypothesis that a recent clonal transfer of resistant *E. coli* might have occurred from a poultry reservoir to human hosts. With the low number of SNP differences (<6 SNPs per Mbp core genome) and close genetic relatedness between the ST38 strains from human and poultry, our results differ from the data from the Netherlands [19]. De Been *et al.* identified 1263 SNP differences per Mbp core genome between their most closely related human and poultry isolates collected in 2006–2011 [19]. The dissimilar results of the two studies may be explained by different sampling strategies and selection of isolates.

We found similar IncK plasmid variants with *bla*_{CMY-2} both in the clonally related human clinical isolates (*n* = 5) and the poultry strains (*n* = 10), and in the rest of the clinical isolates (*n* = 24) and the poultry strains (*n* = 7). This suggests that horizontal transfer of resistance plasmids seems to occur with higher frequency than

clonal transmission of resistant bacteria. Our results are in line with the results from other studies, and supports that plasmids are significant vectors for antimicrobial resistance dissemination [19,31]. Building an IncK/*bla*_{CMY-2} plasmid scaffold by positioning the WGS read sets against the 53C unnamed 3 IncK plasmid (86.0 kb), resulted in a common plasmid backbone sequence of 79 kb for all of our 29 ExPEC isolates. This plasmid backbone might be a part of many of the variants of *bla*_{CMY-2}-containing IncK plasmids detected in the *E. coli* population of broilers in parts of Europe during the last few years [11,19,21–25,32]. Notably, in addition to *bla*_{CMY-2}, the poultry-associated IncK plasmid backbone has two plasmid addiction systems that ensure stable maintenance of the plasmid within each of the daughter cells after cell division [11,33]. Plasmid addiction systems increase the potential for persistence regardless of any antimicrobial selection pressure. In addition, they may increase the potential for resistance plasmid transmission to opportunistic pathogens or to other bacteria in the human and poultry gut microbiota or in environmental reservoirs [33].

Even if only a small fraction of pAmpC-producing ExPEC in human infections is attributable to contaminated food, this is a significant public health concern [6]. Our approach based on targeted selection of resistant isolates for WGS-based high-resolution comparisons supports that transmission of antimicrobial resistance may occur both by clonal transfer of *E. coli* from poultry to humans and more frequently by the transfer of plasmids to pathogenic *E. coli* adapted to humans.

In conclusion, 20% of the 29 selected IncK/*bla*_{CMY-2}-positive ExPEC isolates showed very close relatedness to cephalosporin-resistant *E. coli* found in retail chicken fillets. The remaining 80% of the clinical isolates exhibited a pAmpC vector highly similar to the IncK plasmid carried by the cephalosporin-resistant *E. coli* associated with poultry. This study supports the hypothesis that transfer of antimicrobial resistance may occur between poultry and humans both by clonal transmission and, more frequently, by plasmid dissemination.

Acknowledgements

We wish to acknowledge Åse Berg and Elizabeth Fleur Peacocke for help with revising the manuscript and the staff at the Department of Food-borne Diseases, and Department of Bacteriology, Division of Disease Control, NIPH as well as at the Norwegian National Advisory Unit on Detection of Antimicrobial Resistance Department of Microbiology and Infection Control, UNN for technical assistance. The views expressed in this publication are those of the authors, and not necessarily those of NIPH.

Transparency declaration

The authors have no conflict of interests.

Funding

This study was funded by the Norwegian Institute of Public Health and the Research Council of Norway grant no 233632, which had no role in the study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all data in the study and had final responsibility for the decision to submit for publication.

Authors contribution

ESB, ALW, MSt, ØS, URD and MSu designed and conceived the study. ESB, ALW, SSM, JSS, ØS, NG, GSS, IHL, SBJ, ST and MSu performed initial strain selection and analysis. ESB, JA, SSM, JSS, MS,

OL, and MSu did the data analysis. ESB, JA, and OL performed the bioinformatic analysis. ESB, ALW, JA, SSM, JSS, MSt, ØS, NG, GSS, IHL, SBJ, ST, OL, URD and MSu wrote the manuscript.

Appendix A. Supplementary data

Additional Supporting Information may be found in the online version of this article at <http://dx.doi.org/10.1016/j.cmi.2016.12.035>.

References

- [1] Holmes AH, Moore LS, Sundsfjord A, Steinbakk M, Regmi S, Karkey A, et al. Understanding the mechanisms and drivers of antimicrobial resistance. *Lancet* 2016;387:176–87.
- [2] Lazarus B, Paterson DL, Mollinger JL, Rogers BA. Do human extraintestinal *Escherichia coli* infections resistant to expanded-spectrum cephalosporins originate from food-producing animals? A systematic review. *Clin Infect Dis* 2015;60:439–52.
- [3] WHO. Critically Important Antimicrobials for Human Medicine, 3rd Revision. Available at: http://apps.who.int/iris/bitstream/10665/77376/1/9789241504485_eng.pdf; 2011 [Accessed April 2016].
- [4] EFSA. European Food Safety Authority. Scientific Opinion on the public health risks of bacterial strains producing extended-spectrum β -lactamases and/or AmpC β -lactamases in food and food-producing animals. *EFSA Journal* 2011;9(2322). Available at: http://www.efsa.europa.eu/sites/default/files/scientific_output/files/main_documents/2322.pdf [Accessed April 2016].
- [5] Liebana E, Carattoli A, Coque TM, Hasman H, Magiorakos AP, Mevius D, et al. Public health risks of enterobacterial isolates producing extended-spectrum beta-lactamases or AmpC beta-lactamases in food and food-producing animals: an EU perspective of epidemiology, analytical methods, risk factors, and control options. *Clin Infect Dis* 2013;56:1030–7.
- [6] Manges AR. *Escherichia coli* and urinary tract infections: the role of poultry-meat. *Clin Microbiol Infect* 2016;22:122–9.
- [7] NORM/NORM-VET. 2011. Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Norway. Tromsø / Oslo. 2012. ISSN:1502–2307 (print) / 1890-9965 (electronic).
- [8] NORM/NORM-VET. 2012. Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Norway. Tromsø / Oslo. 2013. ISSN:1502–2307 (print) / 1890-9965 (electronic).
- [9] NORM/NORM-VET. 2014. Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Norway. Tromsø / Oslo. 2015. ISSN:1502–2307 (print) / 1890-9965 (electronic).
- [10] Mo SS, Norstrom M, Slettemeas JS, Lovland A, Urdahl AM, Sunde M. Emergence of AmpC-producing *Escherichia coli* in the broiler production chain in a country with a low antimicrobial usage profile. *Vet Microbiol* 2014;171:315–20.
- [11] Mo SS, Slettemeas JS, Berg ES, Norstrom M, Sunde M. Plasmid and host strain characteristics of *Escherichia coli* resistant to extended-spectrum cephalosporins in the Norwegian broiler production. *PLoS One* 2016;11:e0154019.
- [12] Brolund A, Wisell KT, Edquist PJ, Elfstrom L, Walder M, Giske CG. Development of a real-time SYBRGreen PCR assay for rapid detection of acquired AmpC in *Enterobacteriaceae*. *J Microbiol Methods* 2010;82:229–33.
- [13] EUCAST. EUCAST guidelines for detection of resistance mechanisms and specific resistances of clinical and/or epidemiological importance. EUCAST 2011. Available at: http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Resistance_mechanisms/EUCAST_detection_of_resistance_mechanisms_v1.0_20131211.pdf [Accessed April 2016].
- [14] Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005;63:219–28.
- [15] Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012;19:455–77.
- [16] Kaas RS, Leekitcharoenphon P, Aarestrup FM, Lund O. Solving the problem of comparing whole bacterial genomes across different sequencing platforms. *PLoS One* 2014;9:e104984.
- [17] Leimbach A, Hacker J, Dobrindt U. *E. coli* as an all-rounder: the thin line between commensalism and pathogenicity. *Curr Top Microbiol Immunol* 2013;358:3–32.
- [18] Kaas RS, Friis C, Ussery DW, Aarestrup FM. Estimating variation within the genes and inferring the phylogeny of 186 sequenced diverse *Escherichia coli* genomes. *BMC Genomics* 2012;13:577.
- [19] de Been M, Lanza VF, de Toro M, Scharringa J, Dohmen W, Du Y, et al. Dissemination of cephalosporin resistance genes between *Escherichia coli* strains from farm animals and humans by specific plasmid lineages. *PLoS Genet* 2014;10:e1004776.
- [20] Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Mol Biol Evol* 2012;29:1969–73.
- [21] Voets GM, Fluit AC, Scharringa J, Schapendonk C, van den Munckhof T, Levenstein-van Hall MA, et al. Identical plasmid AmpC beta-lactamase genes and plasmid types in *E. coli* isolates from patients and poultry meat in the Netherlands. *Int J Food Microbiol* 2013;167:359–62.
- [22] Dierikx C, van der Goot J, Fabri T, van Essen-Zandbergen A, Smith H, Mevius D. Extended-spectrum-beta-lactamase- and AmpC-beta-lactamase-producing *Escherichia coli* in Dutch broilers and broiler farmers. *J Antimicrob Chemother* 2013;68:60–7.
- [23] Agerso Y, Jensen JD, Hasman H, Pedersen K. Spread of extended spectrum cephalosporinase-producing *Escherichia coli* clones and plasmids from parent animals to broilers and to broiler meat in a production without use of cephalosporins. *Foodborne Pathog Dis* 2014;11:740–6.
- [24] Seiffert SN, Tinguely R, Lupo A, Neuwirth C, Perreten V, Endimiani A. High prevalence of extended-spectrum-cephalosporin-resistant enterobacteriaceae in poultry meat in Switzerland: emergence of CMY-2- and VEB-6-possessing *Proteus mirabilis*. *Antimicrob Agents Chemother* 2013;57:6406–8.
- [25] Day MJ, Rodriguez I, van Essen-Zandbergen A, Dierikx C, Kadlec K, Schink AK, et al. Diversity of STs, plasmids and ESB genes among *Escherichia coli* from humans, animals and food in Germany, the Netherlands and the UK. *J Antimicrob Chemother* 2016;71:1178–82.
- [26] Jakobsen L, Kurasic A, Skjot-Rasmussen L, Ejrnaes K, Porsbo LJ, Pedersen K, et al. *Escherichia coli* isolates from broiler chicken meat, broiler chickens, pork, and pigs share phylogroups and antimicrobial resistance with community-dwelling humans and patients with urinary tract infection. *Foodborne Pathog Dis* 2010;7:537–47.
- [27] Kluytmans JA, Overdevest IT, Willemsen I, Kluytmans-van den Bergh MF, van der Zwaluw K, Heck M, et al. Extended-spectrum beta-lactamase-producing *Escherichia coli* from retail chicken meat and humans: comparison of strains, plasmids, resistance genes, and virulence factors. *Clin Infect Dis* 2013;56:478–87.
- [28] Levenstein-van Hall MA, Dierikx CM, Cohen Stuart J, Voets GM, van den Munckhof MP, van Essen-Zandbergen A, et al. Dutch patients, retail chicken meat and poultry share the same ESB genes, plasmids and strains. *Clin Microbiol Infect* 2011;17:873–80.
- [29] Overdevest I, Willemsen I, Rijnsburger M, Eustace A, Xu L, Hawkey P, et al. Extended-spectrum beta-lactamase genes of *Escherichia coli* in chicken meat and humans, The Netherlands. *Emerg Infect Dis* 2011;17:1216–22.
- [30] Vincent C, Boerlin P, Daignault D, Dozois CM, Dutil L, Galanakis C, et al. Food reservoir for *Escherichia coli* causing urinary tract infections. *Emerg Infect Dis* 2010;16:88–95.
- [31] Borjesson S, Ny S, Egervarn M, Bergstrom J, Rosengren A, Englund S, et al. Limited dissemination of extended-spectrum beta-lactamase- and plasmid-encoded AmpC-producing *Escherichia coli* from food and farm animals, Sweden. *Emerg Infect Dis* 2016;22:634–40.
- [32] Egervarn M, Borjesson S, Byfors S, Finn M, Kaipe C, Englund S, et al. *Escherichia coli* with extended-spectrum beta-lactamases or transferable AmpC beta-lactamases and *Salmonella* on meat imported into Sweden. *Int J Food Microbiol* 2014;171:8–14.
- [33] Unterholzner SJ, Poppenberger B, Rozhon W. Toxin–antitoxin systems: biology, identification, and application. *Mob Genet Elements* 2013;3:e26219.