

Linear Transformations and Their Matrices

Linear Algebra - 24DS

AH Sheikh QUEST -

Spring 2026

Example 11: $\mathbb{R}^4 \rightarrow \mathbb{R}^2$ — Linear functional

Problem: $T(x_1, x_2, x_3, x_4) = (x_1 - x_3, 2x_2 + x_4)$. Find matrix.

Example 11: $\mathbb{R}^4 \rightarrow \mathbb{R}^2$ — Linear functional

Problem: $T(x_1, x_2, x_3, x_4) = (x_1 - x_3, 2x_2 + x_4)$. Find matrix.

$$T(1, 0, 0, 0) = (1, 0), \quad T(0, 1, 0, 0) = (0, 2), \quad T(0, 0, 1, 0) = (-1, 0), \quad T(0, 0, 0, 1) = (0, 1)$$

$$A = \begin{pmatrix} 1 & 0 & -1 & 0 \\ 0 & 2 & 0 & 1 \end{pmatrix} \quad (2 \times 4)$$

Example 11: $\mathbb{R}^4 \rightarrow \mathbb{R}^2$ — Linear functional

Problem: $T(x_1, x_2, x_3, x_4) = (x_1 - x_3, 2x_2 + x_4)$. Find matrix.

$$T(1, 0, 0, 0) = (1, 0), \quad T(0, 1, 0, 0) = (0, 2), \quad T(0, 0, 1, 0) = (-1, 0), \quad T(0, 0, 0, 1) = (0, 1)$$

$$A = \begin{pmatrix} 1 & 0 & -1 & 0 \\ 0 & 2 & 0 & 1 \end{pmatrix} \quad (2 \times 4)$$

Apply to $(2, 3, 4, 5)$:

$$T = \begin{pmatrix} 2 - 4 \\ 2 \cdot 3 + 5 \end{pmatrix} = \begin{pmatrix} -2 \\ 11 \end{pmatrix}$$

Example 12: $\mathbb{R}^4 \rightarrow \mathbb{R}^4$ — Projection onto 2D subspace

Problem: Project onto span of $\mathbf{e}_1, \mathbf{e}_2$ (zero out last two coordinates).

Example 12: $\mathbb{R}^4 \rightarrow \mathbb{R}^4$ — Projection onto 2D subspace

Problem: Project onto span of $\mathbf{e}_1, \mathbf{e}_2$ (zero out last two coordinates).

$$T(x, y, z, w) = (x, y, 0, 0)$$

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Example 12: $\mathbb{R}^4 \rightarrow \mathbb{R}^4$ — Projection onto 2D subspace

Problem: Project onto span of $\mathbf{e}_1, \mathbf{e}_2$ (zero out last two coordinates).

$$T(x, y, z, w) = (x, y, 0, 0)$$

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$T(1, 2, 3, 4) = (1, 2, 0, 0)$$

Example 12: $\mathbb{R}^4 \rightarrow \mathbb{R}^4$ — Projection onto 2D subspace

Problem: Project onto span of $\mathbf{e}_1, \mathbf{e}_2$ (zero out last two coordinates).

$$T(x, y, z, w) = (x, y, 0, 0)$$

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$T(1, 2, 3, 4) = (1, 2, 0, 0)$$

Note: This is an **idempotent** transformation: $A^2 = A$.

Example 13: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ — Orthogonal projection onto plane

Problem: Project onto plane $x + y + z = 0$. Find matrix (harder).

Example 13: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ — Orthogonal projection onto plane

Problem: Project onto plane $x + y + z = 0$. Find matrix (harder).
We need $T(\mathbf{v}) = \mathbf{v} - \text{proj}_{\mathbf{n}}\mathbf{v}$ where $\mathbf{n} = (1, 1, 1)$.

$$\text{proj}_{\mathbf{n}}\mathbf{v} = \frac{\mathbf{v} \cdot \mathbf{n}}{\|\mathbf{n}\|^2} \mathbf{n} = \frac{x + y + z}{3} (1, 1, 1)$$

Example 13: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ — Orthogonal projection onto plane

Problem: Project onto plane $x + y + z = 0$. Find matrix (harder).
We need $T(\mathbf{v}) = \mathbf{v} - \text{proj}_{\mathbf{n}}\mathbf{v}$ where $\mathbf{n} = (1, 1, 1)$.

$$\text{proj}_{\mathbf{n}}\mathbf{v} = \frac{\mathbf{v} \cdot \mathbf{n}}{\|\mathbf{n}\|^2} \mathbf{n} = \frac{x + y + z}{3} (1, 1, 1)$$

$$T(x, y, z) = \left(x - \frac{S}{3}, y - \frac{S}{3}, z - \frac{S}{3} \right), \quad S = x + y + z$$

Example 13: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ — Orthogonal projection onto plane

Problem: Project onto plane $x + y + z = 0$. Find matrix (harder).
We need $T(\mathbf{v}) = \mathbf{v} - \text{proj}_{\mathbf{n}}\mathbf{v}$ where $\mathbf{n} = (1, 1, 1)$.

$$\text{proj}_{\mathbf{n}}\mathbf{v} = \frac{\mathbf{v} \cdot \mathbf{n}}{\|\mathbf{n}\|^2} \mathbf{n} = \frac{x + y + z}{3} (1, 1, 1)$$

$$T(x, y, z) = \left(x - \frac{S}{3}, y - \frac{S}{3}, z - \frac{S}{3} \right), \quad S = x + y + z$$

Matrix:

$$A = \frac{1}{3} \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$$

Example 13: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ — Orthogonal projection onto plane

Problem: Project onto plane $x + y + z = 0$. Find matrix (harder).
We need $T(\mathbf{v}) = \mathbf{v} - \text{proj}_{\mathbf{n}}\mathbf{v}$ where $\mathbf{n} = (1, 1, 1)$.

$$\text{proj}_{\mathbf{n}}\mathbf{v} = \frac{\mathbf{v} \cdot \mathbf{n}}{\|\mathbf{n}\|^2} \mathbf{n} = \frac{x + y + z}{3} (1, 1, 1)$$

$$T(x, y, z) = \left(x - \frac{S}{3}, y - \frac{S}{3}, z - \frac{S}{3} \right), \quad S = x + y + z$$

Matrix:

$$A = \frac{1}{3} \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$$

Apply to $(1, 0, 0)$: $\frac{1}{3}(2, -1, -1)$ which lies in $x + y + z = 0$

Example 14: $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ — Composition

Problem: Rotate by 90° then reflect across $y = x$. Find combined matrix.

Example 14: $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ — Composition

Problem: Rotate by 90° then reflect across $y = x$. Find combined matrix.

$$A_{\text{rot}} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, A_{\text{refl}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Example 14: $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ — Composition

Problem: Rotate by 90° then reflect across $y = x$. Find combined matrix.

$$A_{\text{rot}} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, A_{\text{refl}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$\text{Composition } T(\mathbf{x}) = A_{\text{refl}}(A_{\text{rot}}\mathbf{x}) = (A_{\text{refl}}A_{\text{rot}})\mathbf{x}$$

$$A_{\text{total}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Example 14: $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ — Composition

Problem: Rotate by 90° then reflect across $y = x$. Find combined matrix.

$$A_{\text{rot}} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, A_{\text{refl}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$\text{Composition } T(\mathbf{x}) = A_{\text{refl}}(A_{\text{rot}}\mathbf{x}) = (A_{\text{refl}}A_{\text{rot}})\mathbf{x}$$

$$A_{\text{total}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

That's a reflection across x-axis! Check on $(1, 0)$: rotate $\rightarrow (0, 1)$,
reflect $\rightarrow (1, 0)$; direct: $(1, 0) \rightarrow (1, 0)$

Example 15: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ — Shear in 3D

Problem: Shear along x direction proportional to z :
 $T(x, y, z) = (x + 2z, y, z)$. Matrix.

Example 15: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ — Shear in 3D

Problem: Shear along x direction proportional to z :
 $T(x, y, z) = (x + 2z, y, z)$. Matrix.

$$T(1, 0, 0) = (1, 0, 0), \quad T(0, 1, 0) = (0, 1, 0), \quad T(0, 0, 1) = (2, 0, 1)$$

$$A = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Example 15: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ — Shear in 3D

Problem: Shear along x direction proportional to z :
 $T(x, y, z) = (x + 2z, y, z)$. Matrix.

$$T(1, 0, 0) = (1, 0, 0), \quad T(0, 1, 0) = (0, 1, 0), \quad T(0, 0, 1) = (2, 0, 1)$$

$$A = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Apply to $(1, 2, 3)$:

$$T = (1 + 6, 2, 3) = (7, 2, 3)$$

Kernel and Image

Kernel (Nullspace)

$\ker(T) = \{\mathbf{v} \mid T(\mathbf{v}) = \mathbf{0}\}$ — vectors mapped to zero.

Kernel and Image

Kernel (Nullspace)

$\ker(T) = \{\mathbf{v} \mid T(\mathbf{v}) = \mathbf{0}\}$ — vectors mapped to zero.

Image (Range)

$\text{Im}(T) = \{T(\mathbf{v}) \mid \mathbf{v} \in \mathbb{R}^n\}$ — all possible outputs.

Kernel and Image

Kernel (Nullspace)

$\ker(T) = \{\mathbf{v} \mid T(\mathbf{v}) = \mathbf{0}\}$ — vectors mapped to zero.

Image (Range)

$\text{Im}(T) = \{T(\mathbf{v}) \mid \mathbf{v} \in \mathbb{R}^n\}$ — all possible outputs.

Dimension Theorem:

$$\dim(\ker(T)) + \dim(\text{Im}(T)) = n$$

Kernel and Image

Kernel (Nullspace)

$\ker(T) = \{\mathbf{v} \mid T(\mathbf{v}) = \mathbf{0}\}$ — vectors mapped to zero.

Image (Range)

$\text{Im}(T) = \{T(\mathbf{v}) \mid \mathbf{v} \in \mathbb{R}^n\}$ — all possible outputs.

Dimension Theorem:

$$\dim(\ker(T)) + \dim(\text{Im}(T)) = n$$

Example: $T(x, y) = (x, 0)$ in $\mathbb{R}^2$.

$$\ker(T) = \{(0, y)\} \Rightarrow \dim = 1, \quad \text{Im}(T) = \{(x, 0)\} \Rightarrow \dim = 1, \quad 1 + 1 = 2$$

Example 16: Finding Kernel

Problem: $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3$ to $\mathbb{R}^3$. Find kernel.

Example 16: Finding Kernel

Problem: $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3$ to $\mathbb{R}^3$. Find kernel.

Set $T(x, y, z) = 0$:

$$\begin{cases} x + y = 0 \\ y + z = 0 \\ x - z = 0 \end{cases}$$

Example 16: Finding Kernel

Problem: $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3$ to $\mathbb{R}^3$. Find kernel.

Set $T(x, y, z) = 0$:

$$\begin{cases} x + y = 0 \\ y + z = 0 \\ x - z = 0 \end{cases}$$

From first: $y = -x$. Second: $z = -y = x$. Third: $x - z = x - x = 0$ (auto). So x free: $(x, -x, x) = x(1, -1, 1)$.

Example 16: Finding Kernel

Problem: $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3$ to $\mathbb{R}^3$. Find kernel.

Set $T(x, y, z) = 0$:

$$\begin{cases} x + y = 0 \\ y + z = 0 \\ x - z = 0 \end{cases}$$

From first: $y = -x$. Second: $z = -y = x$. Third: $x - z = x - x = 0$ (auto). So x free: $(x, -x, x) = x(1, -1, 1)$.

$$\ker(T) = \text{span}\{(1, -1, 1)\}, \quad \dim = 1$$

Check: $T(1, -1, 1) = (0, 0, 0)$

Example 17: Finding Image

Same $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3 \rightarrow \mathbb{R}^3$.

Example 17: Finding Image

Same $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3 \rightarrow \mathbb{R}^3$.
Image = span of columns of matrix:

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{pmatrix}$$

Example 17: Finding Image

Same $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3 \rightarrow \mathbb{R}^3$.
Image = span of columns of matrix:

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{pmatrix}$$

Columns: $C_1 = (1, 0, 1)$, $C_2 = (1, 1, 0)$, $C_3 = (0, 1, -1)$.

Example 17: Finding Image

Same $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3 \rightarrow \mathbb{R}^3$.

Image = span of columns of matrix:

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{pmatrix}$$

Columns: $C_1 = (1, 0, 1)$, $C_2 = (1, 1, 0)$, $C_3 = (0, 1, -1)$. Check independence: $C_3 = C_2 - C_1$? $(0, 1, -1) = (1, 1, 0) - (1, 0, 1) = (0, 1, -1)$
So $\dim(\text{Im}) = 2$, basis $\{(1, 0, 1), (1, 1, 0)\}$.

Example 17: Finding Image

Same $T(x, y, z) = (x + y, y + z, x - z)$ from $\mathbb{R}^3 \rightarrow \mathbb{R}^3$.

Image = span of columns of matrix:

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{pmatrix}$$

Columns: $C_1 = (1, 0, 1)$, $C_2 = (1, 1, 0)$, $C_3 = (0, 1, -1)$. Check independence: $C_3 = C_2 - C_1$? $(0, 1, -1) = (1, 1, 0) - (1, 0, 1) = (0, 1, -1)$
So $\dim(\text{Im}) = 2$, basis $\{(1, 0, 1), (1, 1, 0)\}$. Check dimension theorem:
 $\dim \ker = 1$, $\dim \text{Im} = 2$, $1 + 2 = 3$

Example 18: $\mathbb{R}^4 \rightarrow \mathbb{R}^4$ — Rotation in 4D?

Problem: Rotation in $\mathbb{R}^4$ can be block diagonal:

$$A = \begin{pmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & \cos \phi & -\sin \phi \\ 0 & 0 & \sin \phi & \cos \phi \end{pmatrix}$$

Example 18: $\mathbb{R}^4 \rightarrow \mathbb{R}^4$ — Rotation in 4D?

Problem: Rotation in $\mathbb{R}^4$ can be block diagonal:

$$A = \begin{pmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & \cos \phi & -\sin \phi \\ 0 & 0 & \sin \phi & \cos \phi \end{pmatrix}$$

$$\text{For } \theta = 90^\circ, \phi = 0: A = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Example 18: $\mathbb{R}^4 \rightarrow \mathbb{R}^4$ — Rotation in 4D?

Problem: Rotation in $\mathbb{R}^4$ can be block diagonal:

$$A = \begin{pmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & \cos \phi & -\sin \phi \\ 0 & 0 & \sin \phi & \cos \phi \end{pmatrix}$$

For $\theta = 90^\circ$, $\phi = 0$: $A = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ Apply to $(1, 0, 2, 3)$:

$$T = (0, 1, 2, 3)$$

First two coordinates rotate, last two unchanged.

Example 19: Differentiation as Linear Transformation

Problem: $T : P_3 \rightarrow P_2$ (polynomials degree 3 to 2) defined by $T(p) = p'$. Basis $\{1, x, x^2, x^3\}$.

Example 19: Differentiation as Linear Transformation

Problem: $T : P_3 \rightarrow P_2$ (polynomials degree 3 to 2) defined by $T(p) = p'$. Basis $\{1, x, x^2, x^3\}$.

$$T(1) = 0, \quad T(x) = 1, \quad T(x^2) = 2x, \quad T(x^3) = 3x^2$$

Matrix (using basis $\{1, x, x^2\}$ for codomain):

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix} \quad (3 \times 4)$$

Example 19: Differentiation as Linear Transformation

Problem: $T : P_3 \rightarrow P_2$ (polynomials degree 3 to 2) defined by $T(p) = p'$. Basis $\{1, x, x^2, x^3\}$.

$$T(1) = 0, \quad T(x) = 1, \quad T(x^2) = 2x, \quad T(x^3) = 3x^2$$

Matrix (using basis $\{1, x, x^2\}$ for codomain):

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix} \quad (3 \times 4)$$

Apply to $p(x) = 2 + 3x + 4x^2 + 5x^3$: coefficients $(2, 3, 4, 5)^T$.

$$A \begin{pmatrix} 2 \\ 3 \\ 4 \\ 5 \end{pmatrix} = \begin{pmatrix} 3 \\ 8 \\ 15 \end{pmatrix} \Rightarrow 3 + 8x + 15x^2$$

Check derivative: $p'(x) = 3 + 8x + 15x^2$

Example 20: Integration as Linear Transformation

Problem: $T : P_2 \rightarrow P_3$, $T(p) = \int_0^x p(t)dt$. Basis $\{1, x, x^2\}$.

Example 20: Integration as Linear Transformation

Problem: $T : P_2 \rightarrow P_3$, $T(p) = \int_0^x p(t)dt$. Basis $\{1, x, x^2\}$.

$$T(1) = x, \quad T(x) = \frac{x^2}{2}, \quad T(x^2) = \frac{x^3}{3}$$

Matrix (basis $\{1, x, x^2, x^3\}$):

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1/3 \end{pmatrix} \quad (4 \times 3)$$

Example 20: Integration as Linear Transformation

Problem: $T : P_2 \rightarrow P_3$, $T(p) = \int_0^x p(t)dt$. Basis $\{1, x, x^2\}$.

$$T(1) = x, \quad T(x) = \frac{x^2}{2}, \quad T(x^2) = \frac{x^3}{3}$$

Matrix (basis $\{1, x, x^2, x^3\}$):

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1/3 \end{pmatrix} \quad (4 \times 3)$$

Apply to $p(x) = 4 + 5x + 6x^2$: $(4, 5, 6)^T$

$$T = \begin{pmatrix} 0 \\ 4 \\ 5/2 \\ 2 \end{pmatrix} \Rightarrow 0 + 4x + \frac{5}{2}x^2 + 2x^3$$

Check: $\int_0^x (4 + 5t + 6t^2)dt = 4x + \frac{5}{2}x^2 + 2x^3$

Summary Table of Transformations

Type	Matrix (2D)	Effect
Identity	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	No change
Rotation θ	$\begin{pmatrix} c & -s \\ s & c \end{pmatrix}$	Rotate by θ
Scale a, b	$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$	Stretch/compress
Shear x	$\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$	Horizontal skew
Reflect x-axis	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	Flip vertically
Project onto x	$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$	Drop y-coordinate

Summary Table of Transformations

Type	Matrix (2D)	Effect
Identity	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	No change
Rotation θ	$\begin{pmatrix} c & -s \\ s & c \end{pmatrix}$	Rotate by θ
Scale a, b	$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$	Stretch/compress
Shear x	$\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$	Horizontal skew
Reflect x-axis	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	Flip vertically
Project onto x	$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$	Drop y-coordinate

Key: $c = \cos \theta$,

$s = \sin \theta$; k = shear factor.

Matrix of Transformation: Change of Basis

If B and C are bases for domain and codomain:

$$[T]_{C \leftarrow B} = ([T(\mathbf{b}_1)]_C \quad \cdots \quad [T(\mathbf{b}_n)]_C)$$

Matrix of Transformation: Change of Basis

If B and C are bases for domain and codomain:

$$[T]_{C \leftarrow B} = ([T(\mathbf{b}_1)]_C \quad \cdots \quad [T(\mathbf{b}_n)]_C)$$

Example: $T(x, y) = (2x + y, x - y)$ with $B = \{(1, 1), (1, -1)\}$, C standard.

Matrix of Transformation: Change of Basis

If B and C are bases for domain and codomain:

$$[T]_{C \leftarrow B} = ([T(\mathbf{b}_1)]_C \quad \cdots \quad [T(\mathbf{b}_n)]_C)$$

Example: $T(x, y) = (2x + y, x - y)$ with $B = \{(1, 1), (1, -1)\}$, C standard.

$$T(1, 1) = (3, 0) \rightarrow [3, 0]^T, \quad T(1, -1) = (1, 2) \rightarrow [1, 2]^T.$$

$$[T]_{C \leftarrow B} = \begin{pmatrix} 3 & 1 \\ 0 & 2 \end{pmatrix}$$

Matrix of Transformation: Change of Basis

If B and C are bases for domain and codomain:

$$[T]_{C \leftarrow B} = ([T(\mathbf{b}_1)]_C \quad \cdots \quad [T(\mathbf{b}_n)]_C)$$

Example: $T(x, y) = (2x + y, x - y)$ with $B = \{(1, 1), (1, -1)\}$, C standard.

$$T(1, 1) = (3, 0) \rightarrow [3, 0]^T, \quad T(1, -1) = (1, 2) \rightarrow [1, 2]^T.$$

$$[T]_{C \leftarrow B} = \begin{pmatrix} 3 & 1 \\ 0 & 2 \end{pmatrix}$$

$$\text{If } \mathbf{v} = 2(1, 1) + 1(1, -1) = (3, 1): \quad T(\mathbf{v}) = \begin{pmatrix} 3 & 1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 7 \\ 2 \end{pmatrix}. \text{ Direct:}$$

$$T(3, 1) = (7, 2)$$

Inverse Linear Transformations

$T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is invertible if matrix A is invertible ($\det A \neq 0$).

Inverse Linear Transformations

$T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is invertible if matrix A is invertible ($\det A \neq 0$).

$$T^{-1}(\mathbf{y}) = A^{-1}\mathbf{y}$$

Inverse Linear Transformations

$T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is invertible if matrix A is invertible ($\det A \neq 0$).

$$T^{-1}(\mathbf{y}) = A^{-1}\mathbf{y}$$

Example: $T(x, y) = (x + 2y, 3x + 4y)$.

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}, \quad \det = -2 \neq 0$$

Inverse Linear Transformations

$T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is invertible if matrix A is invertible ($\det A \neq 0$).

$$T^{-1}(\mathbf{y}) = A^{-1}\mathbf{y}$$

Example: $T(x, y) = (x + 2y, 3x + 4y)$.

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}, \quad \det = -2 \neq 0$$

$$A^{-1} = -\frac{1}{2} \begin{pmatrix} 4 & -2 \\ -3 & 1 \end{pmatrix} = \begin{pmatrix} -2 & 1 \\ 1.5 & -0.5 \end{pmatrix}$$

Inverse Linear Transformations

$T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is invertible if matrix A is invertible ($\det A \neq 0$).

$$T^{-1}(\mathbf{y}) = A^{-1}\mathbf{y}$$

Example: $T(x, y) = (x + 2y, 3x + 4y)$.

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}, \quad \det = -2 \neq 0$$

$$A^{-1} = -\frac{1}{2} \begin{pmatrix} 4 & -2 \\ -3 & 1 \end{pmatrix} = \begin{pmatrix} -2 & 1 \\ 1.5 & -0.5 \end{pmatrix} \text{ So}$$

$$T^{-1}(u, v) = (-2u + v, 1.5u - 0.5v). \text{ Check: } T(1, 0) = (1, 3),$$

$$T^{-1}(1, 3) = (-2 + 3, 1.5 - 1.5) = (1, 0)$$

Eigenvalues and Eigenvectors of Transformations

For $T : V \rightarrow V$, λ is eigenvalue if $T(\mathbf{v}) = \lambda\mathbf{v}$ for some $\mathbf{v} \neq 0$.

Eigenvalues and Eigenvectors of Transformations

For $T : V \rightarrow V$, λ is eigenvalue if $T(\mathbf{v}) = \lambda\mathbf{v}$ for some $\mathbf{v} \neq 0$.

Example: $T(x, y) = (3x + y, y)$ from earlier.

$$A = \begin{pmatrix} 3 & 1 \\ 0 & 1 \end{pmatrix}$$

Eigenvalues and Eigenvectors of Transformations

For $T : V \rightarrow V$, λ is eigenvalue if $T(\mathbf{v}) = \lambda\mathbf{v}$ for some $\mathbf{v} \neq 0$.

Example: $T(x, y) = (3x + y, y)$ from earlier.

$$A = \begin{pmatrix} 3 & 1 \\ 0 & 1 \end{pmatrix}$$

Eigenvalues: $\lambda = 3$ (eigenvector $(1, 0)$), $\lambda = 1$ (eigenvector $(-1, 2)$).

Eigenvalues and Eigenvectors of Transformations

For $T : V \rightarrow V$, λ is eigenvalue if $T(\mathbf{v}) = \lambda\mathbf{v}$ for some $\mathbf{v} \neq 0$.

Example: $T(x, y) = (3x + y, y)$ from earlier.

$$A = \begin{pmatrix} 3 & 1 \\ 0 & 1 \end{pmatrix}$$

Eigenvalues: $\lambda = 3$ (eigenvector $(1, 0)$), $\lambda = 1$ (eigenvector $(-1, 2)$).

Check $T(-1, 2) = (-3 + 2, 2) = (-1, 2) = 1 \cdot (-1, 2)$

Geometric Transformations in Computer Graphics

Common 3D transformations (homogeneous coordinates):

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & t_x \\ a_{21} & a_{22} & a_{23} & t_y \\ a_{31} & a_{32} & a_{33} & t_z \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

Geometric Transformations in Computer Graphics

Common 3D transformations (homogeneous coordinates):

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & t_x \\ a_{21} & a_{22} & a_{23} & t_y \\ a_{31} & a_{32} & a_{33} & t_z \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

Translation: t_x, t_y, t_z ; Rotation: top-left 3×3 orthogonal matrix.

Geometric Transformations in Computer Graphics

Common 3D transformations (homogeneous coordinates):

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & t_x \\ a_{21} & a_{22} & a_{23} & t_y \\ a_{31} & a_{32} & a_{33} & t_z \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

Translation: t_x, t_y, t_z ; Rotation: top-left 3×3 orthogonal matrix.

Example: Translate (1, 2, 3) by (4, 5, 6):

$$\begin{pmatrix} 1 & 0 & 0 & 4 \\ 0 & 1 & 0 & 5 \\ 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \\ 1 \end{pmatrix} = \begin{pmatrix} 5 \\ 7 \\ 9 \\ 1 \end{pmatrix}$$

Summary and Key Insights

Every linear transformation:

- Is represented by a matrix
- Satisfies
$$T(c\mathbf{u} + \mathbf{v}) = cT(\mathbf{u}) + T(\mathbf{v})$$
- Has kernel and image
- Composition = matrix multiplication

Summary and Key Insights

Every linear transformation:

- Is represented by a matrix
- Satisfies
$$T(c\mathbf{u} + \mathbf{v}) = cT(\mathbf{u}) + T(\mathbf{v})$$
- Has kernel and image
- Composition = matrix multiplication

Dimensions matter:

$$\underbrace{\mathbb{R}^n}_{\text{domain}} \xrightarrow{A_{m \times n}} \underbrace{\mathbb{R}^m}_{\text{codomain}}$$

Summary and Key Insights

Every linear transformation:

- Is represented by a matrix
- Satisfies
$$T(c\mathbf{u} + \mathbf{v}) = cT(\mathbf{u}) + T(\mathbf{v})$$
- Has kernel and image
- Composition = matrix multiplication

Dimensions matter:

$$\underbrace{\mathbb{R}^n}_{\text{domain}} \xrightarrow{A_{m \times n}} \underbrace{\mathbb{R}^m}_{\text{codomain}}$$

Special cases:

- $m = n$: square matrix
- $m < n$: projection/loss of info
- $m > n$: embedding

Summary and Key Insights

Every linear transformation:

- Is represented by a matrix
- Satisfies
$$T(c\mathbf{u} + \mathbf{v}) = cT(\mathbf{u}) + T(\mathbf{v})$$
- Has kernel and image
- Composition = matrix multiplication

Dimensions matter:

$$\underbrace{\mathbb{R}^n}_{\text{domain}} \xrightarrow{A_{m \times n}} \underbrace{\mathbb{R}^m}_{\text{codomain}}$$

Special cases:

- $m = n$: square matrix
- $m < n$: projection/loss of info
- $m > n$: embedding

Linear transformations are the language of data science, graphics, and machine learning!

Final Practice Problem

Find the matrix of $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that:

- 1 Reflects through xz -plane ($y \rightarrow -y$)
- 2 Then rotates 90° around x -axis

Final Practice Problem

Find the matrix of $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that:

- 1 Reflects through xz -plane ($y \rightarrow -y$)
- 2 Then rotates 90° around x -axis

Solution: Reflection: $R = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$

Final Practice Problem

Find the matrix of $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that:

- 1 Reflects through xz -plane ($y \rightarrow -y$)
- 2 Then rotates 90° around x -axis

Solution: Reflection: $R = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ Rotation around x -axis by 90° :

$$Q = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}$$

Final Practice Problem

Find the matrix of $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that:

- 1 Reflects through xz -plane ($y \rightarrow -y$)
- 2 Then rotates 90° around x -axis

Solution: Reflection: $R = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ Rotation around x -axis by 90° :

$Q = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}$ Composition (rotate after reflect):

$$A = QR = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}$$

Final Practice Problem

Find the matrix of $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that:

- 1 Reflects through xz -plane ($y \rightarrow -y$)
- 2 Then rotates 90° around x -axis

Solution: Reflection: $R = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ Rotation around x -axis by 90° :

$Q = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}$ Composition (rotate after reflect):

$A = QR = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}$ Check on $(0, 1, 0)$: reflect $\rightarrow (0, -1, 0)$,
rotate $\rightarrow (0, 0, -1)$. Direct: $A(0, 1, 0) = (0, 0, -1)$

End — Thank You!