

# 1 Background

Consider the PDE and the ODE

$$\frac{\partial}{\partial t}T(x, t) = S(x, t)\frac{\partial^2}{\partial x^2}T(x, t), \quad \frac{\partial}{\partial t}S(x, t) = T(x, t) - S(x, t), \quad \forall x, t$$

Then with symmetric differences

$$\frac{\partial}{\partial t}T(x, t) = S(x, t)\frac{T(x-h, t) - 2T(x, t) + T(x+h, t)}{h^2} + \mathcal{O}(h^4).$$

We now discretize the state space  $\Omega$  by  $[x_0, \dots, x_{n+1}]$  and define  $s_i(t) := S(x_i, t)$  and  $T_i(t) := T(x_i, t)$  and then get

$$\frac{d}{dt} \begin{pmatrix} T_1 \\ S_1 \\ \vdots \\ T_n \\ S_n \end{pmatrix} = \begin{pmatrix} S_1(t)\frac{T_0(t)-2T_1(t)+T_2(t)}{h^2} + \mathcal{O}(h^4) \\ T_1(t) - S_1(t) \\ \vdots \\ S_n(t)\frac{T_{n-1}(t)-2T_n(t)+T_{n+1}(t)}{h^2} + \mathcal{O}(h^4) \\ T_n(t) - S_n(t) \end{pmatrix}$$

which is a system of  $2n$  ODEs.