Diffusion Models on Manifolds

Abstract

bla bla

CONTENTS

Contents **1**

1 Stochastic Processes Recap **2**

1.1 Canonical Construction of Stochastic Processes . . . 2
1.2 Brownian Motion . . . 11
1.3 Bronwian Family . . . 11
1.4 Weak Solutions by Means of the Girsanov Theorem . . . 12
1.5 Stochastic Calculus Backround . . . 13

2 Girsanov Theorem **17**

3 Weak Solutions **19**

3.1 Weak Solutions by Means of the Girsanov Theorem . . . 21

4 The Martingale Problem of Stroock and Varadhan **22**

4.1 Fundamentals and Motiviation: . . . 23
4.2 Weak Solutions and the Martingale Problem . . . 25

5 Markov Processes **30**

SECTION 1

STOCHASTIC PROCESSES RECAP

Theorem 1.1

Let $(X^n)_{n\in\mathbb{N}}$ be a tight sequence of continuous processes such that

$$\forall 0 \le t_1 < ... < t_d < \infty : \quad (X^n_{t_1}, ..., X^n_{t_d})_{n\in\mathbb{N}}$$

converges in distribution, i.e.

$$w - \lim_{n\to\infty} \mathbb{P} \circ (X^n_{t_1}, ..., X^n_{t_d})^{-1} = \lim_{n\to\infty} \int_{\mathbb{R}^d} df \mathbb{P} \circ (X^n_{t_1}, ..., X^n_{t_d})^{-1}, \quad \forall f : \mathbb{R}^d \to \mathbb{R} \text{ bounded and continuous}$$

exists. Then

- $w - \lim_{n\to\infty} \mathbb{P}_n := w - \lim_{n\to\infty} \mathbb{P} \circ (X^n_{t_1}, ..., X^n_{t_d})^{-1} = P$
- On P the coordinate mapping process $W_t(\omega) = \omega(t)$ on $C([0,\infty))$ satisfies

$$w - \lim_{n\to\infty} \mathbb{P} \circ (X^n_{t_1}, ..., X^n_{t_d})^{-1} = P \circ (W_{t_1}, ..., W_{t_d})^{-1}, \quad \forall 0 \le t_1 < ... < t_d < \infty.$$

Proof. Let $(X^{n_m})_{m\in\mathbb{N}}$ be a subseqeuence of $(X^n)_{n\in\mathbb{N}}$. Because this subseqeunce is again tight (subsequences can be chosen arbitrarly) it has another subsequence $(X^{n_{m_l}})_{l\in\mathbb{N}}$ so that

$$w - \lim_{l\to\infty} \mathbb{P} \circ (X^{n_{m_l}})^{-1} = P.$$

Assume there is another subsequence $(X^{n_{m_k}})_{k\in\mathbb{N}}$ that induces a measure $C([0,\infty))$ with

$$Q := w - \lim_{k\to\infty} \mathbb{P} \circ (X^{n_{m_k}}).$$

Then for any $\pi : C([0,\infty)) \to \mathbb{R}^d$ we have

$$P \circ \pi^{-1} = Q \circ \pi^{-1},$$

yielding $Q = P$, which implies the weak convergence of $(P^n)_{n\in\mathbb{N}}$ to P.
Suppose $(\mathbb{P}^n)_{n\in\mathbb{N}}$ did not converge weakly to P, then $\exists f : C([0,\infty)) \to \mathbb{R}$, bounded and continuous such that $\int f d\mathbb{P}_n$ does not exist or does not equal $\int f dP$. In both cases choose a subsequence $(\mathbb{P}^{n_k})_{k\in\mathbb{N}}$ with $\lim_{k\to\infty} \int df d\mathbb{P}^{n_k}$ does not exists or does not equal $\int f dP$. Any further subseqeunce also cannot converge weakly to P. However, in the abstract before it was shown that

$$\lim_{l\to\infty} \int f d\mathbb{P}^{n_{k_l}} = \int f dQ = \int f \, dP,$$

which implies that $(\mathbb{P}^{n_k})_{k\in\mathbb{N}}$ should converge weakly to P, which contradicts the construction before. □

1.1 Canonical Construction of Stochastic Processes

Construction of a stochastic process on I with $|I| = \infty$.

- Define a σ-algebra $\mathcal{E}^{\otimes I}$ on $E^I := \{f : I \to E\}$
- Understand all probability measures on $\mathcal{E}^{\otimes I}$
- Reinterpret stochastic processes as path valued random variables.

For $d < \infty$ the product σ-algebra

$$\mathcal{E}^{\otimes d} := \mathcal{E} \otimes \cdots \otimes \mathcal{E} := \sigma(\{B_1 \times \cdots \times B_d \mid B_i \in \mathcal{E}\}),$$

where $B_1 \times \cdots \times B_d$ are the rectangular sects.

Remark 1.1

If $\mathcal{E} = \sigma(\mathcal{A})$ then

$$\mathcal{E}^{\otimes d} = \sigma(\{B_1 \times \cdots \times B_d \mid B_i \in \mathcal{A}\}).$$

???

We now want to generalize to infinite products.

Definition 1.2

A subset $C \subset E^I$ is called finitely generated cylinder set if

$$C = \pi_J^{-1}(B_1 \times \cdots \times B_n) = \{f : I \to E \mid f(t_1) \in B_1, ..., f(t_n) \in B_n\},$$

where $J := \{t_1, ..., t_n\} \subseteq I$, $B_1, ..., B_n \in \mathcal{E}$ and the finite projection

$$\pi_J : E^I \to E^{|J|}, f \mapsto \pi_J(f) := (f(t_1), ..., f(t_n)).$$

We say the set C of paths is spanned by finitely many time points $t_1, ..., t_n$ and the finite dimensional cylinder $B_1 \times \cdots \times B_n \in \mathcal{E}^{\otimes d}$.

Definition 1.3

The path σ-algebra $\mathcal{E}^{\otimes I}$ on E^I is the smallest σ-algebra that contains all finitely generated cylinder sets. In the case $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ one writes $\mathcal{B}(\mathbb{R}^d), \mathcal{B}(\mathbb{R}^\infty), \mathcal{B}(\mathbb{R}^{[0,T]})$.

A σ-algebra defined using finitely generated cylinder sets can only at most cover countably many restrictions. We call a set finitely generated if there exists $J \subseteq I$ (finite or countable) such that

$$f \in C \quad \iff \quad f\big|_C \in B, \quad B \in \mathcal{E}^{\otimes J}.$$

Note that here we do not only use cylinder sets $B_1 \times \cdots \times B_d$, as in the next proposition shows that we need more than these cylinders sets to form a σ-algebra.

Proposition 1.4

The set

$$\mathcal{V} := \{C \subseteq E^I \mid \exists J \subseteq I \text{ countable}, B \in \mathcal{E}^{\otimes J}, \forall f \in E^I : \quad f \in C \iff f\big|_J \in B\}$$

of all countably generated sets is a σ-algebra on E^I that contains $\mathcal{E}^{\otimes I}$.

Example 1.1

If $E = \mathbb{R}$ and $I \subseteq \mathbb{R}$ such that $|I| = \infty$, then $\{f\}, \{f \equiv c \in E\}, \{f : I \to E \mid f \geq 0\}, C(I, E)$ are not measurable sets. If any of those sets were measurable, then that set should be with the above proposition in $\mathcal{V}$ as well. This means that there should exist $J \subseteq I$ countable and $B \in \mathcal{E}^{\otimes J}$ so that any $f\big|_J \in B$, meaning for $f \in C$, we only need finite many points to be inside a measurable set of E^I. This however fails the definition of these sets: Choose $i \notin J$ and set $f(i)$ in such a way, that it does not satisfy the defining property anymore, e.g. a discontinuous point for example. Then f is no longer in C even though it is measurable. So statements of the $\mathbb{P}$-a.s. kind need to be done with caution.

We now construct the measures on $\mathcal{E}^{\otimes I}$. The idea is as follows

- Downsize $\mathbb{P}$ to only finite cylinder sets (that are $\cap$-stable). This leads to finite-dimensional distributions
- Check that $\{\mathbb{P}_J \mid |J| < \infty\}$ satisfies a natural property, which will be consistency.
- Show that for these finite dimensional consistent families there exists an infinite dimensional probability measure.

Definition 1.5

If $\mathbb{P}$ is a probability measure on $\mathcal{E}^{\otimes I}$ and $J \subseteq I$ is finite, then the unique probability measure on $\mathcal{E}^{\otimes J}$ is given by

$$\mathbb{P}_J(B_1 \times \cdots \times B_{|J|}) := \mathbb{P}(\pi_J^{-1}(B_1 \times \cdots \times B_{|J|})), \quad B_i \in \mathcal{E},$$

is called finite-dimensional marginal distribution of $\mathbb{P}$.

On the one hand we have the boxes, i.e. $B_1, ..., B_n \in \mathcal{E}$ and cylinder sets $\{f : I \to E \mid f(t_1) \in B_1, ..., f(t_n) \in B_n\}$. The finite-dimensional marginals are the measures that we get by crossing out the functions away from J, meaning it is the distribution of $\pi_J(f)$ on these boxes.

Proposition 1.6

Every probability measure $\mathbb{P}$ on $\mathcal{E}^{\otimes I}$ is uniquely determined through the family of all finite-dimensional marginal distributions $\{\mathbb{P}_J \mid J \subseteq I \text{ finite }\}$.

The next result will lead to the characterizing property of finite dimensional families related to measures on $\mathcal{E}^{\otimes I}$.

Proposition 1.7

If $\mathbb{P}$ is a probability measure on $\mathcal{E}^{\otimes I}$ then the family of all finite-dimensional marginal distributions $\{\mathbb{P}_J \mid J \subseteq I \text{ finite }\}$ satisfies for $\pi_{J,J'} : E^J \to E^{J'}$

$$\forall J' \subseteq J \subseteq I \text{ finite with } \pi_{J,J'}(f) = f\big|_{|J|} : \quad \mathbb{P}_J \circ \pi_{J,J'}^{-1} = \mathbb{P}_{J'}.$$

Definition 1.8

A set of finite-dimensional distributions $\{\mathbb{P}_J \mid |J| < \infty\}$ on I is called consistent if

$$\forall J' \subseteq J : \quad \mathbb{P}_J \circ \pi_{J,J'}^{-1} = \mathbb{P}_{J'}.$$

This means "adding" E's to a product $B_1 \times \cdots \times B_n \times E$ does not change the measure. For the infinite dimensional distribution we do not need more properties for the existence of the infinite dimensional distribution.

Theorem 1.9

Let E be a polish metric space, $\mathcal{E} = \mathcal{B}(E)$ and I an index set. If $\{\mathbb{P}_J \mid J \subseteq I \text{ finite}\}$ is a consistent family then there exists a unqiue $\mathbb{P}$ on $\mathcal{E}^{\otimes}$ such that

$$\forall J \subseteq I \text{ finite:} \quad \mathbb{P} \circ \pi_J^{-1} = \mathbb{P}_J.$$

How should we know $\mathbb{P}_J$ without knowing $\mathbb{P}$?

Definition 1.10

Suppose $(E, \mathcal{E})$ is a measurable space and $\mathbb{P}$ a probability measure on $\mathcal{E}$. Then, a measure μ on $\mathcal{E}^{\otimes \mathbb{N}}$ is called infinite dimensional product measure if

$$\forall n \in \mathbb{N}, B_i \in \mathcal{E}: \quad \mu(B_1 \times \cdots \times B_n \times E \times \cdots) = \mathbb{P}(B_1) \cdots \mathbb{P}(B_n).$$

We typically write for an infinite product measure $\mathbb{P}^{\otimes \infty}$ or $\mathbb{P}^{\otimes \mathbb{N}}$.

Theorem 1.11

If E is polish metric space and $\mathbb{P}$ a probability measure on $\mathcal{B}(E)$, then the infinite dimensional product measure $\mathbb{P}^{\otimes \infty}$ exists and is unique.

Next, we want to understand the connection of stochastic processes and path valued random variables.

Proposition 1.12

$X := (X_t)_{t \in I}$ is E-valued stochastic process on $(\Omega, \mathcal{F}, \mathbb{P})$ if and only if

$$X : \omega \mapsto (t \mapsto X_t(\omega))$$

is a path valued random variable.

Proof. "$\Leftarrow$"
For $B \in \mathcal{E}$

$$X_t^{-1}(B) = X^{-1}(\pi_t^{-1}(B)) \in \mathcal{A},$$

because $\pi_t^{-1}(B)$ is a finitely generated cylinder set and X is $(\mathcal{A}, \mathcal{E}^{\otimes I})$-measurable.
"$\Rightarrow$"
We only need to check measurability for an arbitrary generator. We use the cylinder sets $\pi_t^{-1}(B)$. Then

$$X^{-1}(\pi_t^{-1}(B)) = \{\omega \in \Omega \mid X_t(\omega) \in B\} = X_t^{-1}(B) \in \mathcal{A}.$$

□

Definition 1.13

Suppose X is an E-valued stochastic process on $(\Omega, \mathcal{F}, \mathbb{P})$, then

- The image measure

$$\mathbb{P}^X(A) := \mathbb{P}(X \in A), \quad A \in \mathcal{E}^{\otimes I},$$

is called law of X.

- For $J := \{t_1, ..., t_n\} \subseteq I$ we call

$$\mathbb{P}^X_{t_1,...,t_n} \sim (X_{t_1}, ..., X_{t_n})$$

is finite dimensional distribution (fdd) of X.

Usually, the law does not play a big role and we exclusively deal with the fdds.

Proposition 1.14

The law of a stocahstic process X is uniquely determined by all its finite-dimensional distributions

$$\mathbb{P}^X_{t_1,...,t_n} = \mathbb{P}(X_{t_1} \in \cdot, ..., X_{t_n} \in \cdot).$$

Definition 1.15

Two stochastic processes X and Y with index set I have the same law if $\mathbb{P}^X = \mathbb{P}^Y$. We say they have the same finite-dimensional distribution if

$$\forall \{t_1, ..., t_n\} \subseteq I : \quad \mathbb{P}^X_{t_1,...,t_n} = \mathbb{P}^Y_{t_1,...,t_n}$$

and write $\mathbb{P}^X \overset{fdd}{=} \mathbb{P}^Y$.

Theorem 1.16

Suppose E is a polish metric space, I an index set and $\{\mathbb{P}_J \mid J \subseteq I \text{ finite}\}$ a consistent family of finite-dimensional distributions, then there exists a stochastic process X on $(\Omega, \mathcal{F}, \mathbb{P})$ so that

$$\forall J \subseteq I : \quad \mathbb{P}^X_J = \mathbb{P}_J.$$

One tries to avoid the canonical construction, because statements such as "X is $\mathbb{P}$-a.s. continuous/..." are problematic on the canonical space. This is why we now want to derive continuous modifications. For that we need the stochastic process to be a random variable in a polish space. First, we consider the path space of continuous functions and stochastic processes with continuous oaths. Second, we show how to construct these. We cannot use Kolmogorov's extension theorem, as it breaks down for continuous functions. Note that it is already known that $(C([0,T]), \|\cdot\|_\infty)$ is a polish space.

Proposition 1.17

- $$d(f,g) := \sum_{i=1}^{k} \frac{d : k(f,g) \vee 1}{2^k}, \quad d_k(f,g) := \sup_{x \in [0,k]} |f(x) - g(x)|,$$

is a metric on $C([0,\infty))$.

- $\lim_{n\to\infty} d(f_n, g) = 0 \iff \lim_{n\to\infty} \sup_{x\in K} |f_n(x) - g(x)| = 0$ for every K compact.
- $(C([0,\infty)), d)$ is a polish metric space.

The product σ-algebra for $\mathbb{R}^{[0,\infty)}$ has no structure, however, the Borel σ-algebra generated by open sets

for $C([0,\infty))$ has more, because for the product σ-algebra can only have measurable sets, that have finite restrictions. Because the continuous functions are seperable, there are not any for the Borel σ-algebra on open sets.
Cylinder sets of continuous functions are also the generators now:

Proposition 1.18

$\mathcal{B}(C([0,\infty)))$ is generated by cylinder sets of continuous functions, i.e.

$$\begin{aligned}\mathcal{B}(C([0,\infty))) &= \sigma(\{\pi_t^{-1}(B) \mid t \geq 0, B \in \mathcal{B}(\mathbb{R})\}) \\ &= \sigma(\{\text{ finitely generated cylinder sets of continuous functions}\}),\end{aligned}$$

where $\pi_t : C([0,\infty)) \to \mathbb{R}, f \mapsto f_t$.

Because $\mathcal{B}(C([0,\infty)))$ has the same type of generator as $\mathcal{E}^{\otimes I}$ we can copy ideas for probability measures from before. If $\mathbb{P}$ is a probability measure on $\mathcal{B}(C([0,\infty)))$, then for every $t_i \geq 0$ and $B_i \in \mathcal{B}(\mathbb{R})$ we have

$$\mathbb{P}_{t_1,\dots,t_n}(B_1 \times \cdots \times B_k) := \mathbb{P}(\{f \in C([0,\infty)) \mid f(t_1) \in B_1, \dots, f(t_k) \in B_k\}),$$

is called the finite dimensional distribution. Since this probability measure is defined on a $\cap$-stable generator, then finite dimensional distribution uniquely determines the law.

Corollary 1.19

A probability measure $\mathbb{P}$ on $\mathcal{B}(C([0,\infty)))$ is uniquely determined by the family $\{\mathbb{P}_{t_1,\dots,t_k} \mid k \in \mathbb{N}, t_i \geq 0\}$ of all finite-dimensional distributions.

Proposition 1.20

$\mathcal{B}(\mathbb{R}^{[0,\infty)})\big|_{C([0,\infty))} = \mathcal{B}(C([0,\infty)))$.

we cannot use Kolmogorovs extension theorem to construct measures on $\mathcal{B}(C([0,\infty)))$.

Definition 1.21

Suppose X is a stochastic process on $(\Omega, \mathcal{F}, \mathbb{P})$ then

- X is continuous/differentiable/... if $\omega \mapsto X_t(\omega)$ is continuous/differentiable/... for all $\omega \in \Omega$.
- X is *a.s.* continuous/differentiable/... if there exists $C \in \mathcal{F}$ with $\mathbb{P}(C) = 1$ such that $t \mapsto X_t(\omega)$ is continuous/differentiable/... for every $c \in C$.

We have already defined the law $\mathbb{P}^X(B) := \mathbb{P}(X \in B)$ on the product σ-algebra. It would be nice to define it on $C([0,\infty))$ as well. This is why we need $X : \Omega \to C([0,\infty)) \subset \mathbb{R}^{[0,\infty)}$ to be $(\mathcal{F}, \mathcal{E}^{\otimes I}$ and $(\mathcal{F}, \mathcal{B}(C([0,\infty))))$ measurable. This is because if $B \in \mathcal{B}(C([0,\infty)))$ then there exists $B' \in \mathcal{E}^{\otimes I}$ with $B = B' \cap C([0,\infty))$ and it holds

$$X^{-1}(B) = X^{-1}(B) \cap X^{-1}(C([0,\infty))) = X^{-1}(B') \cap \Omega \in \mathcal{F}.$$

Hence, introducing continuous processes can be interpreted as random variables taking values in $C([0,\infty))$ and inducing a law on $\mathcal{B}(C([0,\infty)))$.

Definition 1.22

If X is a continuous stochastic process then the law $\mathbb{P}^{X,c}$ of X on $\mathcal{B}(C([0,\infty)))$ is

$$\mathbb{P}^{X,c}(B) := \mathbb{P}(X \in B), \quad B \in \mathcal{B}(C([0,\infty))).$$

We now construct the continuous stochastic processes.

Definition 1.23

A function $f : [0,\infty) \to \mathbb{R}$ is called Hölder continuous with index $\gamma \in (0,1]$ and constant K if

$$\forall s,t \in \mathbb{R} : \quad |f(t) - f(s)| \leq k|t-s|^{\gamma}.$$

We call f Hölder continuous on compacts on $\gamma \in (0,1]$ with $(K_n)_{n\in\mathbb{N}}$ if

$$\forall s,t \in \mathbb{N} : \quad |f(t) - f(s)| \leq K_N|t-s|^{\gamma},$$

for all $N \in \mathbb{N}$. f is locally Hölder continuous of index $\gamma \in (0,1]$ if

$$\forall t \in \mathbb{N} \exists \mathcal{U}_t \subseteq \mathbb{N}, \text{ with } t \in \mathcal{U}_t \forall s \in \mathcal{U}_t : \quad |f(t) - f(s)| \leq K_t|t-s|^{\gamma}.$$

f is Hölder continuous in t of index $\gamma \in (0,1]$ if

$$\exists \mathcal{U}_t \ni t \in \mathbb{N} \text{ and } K_t : \quad |f(t) - f(s)| \leq K_t|t-s|^{\gamma}, \quad s \in ???.$$

Definition 1.24

Let X and $\tilde{X}$ be a stochastic process with index set I on $(\Omega, \mathcal{F}, \mathbb{P})$. Then $\tilde{X}$ is a modification of X if

$$\forall t \in I : \quad \mathbb{P}(X_t = \tilde{X}_t) = 1.$$

Definition 1.25

Let X and $\tilde{X}$ be a stochastic process with index set I on $(\Omega, \mathcal{F}, \mathbb{P})$. Then X and $\tilde{X}$ are called indistinguishable if there exists $C \in \mathcal{F}$ with $\mathbb{P}(C) = 1$ so that for every $\omega \in \Omega$: $X_t(\omega) = \tilde{X}_t(\omega)$.

Modifications of erach other is stronger than equality in law. Indistinguishable is the strongest here.

Example 1.2

Suppose $U \sim \mathcal{U}([0,1])$ on $(\Omega, \mathcal{F}, \mathbb{P})$ and define $X_t := 0$ and $\tilde{X}_t := \mathbf{1}_U(f)$ on $I = [0,1]$. Then $\tilde{X}$ is a modification of X, because

$$\mathbb{P}(X_t = \tilde{X}_t) = \mathbb{P}(U \neq t) = 1.$$

However, their paths are a.s. different.

Theorem 1.26

Suppose X is a stochastic process on $(\Omega, \mathcal{F}, \mathbb{P})$ such that for all $T > 0$ exists c, α, β

$$\forall s,t \leq T : \quad \mathbb{E}[|X_t - X_s|^{\alpha}] \leq C|t-s|^{1+\beta}.$$

Then there exists $\tilde{X}$ a continuous modification of X s.t. $\tilde{X}$ has Hölder continuous paths of order $\gamma < \frac{\beta}{\alpha}$.

Finally, we now consider weak convergence of continuous processes

Proposition 1.27

Let $\mu, \mu^1, ...$ be sequence of probability measures on $\mathcal{B}(C([0,\infty)))$, then $\mu^n \overset{weak}{\to} \mu$, $n \to \infty$ if and only if

- $\{\mu^n \mid n \in \mathbb{N}\}$ is tight in $\mathcal{M}_1(C([0,\infty)))$
- All finite-dimensional distributions converge, i.e.

$$\forall J \subseteq [0,\infty) \text{ finite:} \quad \mu_J^n := \mu^n \circ \pi_J^{-1} \overset{weak}{\to} \mu \circ \pi_J^{-1} = \mu_J, \quad n \to \infty.$$

We are mostly interested in the stochastic reformulation and not in the abstract measure theory.

Definition 1.28

Suppose $X, X^1, ...$ are stochastic processes with *a.s.* continuous paths and laws $\mathbb{P}^X, \mathbb{P}^{X^1}, ...$ on $C([0,\infty))$.

- $(X^n)_{n\in\mathbb{N}}$ converges weakly to X if $\mathbb{P}_{X^n} \overset{weak}{\to} \mathbb{P}_X$, $n \to \infty$. We write $X^n \Rightarrow X$.
- The finite-dimensional distributions (fdds) of $(X^n)_{n\in\mathbb{N}}$ converge to those of X if for every $0 \le t_1 \le ... \le t_n$

  $$(X_{t_1}^n, ..., t_{t_k}^n) \overset{(d)}{\to} (X_{t_1}, ..., X_{t_k}), n \to \infty$$

  or equivalently

  $$\mathbb{P}^{X^n}_{t_1,...,t_k} \overset{weak}{\to} \mathbb{P}_{t_1,...,t_k}, n \to \infty.$$

  We write $X \overset{fdd}{\Rightarrow} X$.

Remark 1.2

$$X^n \Rightarrow X, \quad n \to \infty \quad \overset{cont.MappingThm}{\Rightarrow} \quad X^n \overset{fdd}{\Rightarrow} X, n \to \infty.$$

The continuous mapping theorem yields that weak convergence implies that

$$\forall F : C([0,\infty)) \to \mathbb{R} : \quad F(X^n) \overset{(d)}{=} F(X),$$

whereas convergence in ffd only implies that

$$\forall F : \mathbb{R}^k \to \mathbb{R} \text{ continuous:} \quad F(X_{t_1}^n, ..., t_{t_k}^n) \overset{(d)}{=} F(X_{t_1}, ..., X_{t_k}).$$

Theorem 1.29

A sequence $(X^n)_{n\in\mathbb{N}}$ of continuous stochastic processes weakly converges to X if and only if

- $\{\mathbb{P}^{X^n} \mid n \in \mathbb{N}\} \subseteq \mathcal{M}_1(C([0,\infty)))$ is tight

- $X^n \overset{ffd}{\Longrightarrow} X$, $n \to \infty$.

Recall A is called relatively compact if all sequences have convergent subsequences in A. It is relatively compact if all sequences have convergent subsequences with the limit not needing to belong to A.

Remark 1.3

Ascela-Ascoli: $A \subseteq C([0,\infty))$ is relatively compact if and only if

- $\{f(0) \mid f \in A\} \subseteq \mathbb{R}$ is bounded
- For every $N \in \mathbb{N}$ and $V_f^N(\delta) := \sup_{\substack{|t-s|<\delta \\ s,t \leq N}} |f(t) - f(s)| : \lim_{\delta \to 0} \sup_{f \in A} V_f^N(\delta) = 0$.

So two things can go wrong. Either functions in A are not sufficiently bounded or there are regularity holes/not all functions have the same kind of regularity.

Example 1.3

The function

$$g_n(x) := \begin{cases} x^{1/n} & , x \leq 1 \\ 2 - x & , x \in (1,2) \\ 0 & , x \geq 2 \end{cases}$$

The set $A := \{g_n \mid n \in \mathbb{N}\}$ is neither compact nor relatively compact.

- We need that every sequence has a subsequence that convergence to a $g \in C([0,\infty])$ however,

$$\lim_{n \to \infty} g_n(0) = 0 \text{ and } \forall x \in (0,1] : \quad \lim_{n \to \infty} g_n(x) = 1.$$

  So the limit is discontinuous at zero. So its not compact

- It also fails the second bullet point of Ascela-Ascoli:

$$\forall \epsilon > 0, s \in (0, \epsilon] : \quad \sup_{n \in \mathbb{N}} |g_n(s) - g_n(0)| = 1.$$

We see that this family of functions is for every $n \in \mathbb{N}$ Hölder continuous with index $\gamma = \frac{1}{n}$:

$$|g_n(y) - g_n(x)| \leq |x - y|^{\frac{1}{n}}, \quad \forall x, y \in [0,1].$$

What goes wrong here? Is it the boundedness or regularity? The missing piece is Hölder continuity where the index is independent of $n \in \mathbb{N}$.

Lemma 1.30

Suppose $(K_N)_{N \in \mathbb{N}}$ are non-negative and $\gamma \in (0,1]$, then

$A_K^\gamma := \{f : [0,\infty) \to \mathbb{R} \mid f$ Hölder continuous on compacts with index γ, constant K_N and $|f(0)| \leq K_0\}$

is compact in $C([0,\infty))$.

Theorem 1.31

Suppose $(X^n)_{n\in\mathbb{N}}$ is a sequence of continuous stochastic processes on $(\Omega, \mathcal{F}, \mathbb{P})$ such that

- $(\mathbb{P}^{X_0^n})_{n\in\mathbb{N}}$ is tight in $\mathcal{M}_1(\mathbb{R})$
- For every $n \in \mathbb{N}$ and $T > 0$ exists c, α, β independent of $n \in \mathbb{N}$ with

$$\forall s, t \leq T: \quad \mathbb{E}[|X_t^n - X_s^n|^\alpha] \leq C|t-s|^{1+\beta}.$$

Then $(\mathbb{P}^{X^n})_{n\in\mathbb{N}}$ is tight in $\mathcal{M}_1(C([0,\infty)))$.

So previously we had Ascela-Ascoli as a tool to construct compact sets to check for tightness, i.e.

$$\forall \epsilon > 0 \exists K \text{ compact subsets: } \inf_{n\in\mathbb{N}} \mathbb{P}^{X^n}(K) > 1-\epsilon \quad \iff \quad (\mathbb{P}^{X^n})_{n\in\mathbb{N}} \text{ tight in } \mathcal{M}_1(C([0,\infty))).$$

There we would consider

$$K := \{f \in C([0,\infty)) \mid |f(0)| \leq a, \forall N \in \mathbb{N}, \delta > 0 : V_f^N(\delta) \to 0, \delta \to 0\}$$

which leads with the above to an infinite dimensional joint distribution of $(X_t^n)_{t\geq 0}$ integral, when computing the tail probability. Kolmogorov Chensov avoids this, by only considering a bi variate expectation.

1.2 Brownian Motion

1.3 Bronwian Family

Theorem 1.32

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space on which $(\xi_i)_{i\in\mathbb{N}}$ is an iid sequence with mean zero and finite variance. Define $X_t^n := \frac{Y_{nt}}{\sigma\sqrt{n}}$ and $P_n := \mathbb{P} \circ (X^n)^{-1}$ on $(C([0,\infty)), \mathcal{B}(C([0,\infty)))$. Then $w - \lim_{n\to\infty} P_n = P_*$, where $W \sim P_*$ is the coordinate mapping process on $C([0,\infty))$ is a one-dimensional Brownian.

Definition 1.33

The probability measure P_* on $(C([0,\infty)), \mathcal{B}(C[0,\infty)))$ under which $W_t(\omega) = \omega(t)$, $t \in [0,\infty)$, is a standard one dimensional Brownian motion, called Wiener measure.

Let $P^1, ..., P^d$ be d copies of the Wiener measure, then $P^0 := P^1 \times \cdots \times P^d$ is a measure on $(C([0,\infty)^d), \mathcal{B}(C([0,\infty)^d)))$ and $B_t(\omega) = \omega(t)$ on P^0 with $(\mathcal{F}_t^B)_{t\geq 0}$ is a d-dimensional Brownian motion starting at the origin. Define

$$\forall x \in \mathbb{R}^d, F \in \mathcal{B}(C([0,\infty)^d)): \quad P^x(F) := P^0(F - x),$$

a prbability measure on $(C([0,\infty)^d), \mathcal{B}(C([0,\infty)^d)))$, where

$$F - x := \{\omega \in C([0,\infty)^d) \mid \omega(\cdot) + x \in F\}.$$

Then under P^x, $(B_t)_{t\geq 0}$ is a d dimensional Brownian motion starting in x. For a probability measure μ on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ on $\mathcal{B}(C([0,\infty)^d))$ define

$$P^\mu(F) := \int_{\mathbb{R}^d} P^x(F) d\mu, \quad \text{on } \mathcal{B}(C([0,\infty)^d)).$$

Lemma 1.34

$\forall F \in \mathcal{B}(C([0,\infty)^d))$ the mapping $x \mapsto P^x F$ is $\mathcal{B}(\mathbb{R}^d)$, $\mathcal{B}([0,1])$ measurable.

Proof. Define $\mathcal{C}$ as the family of all cylinder sets in Ω, constructed by

$$C := \{\omega \in \Omega \mid \omega(t_1) \in B_1, ..., \omega(t_n) \in B_n\}, \quad \forall 0 \leq t_1 < ... < t_n < \infty, B_1, ..., B_n \in \mathcal{B}(\mathbb{R}^d).$$

Then $\mathcal{C}$ is $\cap$-stable and generates $\sigma(\mathcal{C}) = \mathcal{F}$. For, fixed $C \in \mathcal{C}$ $x \mapsto P^x(C)$ is $\mathcal{B}(\mathbb{R}^d)$ measurable, because we can formulate $P^x(C)$ as an integral with continuous densities. Finally, define

$$\mathcal{D} := \{F \in \mathcal{F} \mid x \mapsto P^x(F) \text{ is } \mathcal{B}(\mathbb{R}^d) \text{ measurable}\}.$$

- $\Omega \in \mathcal{D}$, because $x \mapsto P^x(\Omega) = 1$ is a constant and thus measurable
- Let $F \in \mathcal{D}$ then $x \mapsto P^x(F^C) = 1 - P^x(F)$, is just a translation and thus measurable.
- Let $(F_n)_{n\in\mathbb{N}} \subset \mathcal{D}$ then $x \mapsto P^x(\bigcup_{n\in\mathbb{N}} F_n) = \sum_{n=1}^{\infty} P^x(F_n)$ is again measurable.

Thus,

$$\mathcal{F} = \sigma(\mathcal{C}) \subseteq \mathcal{D} \subseteq \mathcal{F} \quad \Longrightarrow \quad \mathcal{D} = \mathcal{F}.$$

□

Definition 1.35

A d-dimensional Brownian family is an adapted, d-dimensional process $(W_t)_{t\geq 0}$ on a filtered measurable space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0})$ and a family of probability measures $(P^x)_{x\in\mathbb{R}^d}$ such that

- $\forall F \in \mathcal{F}$ the mapping $x \mapsto P^x(F)$ is (universally ???) measurable
- $\forall x \in \mathbb{R}^d$: $P^x(W_0 = x) = 1$
- $\forall P^x$ the process W is a d-dimensional Brownian motion starting in x.

1.4 Weak Solutions by Means of the Girsanov Theorem

Proposition 1.36

For $T > 0$ fixed consider

$$dX_t = b(t, X_t)dt + dW_t, \quad t \in [0, T],$$

where W is d-dimensional Brownian motion, $b : [0,T] \times \mathbb{R}^d \to \mathbb{R}^d$, is Borel measurable and satisfies

$$\forall x \in \mathbb{R}^d, t \in [0,T] : \quad \|b(x,t)\| \leq K(1 + \|x\|),$$

for some constant K. Then for any initial distribution μ on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ the SDE has a weak solution with initial distribution μ.

Proof. Beginning with a Brownian family $(W_t)_{t\geq 0}$, $(\mathcal{F}_t)_{t\geq 0}$ and $(\mathbb{P}^x)_{x\in\mathbb{R}^d}$ then

$$Z_t := e^{\sum_{j=1}^d \int_0^t b_j(s,X_s)dX_s^j - \frac{1}{2}\int_0^t \|b(s,X_s)\|^2 ds}$$

is a martingale under each $\mathbb{P}^x$. Under Q^x given by $\frac{dQ^x}{d\mathbb{P}^x} =: Z_T$

$$W_t := X_t - X_0 - \int_0^t b(s, X_s)ds, \quad t \in [0,T],$$

is a Brownian motion with $Q^x(W_0 = 0) = 1$, $\forall x \in \mathbb{R}^d$. The above is equivalent to

$$X_t = X_0 - \int_0^t b(s, X_s)ds + W_t, \quad t \in [0, T].$$

With $Q^\mu(\cdot) := \int_{\mathbb{R}^d} Q^x(\cdot)d\mu(x)$, then $((X, W), (\Omega, \mathcal{F}, Q^\mu), (\mathcal{F}_t))$ is a weak solution. □

Remark 1.4

To find a weak solution for $t \in [0, \infty)$, the finite-time Girsanov theorem needs to be extended. The Radon-Nikodym derivative $Z_T := \frac{dQ^x}{d\mathbb{P}^x}$ is only defined on the bounded σ-algebra $\mathcal{F}_T$.
For $M_t^j := \int_0^t b(s, X_s)dX_s^j$ yields that $\int_0^t \|b(s, X_s)\|^2 ds = \langle M \rangle_t$. Then

$$Z_\infty = \lim_{t\to\infty} e^{\sum_{j=1}^d M_t^j - \frac{1}{2}\langle M\rangle_t} \notin L^1(\mathbb{P}).$$

So $(Z_t)_{t\in[0,\infty)}$ is not a martingale. Furthermore, with Fatou

$$\mathbb{E}[\liminf_{t\to\infty} Z_t] \le \liminf_{t\to\infty} \mathbb{E}[Z_t] = 1.$$

This means we cannot find a Girsanov measure Q^x such that $Q^x(A) = \int_A Z_\infty d\mathbb{P}^x$ for any $A \in \mathcal{B}(\mathbb{R}^d)$, because

$$1 = Q^x(\Omega) \neq \int_\Omega Z_\infty d\mathbb{P}^x < 1.$$

So Z_∞ also cannot be Radon derivative.
Let Ω be infinite set and $\mathcal{A}$ an algebra. By Zorn there exists $\mathcal{U} \subset \mathcal{A}$ such that $\forall A, B \in \mathcal{U} : A \cap B \neq \emptyset$ but $\bigcap_{A\in\mathcal{U}} A = \emptyset$. Define

$$Q_p : \mathcal{A} \to \{0, 1\}, \quad A \mapsto Q_p(A) = \begin{cases} 1 & , A \in \mathcal{U} \\ 0 & , A \notin \mathcal{U} \end{cases}.$$

Because $\bigcap_{A\in\mathcal{U}} A = \emptyset$ there exist $A_1 \supset A_2 \supset \ldots$ in $\mathcal{U}$ such that $\bigcap_{n\in\mathbb{N}} A_n = \emptyset$, with $Q_p(A_n) = 1$ for every $n \in \mathbb{N}$. Thus, $\lim_{n\to\infty} Q_p(A_n) = 1 \neq 0$, which is a contradition. So Q_p cannot be extended to $\sigma(A)$.

1.5 Stochastic Calculus Backround

Remark 1.5

We discuss the notion of an extended probability space. Let $(X_t)_{t\ge0}$ be $\mathcal{F}_t$-adapted process on $(\Omega, \mathcal{F}, \mathbb{P})$. If one requires a d-dimensional Brownian motion independent of X, one might run into the issue that the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ might not be rich enough to support this Brownian motion.

Example 1.4

Consider $\Omega := C([0, \infty), \mathbb{R}^d)$ and $\mathcal{F} := \sigma(X_s \mid s \ge 0) = \sigma(X)$, then $X_t(\omega) = \omega(t)$ is a stochastic process on $(\Omega, \mathcal{F}, \mathbb{P})$. Any Brownian motion on that probability space satisfies $\sigma(W) \subseteq \mathcal{F} = \sigma(X)$. Doobs factorization lemma yields that for every $t \ge 0$ there exists a $\mathbb{P}_X$-a.s. unique measurable mapping $\varphi_t := C([0, \infty), \mathbb{R}^d) \to \mathbb{R}^d$ such that $W_t = \varphi_t(X)$. So W is just a transformation of X.

Remark 1.6

To extend the probability space to construct a Brownian motion independent of X, let $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$ be another probability space on which $\hat{W}$ is a Brownian motion and set $\tilde{\Omega} := \Omega \times \hat{\Omega}$, $\tilde{\mathcal{G}} := \mathcal{F} \otimes \hat{\mathcal{F}}$, $\tilde{\mathbb{P}} := \mathbb{P} \otimes \hat{\mathbb{P}}$ and the filtration $\tilde{\mathcal{G}}_t := \mathcal{F}_t \otimes \hat{\mathcal{F}}_t$. We make the filtration right continuous by defining

$$\tilde{\mathcal{F}}_t := \bigcap_{s>t} \sigma(\tilde{\mathcal{G}} \cup \mathcal{N}),$$

where $\mathcal{N}$ are the $\tilde{\mathbb{P}}$-nullsets in $\tilde{\mathcal{G}}$. We make the σ-algebra complete by $\tilde{\mathcal{F}} := \sigma(\tilde{\mathcal{G}} \cup \mathcal{N})$. Finally, we extend X and W to be $\tilde{\mathcal{F}}_t$-adapted processes on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ by defining

$$\forall (\omega, \hat{\omega}) \in \tilde{\Omega}: \quad \tilde{X}_t(\omega, \hat{\omega}) := X_t(\omega) \quad \text{and} \quad \tilde{W}_t(\omega, \hat{\omega}) := W_t(\hat{\omega}).$$

Then $(\tilde{W}_t)_{t\geq 0}$ is independent of $(\tilde{X}_t)_{t\geq 0}$.

Theorem 1.37

Let $(M_t := (M_t^1, ..., M_t^d))_{t\geq 0}$ defined on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$ with M^i being a continuous local martingale for every $i = 1, ..., d$. Also suppose that the cross-variation $\langle M^i, M^j \rangle_t$ is $\mathbb{P}$-a.s. an absolutely continuous function in t for every $i, j = 1, ..., d$. Then there exists an extension $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ of $(\Omega, \mathcal{F}, \mathbb{P})$ on which we define a Brownian motion W and $((X_t^{i,k})_i, k = 1, ..., d)_{t\geq 0}$ a metric of measurable adapted processes with

$$\forall t \in [0, \infty), i, k = 1, ..., d : \tilde{\mathbb{P}}(\int_0^t |X_s^{i,k}|^2 ds < \infty) = 1,$$

such that

$$M_t^i = \sum_{k=1}^d \int_0^t X_s^{i,k} dW_s^k \quad i = 1, ..., d, t \in [0, \infty) \quad \tilde{\mathbb{P}} - a.s.$$

and

$$\langle M^i, M^j \rangle_t = \sum_{k=1}^d \int_0^t X_s^{i,k} X_s^{j,k} ds, \quad i, j = 1, ..., d, t \in [0, \infty) \quad \tilde{\mathbb{P}} - a.s.$$

Proof. We reduce the d-dimensional process by a random, time dependent rotation of coordinates to one-dimensional processes. Define

$$z_t^{i,j} := \frac{d}{dt} \langle M^i, M^j \rangle_t = \lim_{n\to\infty} n(\langle M^i, M^j \rangle_t - \langle M^i, M^j \rangle_{(t-\frac{1}{n})^+}).$$

As $z_t^{i,j} = z_t^{j,i}$, because $\langle cM^i \rangle_t = c^2 \langle M^i \rangle_t$ yields

$$\langle M^i, M^j \rangle_t = \frac{1}{4}(\langle M^i + M^j \rangle_t - \langle M^i - M^j \rangle_t) = \langle M^j, M^i \rangle_t,$$

we get that the matrix process $Z := (Z_t := (z_t^{ij})_{i,j=1,...,d})_{t\geq 0}$ is symmetric and $(s, \omega) \mapsto Z_s(\omega)$ is $\mathcal{B}([0, t])???\otimes$

$\mathcal{F}_t$-measurable, i.e. progressively measurable. Additionally, we have

$$\begin{aligned}\forall \alpha \in \mathbb{R}^d : \ \alpha^T Z_t \alpha &= \lim_{n\to\infty} n(\sum_{i=1}^d \sum_{j=1}^d \alpha_i(\langle M^i, M^j\rangle_t - \langle M^i, M^j\rangle_{(t-\frac{1}{n})^+})\alpha_j) \\ &= \lim_{n\to\infty} n(\langle \sum_{i=1}^d \alpha_i M^i\rangle_t - \langle \sum_{i=1}^d \alpha_i M^j\rangle_{(t-\frac{1}{n})^+}) \\ &= \frac{d}{dt}\langle \sum_{i=1}^d \alpha_i M^i\rangle_t \geq 0.\end{aligned}$$

Because the quadratic variation is non-decreasing, for fixed $\omega \in \Omega$ it is $\lambda - a.e.$ differentiable in $t \geq 0$ with Lebesgues theorem.
Because Z is progressively measurable the set

$$E := \{(t,\omega) \in [0,\infty) \times \Omega \mid Z_t(\omega) \text{ is not positive semi definite}\} \in \mathcal{B}([0,\infty)) \otimes \mathcal{F}.$$

This yields

$$(\lambda \otimes \mathbb{P})(E) = \int_\Omega \lambda(\{t \geq 0 \mid (t,\omega) \in E\}) d\mathbb{P}(\omega) = \int_\Omega 0 d\mathbb{P}(\omega) = 0.$$

So Z_t is positive semi definite for λ-a.e $t \geq 0$ $\mathbb{P}$-a.s.
It is well known that any symmetric, positive definite matrix can be diagonalized by an orthogonal matrix, i.e. there exists Q such that

$$Q^{-1} Z Q = \Lambda,$$

where Λ is the daigonal matrix with the non negative eigenvalues of Z. So there exists progressively measurable matrix valued processes

$$\left(Q_t := (q_t^{i,j})_{i,j=1,\dots,d}\right)_{t\geq 0} \quad \text{and} \quad \left(\Lambda_t := (\delta_{ij}\lambda_t^i)_{i,j=1,\dots,d}\right)_{t\geq 0},$$

such that for λ-a.e. $t \geq 0$ it holds

$$\sum_{k=1}^d q_t^{k,i} q_t^{k,j} = (q_t^{\cdot,i})^T q_t^{\cdot,j} = (q_t^{i,\cdot})^T q_t^{j,\cdot} = \sum_{k=1}^d q_t^{i,k} q_t^{j,k} = \delta_{ij}, \ i,j = 1,\dots,d \quad \mathbb{P} - a.s.$$

and

$$(q^{\cdot,i})^T Z_t (q^{\cdot,j}) = \sum_{k=1}^d \sum_{l=1}^d q_t^{k,i} z_t^{k,l} q_t^{l,j} = \delta_{ij}\lambda_t^i \geq 0, i,j = 1,\dots,d \quad \mathbb{P} - a.s.$$

From the orthogonality, we see that $(q_t^{k,i})^2 \leq 1$ and thus

$$\int_0^t (q_s^{k,i})^2 d\langle M^k\rangle_s \leq \langle M^k\rangle_t < \infty.$$

Then, $N_t^i := \sum_{k=1}^d \int_0^t q_s^{k,i} dM_s^k, \quad i = 1,\dots,d, t \in [0,\infty)$ is a continuous local martingale.
To finish the proof our goal is to represent M^i in terms of an independent Brownian motion on an extended probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$. The next step is to represent N^i in terms of that independent Brownian motion. First, see that

$$\begin{aligned}\langle N^i, N^j\rangle_t &= \sum_{k=1}^d \sum_{l=1}^d \int_0^t q_s^{k,i} q_s^{l,j} d\langle M^k, M^l\rangle_s \\ &= \sum_{k=1}^d \sum_{l=1}^d \int_0^t q_s^{k,i} q_s^{l,j} \frac{d}{dt}\langle M^k, M^l\rangle_s ds \\ &= \sum_{k=1}^d \sum_{l=1}^d \int_0^t q_s^{k,i} z_s^{k,l} q_s^{l,j} ds = \delta_{ij} \int_0^t \lambda_s^i ds\end{aligned}$$

Because N^i is a continuous local martingale $\langle N^i\rangle_t < \infty$, which implies that the Lebesgues-Stieltjes integral $\int_0^t \mathbf{1}_{\lambda_s^i>0}\frac{1}{\lambda_s^i}d\langle N^i\rangle_s$ exists. As

$$\langle N^i\rangle_t = \langle N^i, N^i\rangle_t = \int_0^t \lambda_s^i ds,$$

it follows $\frac{d}{dt}\langle N^i\rangle_t = \lambda_s^i$, which yields

$$\int_0^t \mathbf{1}_{\lambda_s^i>0}\frac{1}{\lambda_s^i}d\langle N^i\rangle_s = \int_0^t \mathbf{1}_{\lambda_s^i>0}ds \leq t.$$

This gives us that

$$W_t^i := \int_0^t \mathbf{1}_{\lambda_s^i>0}\frac{1}{\sqrt{\lambda_s^i}}dN_s^i + \int_0^t \mathbf{1}_{\lambda_s^i=0}d\tilde{W}_s^i, \quad i = 1, ..., d,$$

is a continuous local martingale, where $\tilde{W}$ is a Brownian motion on the extended probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$. Because

$$\forall t \in [0,\infty), i, j = 1, ..., d: \quad \langle W^i, W^j\rangle_t = \delta_{ij}t,$$

we have that W is a Brownian motion wrt. $(\tilde{\mathcal{F}}_t)_{t\geq 0}$. Moreover,

$$\int_0^t \sqrt{\lambda_s^i}dW_s^i = \int_0^t \mathbf{1}_{\lambda_s^i>0}dN_s^i + \int_0^t \sqrt{\lambda_s^i}\mathbf{1}_{\lambda_s^i=0}d\tilde{W}_s^i = \int_0^t \mathbf{1}_{\lambda_s^i>0}dN_s^i \overset{(*)}{=} \int_0^t \mathbf{1}_{\lambda_s^i\geq 0}dN_s^i = N_t^i, \quad \tilde{\mathbb{P}} - a.s.$$

although we used in (*) that the martingale $\int_0^t \mathbf{1}_{\lambda_s^i=0}dN_s^i$ has quadratic variation

$$\langle \int_0^\cdot \mathbf{1}_{\lambda_s^i=0}dN_s^i\rangle_t = \int_0^t \mathbf{1}_{\lambda_s^i=0}d\langle N^i\rangle_s = \int_0^t \mathbf{1}_{\lambda_s^i=0}\lambda_s^i ds = 0$$

and starting in zero yields that it is the trivial martingale, i.e.

$$\int_0^t \mathbf{1}_{\lambda_s^i=0}dN_s^i = 0 \quad , t \geq 0 \quad \tilde{\mathbb{P}} - a.s.$$

Setting $X_t^{i,k} := q_t^{i,k}\sqrt{\lambda_t^k}$ we get that

$$\begin{aligned}\forall t \geq 0, i, k = 1, ..., d: \quad \tilde{P}(\int_0^t (X_s^{i,k})^2 ds < \infty) &= \tilde{P}(\int_0^t (q_s^{i,k})^2\lambda_s^k ds < \infty)\\ &= \tilde{P}(\int_0^t \lambda_s^k ds < \infty) = \tilde{P}(\langle N^k\rangle_t < \infty) = 1.\end{aligned}$$

Finally,

$$\begin{aligned}\sum_{k=1}^d \int_0^t X^{i,k}dW_s^k &= \sum_{k=1}^d \int_0^t q_s^{i,k}\sqrt{\lambda_s^k}dW_s^k = \sum_{k=1}^d \int_0^t q_s^{i,k}dN_s^k\\ &= \sum_{j=1}^d\sum_{k=1}^d \int_0^t q_s^{i,k}q_s^{j,k}dM_s^j \overset{orthogonal}{=} \sum_{j=1}^d \delta_{ij}\int_0^t dM_s^j = M_t^i.\end{aligned}$$

The last equation follows directly from this one. □

Remark 1.7

Note that if the matric process $Z_t := (z_t^{i,j})_{i,j=1,\dots,d}$ has constant rank r $\mathbb{P}$-a.s. λ-almoust every $t \geq 0$, then

$$\forall i \leq r: \quad W_t^i = \int_0^t \mathbf{1}_{\lambda_s^i > 0} \frac{1}{\sqrt{\lambda_s^i}} dN_s^i,$$

is a Brownian motion on $(\Omega, \mathcal{F}, \mathbb{P})$ and there is no need to extend the probability space, as it is independently defined from $\tilde{W}$.

SECTION 2

GIRSANOV THEOREM

We study local martingales if a different equivalent probability measure Q on $(\Omega, \mathcal{F})$ is chosen.

Definition 2.1

We call two probability measures Q and $\mathbb{P}$ on $(\Omega, \mathcal{F})$ equivalent if

$$\forall A \in \mathcal{F}: \quad \mathbb{P}(A) = 0 \quad \Longleftrightarrow \quad Q(A) = 0.$$

By Radon-Nikodym there exists a Radon density L_T on $(\Omega, \mathcal{F})$ with $L_T := \frac{dQ}{d\mathbb{P}}$ such that

$$\forall A \in \mathcal{F}: \quad \mathbb{E}^Q[\mathbf{1}_A] = \mathbb{E}[\mathbf{1}_A L_T].$$

If $L_T \in L^1(\Omega, \mathcal{F}, \mathbb{P})$ define $L_t := \mathbb{E}[L_T \mid \mathcal{F}_t]$ for every $t \in [0, T]$, then $(L_t)_{t\in[0,T]}$ is a martingale (???) and is called density process of Q wrt. $\mathbb{P}$.

Assumption 2.2

The stochastic process $(L_t)_{t\in[0,T]}$ satisfies:

- $(L_t)_{t\in[0,T]}$ is a continuous $\mathbb{P}$-martingale
- $\forall t \in [0, T]$: $L_t > 0$ $\mathbb{P}$-a.s. and $L_0 = 1$.

Under these assumptions we can define the associated probability measure

$$Q(A) := \mathbb{E}^Q[\mathbf{1}_A] = \mathbb{E}[\mathbf{1}_A L_T], \quad \forall A \in \mathcal{F}.$$

Then Q is equivalent to $\mathbb{P}$ and with Radon-Nikodym the density is $\frac{dQ}{d\mathbb{P}} = L_T$.

Proposition 2.3

Let $(L_t)_{t\in[0,T]}$ satisfy the assumptions above, where Q is the associated probability measure and $(X_t)_{t\in[0,T]}$ an adapted, continuous stochastic process, then

$$(L_t X_t)_{t\in[0,T]} \text{ is a (local) } \mathbb{P}\text{-martingale} \quad \Longleftrightarrow \quad (X_t)_{t\in[0,T]} \text{ is a (local) } Q\text{-martingale.}$$

Proof. "$\Rightarrow$"
Because X is adapted and Q integrable, it holds

$$\mathbb{E}^Q[|X_t|] = \mathbb{E}[L_T|X_t|] = \mathbb{E}[\mathbb{E}[L_T|X_t| \mid \mathcal{F}_t]] = \mathbb{E}[L_t|X_t|] = \mathbb{E}[|L_tX_t|] < \infty,$$

because L and LX are $\mathbb{P}$-martingales. In order to show the martingale property it suffices to show:

$$\forall A \in \mathcal{F}_s : \quad \mathbb{E}^Q[\mathbf{1}_A(X_t - X_s)] = 0.$$

This immediately follows from

$$\begin{aligned}\mathbb{E}^Q[\mathbf{1}_A(X_t - X_s)] &= \mathbb{E}[L_t\mathbf{1}_A(X_t - X_s)] \\ &= \mathbb{E}[\mathbb{E}[L_T\mathbf{1}_AX_t \mid \mathcal{F}_t] - \mathbb{E}[L_T\mathbf{1}_AX_s \mid \mathcal{F}_s]] \\ &= \mathbb{E}[L_t\mathbf{1}_AX_t - L_s\mathbf{1}_AX_s] = \mathbb{E}[\mathbf{1}_A(L_tX_t - L_sX_s)] \overset{martingale}{=} 0.\end{aligned}$$

"$\Leftarrow$"
Same arguments. □

Theorem 2.4

If

$$L_t := e^{\int_0^t X_s dB_s - \frac{1}{2}\int_0^t X_s^2 ds}, \quad t \in [0,T]$$

and $(L_t)_{t\in[0,T]}$ is martingale on $(\Omega, \mathcal{F}_T, \mathbb{P})$ wrt. the Brownian standard filtration $(\mathcal{F}_t)_{t\in[0,T]}$, then

$$\tilde{B}_t := B_t - \int_0^t X_s ds, \quad t \in [0,T],$$

is Brownian motion wrt. $(\Omega, \mathcal{F}_T, (\mathcal{F}_t)_{t\in[0,T]}, Q)$ where $\frac{dQ}{d\mathbb{P}} = L_T$.

Proof. We need to show that $\tilde{B}$ is continuous local Q-martingale with $\langle \tilde{B} \rangle_t = t$, $t \in [0,T]$.

- To show: $\tilde{B}$ is continuous Q-martingale.
  With proposition before we need to show $(\tilde{B}_tL_t)_{t\in[0,T]}$ is continuous local $\mathbb{P}$-martingale. We can apply Ito Formula for L_t, because we can write it as

  $$L_t = f(U_t, U_t) := e^{U_t + Y_t}, \quad U_t := \int_0^t X_s dB_s, \quad V_t := -\frac{1}{2}\int_0^t X_s^2 ds.$$

  Because integrating over $\langle U \rangle_\cdot$ and $\langle V \rangle_\cdot$ yields a Lebesgues Stieltjes integral which leads to $X_s^2 ds$ and 0 respectively. $d\langle U, V \rangle_s = X_s^2 \cdot 0 ds = 0$. $f(0,0) = 1$ and Itos formula yields

  $$\begin{aligned}L_t &= 1 + \int_0^t L_s dU_s + \int_0^t L_s dV_s + \frac{1}{2}\int_0^t L_s X_s^2 ds \\ &= 1 + \int_0^t L_s X_s dB_s - \frac{1}{2}\int_0^t L_s X_s^2 ds + \frac{1}{2}\int_0^t L_s X_s^2 ds \\ &= 1 + \int_0^t L_s X_s dB_s\end{aligned}$$

Thus, $dL_t = L_s X_s dB_s$, which yields with the product rule

$$\begin{aligned}\tilde{B}_t L_t &= \tilde{B}_0 L_0 + \int_0^t \tilde{B}_s dL_s + \int_0^t L_s d\tilde{B}_s + \langle B, L\rangle_t \\ &= 0 + \int_0^t \tilde{B}_s L_s X_s dB_s + \int_0^t L_s dB_s - \underbrace{\int_0^t L_s X_s ds + \int_0^t 1 \cdot L_s X_s ds}_{=0} \\ &= \int_0^t (\tilde{B}_s X_s + 1) L_s dB_s,\end{aligned}$$

which is a continuous local martingale wrt. $\mathbb{P}$.

- $(\tilde{B}_t)_{t\in[0,T]}$ is a continuous local Q-martingale.
  Let $((\tilde{B}L)_{t\wedge\tau_n})_{t\in[0,T]}$ be a $\mathbb{P}$-martingale for some suitable localizing stopping times $\tau_n \nearrow T$. With the first step and the proposition from before $(\tilde{B}_{t\wedge\tau_n})_{t\in[0,T]}$ is a continuous Q-martingale wrt. $(\mathcal{F}_t)_{t\in[0,T]}$ it follows that $(\tilde{B}_t)_{t\in[0,T]}$ is a continuous local Q-martingale.

- $\forall t \in [0,T]$: $\langle \tilde{B}\rangle_t = t$ under Q.
  To this end, we need to show that $(\tilde{B}_t - t)_{t\in[0,T]}$ is a continuous local martingale under Q. With the proposition from before it suffices to show that $((\tilde{B}_t^2 - t)L_t)_{t\in[0,T]}$ is a continuous local $\mathbb{P}$-martingale. Then, because the quadratic variation of any finite variation process, e.g. $\int_0^\cdot X_s ds$, is zero and and its co-variation with a continuous martingale is zero, under $\mathbb{P}$ we have

  $$\langle \tilde{B}\rangle_t = \langle B - \int_0^\cdot X_s ds\rangle_t = \langle B\rangle_t - 2\langle B, \int_0^\cdot X ds\rangle_t + \langle \int_0^\cdot X ds\rangle_t = \langle B\rangle_t - 0 + 0 = t.$$

  Finally, with Itos formula for Ito processes for $f(x) = x^2$ we have

  $$\tilde{B}_t^2 = \int_0^t 2\tilde{B}_s d\tilde{B}_s + \int_0^t \langle \tilde{B}\rangle_s = \int_0^t 2\tilde{B}_s d\tilde{B}_s + t \overset{Def.}{=} \int_0^t 2\tilde{B}_s dB_s - \int_0^t 2\tilde{B}_s X_s ds + t.$$

  This yields with the product formula:

  $$\begin{aligned}(\tilde{B}_t^2 - t)L_t &= \int_0^t (B_s^2 - s) dL_s + \int_0^t L_s d(\tilde{B}_s - s) + \langle \tilde{B}^2 - \cdot, L\rangle_t \\ &= \int_0^t (B_s^2 - s) dL_s + \int_0^t L_s 2\tilde{B}_s d\tilde{B}_s + \int_0^t 2\tilde{B}_s L_s X_s ds \\ &= \int_0^t (B_s^2 - s) L_s X_s dB_s + \int_0^t 2L_s \tilde{B}_s dB_s - \int_0^t 2L_s \tilde{B}_s X_s ds + \int_0^t 2\tilde{B}_s L_s X_s ds \\ &= \int_0^t ((\tilde{B}_s - s)X_s + 2\tilde{B}_s) L_s dB_s,\end{aligned}$$

  which is a continuous local $\mathbb{P}$-martingale.

□

SECTION 3

WEAK SOLUTIONS

The Brownian motion is now denoted by (W_t).
One can prove the existence of a solution in a weaker sense for less restrictive assumptions on the drift term

Definition 3.1

Let $a : [0,\infty) \times \mathbb{R}^d \to \mathbb{R}^d$ and $b : [0,\infty) \times \mathbb{R}^d \to \mathbb{R}^{d\times r}$ be Borel measurable functions. For a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$, a r-dimensional $(\mathcal{F}_t)$-Brownian motion W and a $\mathcal{F}_0$ measurable random variable ξ, we call call $(X_t)_{t\geq 0}$ strong solution to the SDE

$$dX_t = a(t, X_t)dt + b(t, X_t)dW_t,$$

if

- $(X_t)_{t\geq 0}$ is $(\mathcal{F}_t^W \vee \sigma(\xi) \vee \mathcal{N})_{t\geq 0}$ adapted to the augmented filtration
- $$\forall i = 1, ..., d, j = 1, ..., r \text{ and } t \in [0,\infty) : \quad \mathbb{P}(\int_0^t |a_i(s, X_s)| + b_{ij}^2(s, X_s)ds < \infty) = 1$$
- $$\forall t \in [0, T], i = 1, ..., d : \quad X_t^i = X_0^i + \int_0^t a_i(s, X_s)ds + \sum_{j=1}^r \int_0^t b_{ij}(s, X_s)dW_s^j.$$

Definition 3.2

A weak solution, with initial distribution μ on $\mathcal{B}(\mathbb{R}^d)$ of

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dW_t$$

is a tuple $((\Omega, \mathcal{F}, \mathbb{P}), (\mathcal{F}_t)_{t\geq 0}, X, W)$ where $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$ is a filtered probability space, where the filtration satisfies the usual conditions, X is $(\mathcal{F}_t)$-adapted, $\mathbb{R}^d$ valued process and W a r-dimensional $(\mathcal{F}_t)$-Brownian motion W

- $$\forall i = 1, ..., d, j = 1, ..., r \text{ and } t \in [0,\infty) : \quad \mathbb{P}(\int_0^t |a_i(s, X_s)| + b_{ij}^2(s, X_s)ds < \infty) = 1$$
- $$\forall t \in [0, T], i = 1, ..., d : \quad X_t^i = X_0^i + \int_0^t a_i(s, X_s)ds + \sum_{j=1}^r \int_0^t b_{ij}(s, X_s)dW_s^j.$$

Remark 3.1

For a strong solution the probability space was given. For weak solutions, the probability space is provided as part of the solution. Further, the randomness in a strong solution only comes from the randomness of the augmented standard Brownian filtration, whereas the randomness in a weak solution comes from any filtration, with the restriction that W has to be a $(\mathcal{F}_t)$-Brownian motion. This means, that the solution at time t can not only be given by the initial condition, a measurable functional given by the Brownian path and the drift, but also from randomness independent from the initial condition and the Brownian motion.

Example 3.1

Consider

$$dX_t = b(t, X_t)dt + dW.t, \quad t \in [0, T], \quad T < \infty$$

and $b : \mathbb{R}^d \to \mathbb{R}^d$ is a measurable function with sub-linear growth

$$\forall x \in \mathbb{R}^d, t \in [0, T] : \quad \|b(t, x)\| \leq K(1 + \|x\|).$$

Let X be a Brownian motion on $(\Omega, \mathcal{F}, \mathbb{P})$ and let $(\mathcal{F}_t)_{t\geq 0}$ be the augmented filtration generated by X, i.e.

$$\forall t \geq 0 : \quad \mathcal{F}_t := \sigma(\mathcal{F}_t^X \cup \mathcal{N}), \quad \mathcal{N} := \{A \subseteq \Omega \mid \exists C \in \mathcal{F}, \mathbb{P}(C) = 0 : \quad A \subseteq C\}.$$

Define $\forall t \in [0, T]$:

$$Z_t := e^{\int_0^t b(s,X_s)dX_s - \frac{1}{2}\int_0^t \|b(s,X_s)\|^2 ds},$$

which is a martingale under $\mathbb{P}$. Define Q via the Radon-Nikodym derivative

$$\frac{dQ}{d\mathbb{P}} = Z_T, \quad Q(\cdot) := \mathbb{E}^Q[\mathbf{1}.].$$

With Girsanov, the process

$$\tilde{W}_t := X_t - X_0 - \int_0^t b(s, X_s)ds, \quad t \in [0, T]$$

is Brownian motion with $Q(\tilde{W}_0 = 0) = 1$. Then $(X, \tilde{W}_t)$ and $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, Q)$ is weak solution.

3.1 Weak Solutions by Means of the Girsanov Theorem

Proposition 3.3

For $T > 0$ fixed consider

$$dX_t = b(t, X_t)dt + dW_t, \quad t \in [0, T],$$

where W is d-dimensional Brownian motion, $b : [0, T] \times \mathbb{R}^d \to \mathbb{R}^d$, is Borel measurable and satisfies

$$\forall x \in \mathbb{R}^d, t \in [0, T] : \quad \|b(x, t)\| \leq K(1 + \|x\|),$$

for some constant K. Then for any initial distribution μ on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ the SDE has a weak solution with initial distribution μ.

Proof. Beginning with a Brownian family $(W_t)_{t\geq 0}$, $(\mathcal{F}_t)_{t\geq 0}$ and $(\mathbb{P}^x)_{x\in\mathbb{R}^d}$ then

$$Z_t := e^{\sum_{j=1}^d \int_0^t b_j(s,X_s)dX_s^j - \frac{1}{2}\int_0^t \|b(s,X_s)\|^2 ds}$$

is a martingale under each $\mathbb{P}^x$. Under Q^x given by $\frac{dQ^x}{d\mathbb{P}^x} =: Z_T$

$$W_t := X_t - X_0 - \int_0^t b(s, X_s)ds, \quad t \in [0, T],$$

is a Brownian motion with $Q^x(W_0 = 0) = 1$, $\forall x \in \mathbb{R}^d$. The above is equivalent to

$$X_t = X_0 - \int_0^t b(s, X_s)ds + W_t, \quad t \in [0, T].$$

With $Q^\mu(\cdot) := \int_{\mathbb{R}^d} Q^x(\cdot) d\mu(x)$, then $((X,W),(\Omega,\mathcal{F},Q^\mu),(\mathcal{F}_t))$ is a weak solution. □

Remark 3.2

To find a weak solution for $t \in [0,\infty)$, the finite-time Girsanov theorem needs to be extended. The Radon-Nikodym derivative $Z_T := \frac{dQ^x}{d\mathbb{P}^x}$ is only defined on the bounded σ-algebra $\mathcal{F}_T$.
For $M_t^j := \int_0^t b(s,X_s) dX_s^j$ yields that $\int_0^t \|b(s,X_s)\|^2 ds = \langle M \rangle_t$. Then

$$Z_\infty = \lim_{t\to\infty} e^{\sum_{j=1}^d M_t^j - \frac{1}{2}\langle M\rangle_t} \notin L^1(\mathbb{P}).$$

So $(Z_t)_{t\in[0,\infty)}$ is not a martingale. Furthermore, with Fatou

$$\mathbb{E}[\liminf_{t\to\infty} Z_t] \le \liminf_{t\to\infty} \mathbb{E}[Z_t] = 1.$$

This means we cannot find a Girsanov measure Q^x such that $Q^x(A) = \int_A Z_\infty d\mathbb{P}^x$ for any $A \in \mathcal{B}(\mathbb{R}^d)$, because

$$1 = Q^x(\Omega) \neq \int_\Omega Z_\infty d\mathbb{P}^x < 1.$$

So Z_∞ also cannot be Radon derivative.
Let Ω be infinite set and $\mathcal{A}$ an algebra. By Zorn there exists $\mathcal{U} \subset \mathcal{A}$ such that $\forall A,B \in \mathcal{U} : A \cap B \neq \emptyset$ but $\bigcap_{A\in\mathcal{U}} A = \emptyset$. Define

$$Q_p : \mathcal{A} \to \{0,1\}, \quad A \mapsto Q_p(A) = \begin{cases} 1 & , A \in \mathcal{U} \\ 0 & , A \notin \mathcal{U} \end{cases}.$$

Because $\bigcap_{A\in\mathcal{U}} A = \emptyset$ there exist $A_1 \supset A_2 \supset \ldots$ in $\mathcal{U}$ such that $\bigcap_{n\in\mathbb{N}} A_n = \emptyset$, with $Q_p(A_n) = 1$ for every $n \in \mathbb{N}$. Thus, $\lim_{n\to\infty} Q_p(A_n) = 1 \neq 0$, which is a contradition. So Q_p cannot be extended to $\sigma(A)$.

SECTION 4

THE MARTINGALE PROBLEM OF STROOCK AND VARADHAN

Weak solutions is finding a probability space, a Brownian motion and a process with the right law (determined by its finite- dimensional distributions).

Definition 4.1

We say a weak solution is unique in sense of probability law if for any two weak solutions $((X,W),(\Omega,\mathcal{F},\mathbb{P}),(\mathcal{F}_t)_{t\ge0})$ and $((\tilde{X},\tilde{W}),(\tilde{\Omega},\tilde{\mathcal{F}},\tilde{\mathbb{P}}),(\tilde{\mathcal{F}}_t)_{t\ge0})$ with

$$\mathbb{P}(X_0 \in A) = \tilde{\mathbb{P}}(\tilde{X}_0 \in A), \quad A \in \mathcal{B}(\mathbb{R}^d),$$

holds that X and $\tilde{X}$ have the same law, i.e.

$$\mathbb{P} \circ X^{-1}(A) = \tilde{\mathbb{P}} \circ \tilde{X}^{-1}(A), \quad \forall A \in \mathcal{B}(C([0,\infty),\mathbb{R}^d)).$$

There is also a different approach: Search for the law of a diffions process with drift and dispersion coefficients in terms of a martingale problem, which will be equivalent to solving the related SDE in a weak sense. This formulation is useful in terms of continuity and weak convergence arumgents that yield exitnece results and "invariance principles" (i.e. the convergence of Markov chains to diffion processes).

4.1 Fundamentals and Motiviation:

Assume $((X,W),(\Omega,\mathcal{F},\mathbb{P}),(\mathcal{F}_t)_{t\geq 0})$ is a weak solution to the SDE.

Definition 4.2

We call for every u in some function space

$$L : u \mapsto F(x, D^{k_1}, ..., D^{k_s}), \quad \text{where} \quad F : E \times \mathbb{R}^s \to \mathbb{R},$$

although $D^{k_i} := (\partial^{k_1} \cdots \partial^{k_i})$, differential operator on the domain Ω of order s.

For every $t \geq 0$ we introduce the second order differential operator for any $f \in C^2(\mathbb{R}^d)$:

$$(A_t f)(\cdot) := \frac{1}{2}\sum_{i=1}^{d}\sum_{k=1}^{d} a_{ik}(t,\cdot)\frac{\partial^2}{\partial x_i \partial x_k} f(\cdot) + \sum_{i=1}^{d} b_i(t,\cdot)\frac{\partial}{\partial x_i} f(\cdot), \quad \forall f \in C^2(\mathbb{R}^d).$$

Proposition 4.3

Let $(X_t)_{t\geq 0}$ a continuous Ito diffusion on $(\Omega,\mathcal{F},(\mathcal{F}_t)_{t\geq 0},\mathbb{P})$ with generator A_t and diffusion matric $a(t,x) = \sigma(t,x)^T\sigma(t,x)$. For every $f \in C^{1,2}([0,\infty)\times\mathbb{R}^d,\mathbb{R})$ the process $(M_t^f)_{t\geq 0}$ defined by

$$M_t^f := f(t,X_t) - f(0,X_0) - \int_0^t (\frac{\partial}{\partial s} + A_s) f(s,X_s) ds,$$

a continuous local martingale. If g is another Member of $C^{1,2}([0,\infty)\times\mathbb{R}^d,\mathbb{R})$ then M^g is a continuous local martingale and

$$\langle M^f, M^g\rangle_t = \sum_{i=1}^{d}\sum_{k=1}^{d}\int_0^t a_{ik}(s,X_s)\frac{\partial}{\partial x_i} f(s,X_s)\frac{\partial}{\partial x_k} g(s,X_s) ds.$$

Finally, if $f \in C_0([0,\infty)\times\mathbb{R}^d,\mathbb{R})$ (chat sagt $C_c^{1,2}$???) and σ_{ij} are bounded on $suppf$ then M_2^f is a continuous square integrable martingale.

Proof. Using Itos formula for the Ito diffusion

$$dX_t^i = b_i(t,X_t)dt + \sum_{i=1}^{r}\sigma_{ij}(t,X_t)dW_t^j$$

and reminding that

$$d\langle X^i, X^k\rangle_t \overset{???}{=} \sum_{j=1}^{r}\sigma_{ij}(t,X_t)\sigma_{kj}(t,X_t)dt =: a_{ik}(t,X_t)dt$$

yields that

$$\begin{aligned}
df(t,X_t) &= \frac{\partial}{\partial t}f(t,X_t)dt + \sum_{i=1}^{d}\frac{\partial}{\partial x_i}f(t,X_t)\left(b_i(t,X_t)dt + \sum_{j=1}^{r}\sigma_{ij}(t,X_t)dW_t^j\right)\\
&\quad + \frac{1}{2}\sum_{i,k=1}^{d}\frac{\partial^2}{\partial x_i\partial x_k}f(t,X_t)a_{ik}(t,X_t)dt\\
&= \left(\frac{\partial}{\partial t}f(t,X_t) + \sum_{i=1}^{d}b_i(t,X_t)\frac{\partial}{\partial x_i}f(t,X_t) + \frac{1}{2}\sum_{i,k=1}^{d}a_{ik}(t,X_t)\frac{\partial^2}{\partial x_i\partial x_k}f(t,X_t)\right)dt\\
&\quad + \sum_{i=1}^{d}\frac{\partial}{\partial x_i}f(t,X_t)\sum_{j=1}^{r}\sigma_{ij}(t,X_t)dW_t^j\\
&= (\frac{\partial}{\partial t} + \mathcal{A}_t)f(t,X_t)dt + \sum_{i=1}^{d}\sum_{j=1}^{r}\frac{\partial}{\partial x_i}f(t,X_t)\sigma_{ij}(t,X_t)dW_t^j\\
&\iff f(t,X_t) - f(0,X_0) - \int_0^t(\frac{\partial}{\partial s} + \mathcal{A}_s)f(s,X_s)ds = \sum_{i=1}^{d}\sum_{j=1}^{r}\int_0^t\sigma_{ij}(s,X_s)\frac{\partial}{\partial x_i}f(s,X_s)dW_s^j.
\end{aligned}$$

Define the stopping time

$$T_n := \inf\{t\geq 0 \mid \|X\|_t \geq n \vee \exists(i,j): \quad \int_0^t \sigma_{ij}^2(s,X_s)ds \geq n\}.$$

By definition of the weak solution we have

$$\mathbb{P}(\int_0^t \sigma_{ij}^2(s,X_s)ds < \infty) \geq \mathbb{P}(\int_0^t |a_i(s,X_s)| + \sigma_{ij}^2(s,X_s)ds < \infty) = 1,$$

yields $\lim_{n\to\infty} T_n = \infty$ $\mathbb{P}-a.s.$ Then the process defined through $M_t^{f,n} := M_{t\wedge T_n}^f$ is a continuous martingale, because a stochastic intefral $I_t = \int_0^t H_s dW_s$ is a continuous martignale if $\mathbb{E}[\int_0^t H_s^2 ds] < \infty$. This is the case! So M_t^f is a continuous local martingale. The cross variation is

$$\begin{aligned}
\langle M^f, M^g\rangle_t &= \sum_{i=1}^{d}\sum_{j=1}^{r}\langle\int_0^{\cdot\wedge S_n}\sigma_{ij}(s,X_s)\frac{\partial}{\partial x_i}f(s,X_s)dW_s^j, \int_0^{\cdot\wedge S_n}\sigma_{ij}(s,X_s)\frac{\partial}{\partial x_i}g(s,X_s)dW_s^j\rangle_t\\
&\overset{???}{=} \sum_{i=1}^{d}\sum_{k=1}^{d}\int_0^t\sigma_{ij}^2(s,X_s)\frac{\partial}{\partial x_i}f(s,X_s)\frac{\partial}{\partial x_k}g(s,X_s)ds.
\end{aligned}$$

Finally, if f has compact support on which σ_{ij} is bounded, then the integrand $\sigma_{ij}(s,X_s)\frac{\partial}{\partial x_i}f(s,X_s)$ is bounded. So M^f is a continuous and square integrable martingale. □

Remark 4.1

Problem:
Suppose that $((X,W),(\Omega,\mathcal{F},\mathbb{P}),(\mathcal{F}_t)_{t\geq 0})$ is a weak solution to the SDE on $C([0,\infty)^d)$. Set

$$(\mathcal{A}_t'u)(y) := \frac{1}{2}\sum_{i=1}^{d}\sum_{k=1}^{d}a_{ik}(t,y)\frac{\partial^2}{\partial x_i\partial x_k}u(y(t)) + \sum_{i=1}^{d}b_i(t,y)\frac{\partial}{\partial x_i}u(y(t)), t\in[0,\infty), y\in C([0,\infty)^d), u\in C^2(\mathbb{R}^d).$$

For any $f, g \in C^{1,2}([0,\infty) \times \mathbb{R}^d; \mathbb{R})$ the process

$$M_t^f := f(t, X_t) - f(0, X_0) - \int_0^t (\frac{\partial}{\partial s} + \mathcal{A}'_s) f(s, X_s) ds$$

is a continuous local martingale and

$$\langle M^f, M^g \rangle_t = \sum_{i=1}^d \sum_{k=1}^d \int_0^t a_{ik}(s, X_s) \frac{\partial}{\partial x_i} f(s, X_s) \frac{\partial}{\partial x_k} g(s, X_s) ds.$$

Finally, if the first derivative of f is bounded, then for every $0 \leq T < \infty$:

$$\forall y \in C([0,\infty)^d), t \in [0,T]: \quad \|\sigma(t,y)\| \overset{???}{\leq} K_T.$$

So M^f is continuous square integrable martingale.

Remark 4.2

If we set $b_i \equiv 0$ and $\sigma_{ij} \equiv \delta_{ij}$ for every $i, j = 1, ..., d$ then

$$\mathcal{A} f = \frac{1}{2} f = \frac{1}{2} \sum_{i=1}^d \frac{\partial}{\partial x_i^2} f, \quad f \in C^2(\mathbb{R}^d, \mathbb{R}).$$

Remark 4.3

Problem:
A continuous and adapted process W is a d-dimensional Brownian motion if and only if

$$M_t^f = f(W_t) - f(W_0) - \frac{1}{2} \int_0^t \Delta f(W_s) ds$$

is a continuous local martingale for every $f \in C^2(\mathbb{R}^d)$.

4.2 Weak Solutions and the Martingale Problem

The problem above gives a characterization of a Brownian motion. The basic idea of the martingale problem is to employ M^f as before. For now, we consider b_i and σ_{ij} to be progressively measurable functionals. Let $(\mathcal{A}'_t)_{t \geq 0}$ be defined by

$$(\mathcal{A}'_t u)(y) := \frac{1}{2} \sum_{i=1}^d \sum_{k=1}^d a_{ik}(t,y) \frac{\partial^2}{\partial x_i \partial x_k} u(y(t)) + \sum_{i=1}^d b_i(t,y) \frac{\partial}{\partial x_i} u(y(t)), t \in [0,\infty), y \in C([0,\infty)^d), u \in C^2(\mathbb{R}^d).$$

Definition 4.4

A probability measure $\mathbb{P}$ on $(C([0,\infty)), \mathcal{B}(C([0,\infty))))$ is called solution to the martingale problem associated with $(\mathcal{A}'_t)_{t \geq 0}$, if under this probability measure

$$M_t^f = f(y(t)) - f(y(0)) - \int_0^t (\mathcal{A}'_s f)(y) ds, \quad \forall t \in [0,\infty),$$

is a $\mathcal{F}_t$ continuous and local martingale for every $f \in C^2(\mathbb{R}^d)$. Here, $\mathcal{F}_t := \mathcal{G}_{t+} = \bigcap_{s>t} \mathcal{G}_t$, although $(\mathcal{G}_t)_{t\geq 0}$ is the augmentation under $\mathbb{P}$ of the canonical filtration of

$$\mathcal{B}_t := \mathcal{B}_t(C([0,\infty)^d)) := \sigma(z(s) : s \in [0,t]) = \varphi_t^{-1}(\mathcal{B}(C([0,\infty)^d))),$$

where

$$\varphi_t : C([0,\infty)^d) \to C([0,\infty)^d), z \mapsto (\varphi_t z)(s) := z(t \wedge s).$$

We show that weak solutions to the functional SDE is equivalent to the local martingale problem. One direction has already been shown: If there is a weak solution $((X,W),(\Omega,\mathcal{F},\mathbb{P}),(\mathcal{F}_t)_{t\geq 0})$ then with Proposition 4.1

$$M_t^f := f(t,X_t) - f(0,X_0) - \int_0^t \left(\frac{\partial}{\partial s} + \mathcal{A}'_s\right) f(s,X_s)ds, \quad \forall f \in C^{1,2}([0,\infty)\times\mathbb{R}^d)$$

is a continuous local martingale, i.e. a solution to the local martingale problem. The converse holds true with the next proposition.

Proposition 4.5

Let $\mathbb{P}$ be a probability measure on $(C([0,\infty)^d), \mathcal{B}(C([0,\infty)^d))$ under which the process M^f is a continuous local $\mathcal{F}_t$-martingale where $f(x) = x_i$ or $f(x) = x_i x_k$ for $i,k = 1,...,d$. Then there exists a Brownian motion W wrt. $(\tilde{\mathcal{F}}_t)_{t\geq 0}$ defined on the extension $(\tilde{\Omega},\tilde{\mathcal{F}},\tilde{\mathbb{P}})$ of $(C([0,\infty)), \mathcal{B}(C([0,\infty)^d)),\mathbb{P})$ such that $((X_t := y(t), W_t),(\tilde{\Omega},\tilde{\mathcal{F}},\tilde{\mathbb{P}}),(\tilde{\mathcal{F}}_t))$ is a weak solution to the functional SDE.

Proof. For $f(x) = X_i$ the differential operator is $(\mathcal{A}'f)(x) = b_i(s,X_s)$. Then by assumption

$$M_t^i := X_t^i - X_0^i - \int_0^t b_i(s,X_s)ds,$$

is a continuous local $\mathcal{F}_t$-martingale under $\mathbb{P}$. We now want to show that

$$\forall t \geq 0 : \quad \langle M^i, M^k\rangle_t = \int_0^t a_{ik}(s,X_s)ds \quad \mathbb{P}-a.s.$$

We do this by showing that $(M_t^i M_t^k - \int_0^t a_{ik}(s, X_s)ds)_{t\geq 0}$ is a continuous local martingale. See that

$$\begin{aligned}
&\int_0^t (X_s^i - X_t^i)b_k(s, X_s)ds + \int_0^t X_s^k - X_t^k)b_i(s, X_s)ds + \int_0^t b_i(s, X_s)ds \int_0^t b_i(s, X_s)ds \\
&= \int_0^t (M_s^i + X_0^i + \int_0^s b_i(l, X_l)dl - (M_t^i + X_0^i + \int_0^t b_i(l, X_l)dl))b_k(s, X_s)ds \\
&+ \int_0^t (M_s^k - X_0^k + \int_0^s b_k(l, X_l)dl - (M_t^k + X_0^k + \int_0^t b_k(l, X_l)dl))b_i(s, X_s)ds \\
&+ \int_0^t b_i(s, X_s)ds \int_0^t b_k(s, X_s)ds \\
&= \int_0^t (M_s^i - M_t^i)b_k(s, X_s)ds + \int_0^t (M_s^k - M_t^k)b_i(s, X_s)ds \\
&\overset{(*)}{=} M_t^i B_t^k - \int_0^t B_s^i dM_s^k - M_t^i \int_0^t dB_s^k + M_t^k B_t^i - \int_0^t B_s^k dM_s^i - M_t^k \int_0^t dB_s^i \\
&= M_t^i B_t^k + M_t^k B_t^i - M_t^i \int_0^t dB_s^k - M_t^k \int_0^t dB_s^i - \int_0^t B_s^i dM_s^k - \int_0^t B_s^k dM_s^i \\
&= M_t^i B_t^k + M_t^k B_t^i - M_t^i B_t^k - M_t^k B_t^i - \int_0^t B_s^i dM_s^k - \int_0^t B_s^k dM_s^i \\
&= -\left(\int_0^t \int_0^s b(l, X_l)dl dM_s^k + \int_0^t \int_0^s b(l, X_l)dl dM_s^i \right). \\
&(HierSindINdexfehler???)
\end{aligned} \tag{4.1}$$

although we used in $(*)$ the product formula for $dB_t^i = \int_0^t b(s, X_s)ds$:

$$M_t^i B_t^i = \int_0^t M_s^i dB_s^i + \int_0^t B_s^i dM_s^i + \underbrace{\langle M_t^i, B_{t)}^i \rangle}_{=0}.$$

The last term is zero, as $(B_t^i)_{t\geq 0}$ is of finite variation. Because of (???) it holds that

$$\forall i = 1, ..., d: \quad \mathbb{P}(\int_0^t |b_i(s, X_s)|ds < \infty, t \geq 0) = 1,$$

which implies that (4.1) is a local martingale. One can show that

$$\begin{aligned}
M_t^i M_t^k - \int_0^t a_{ik}(s, X_s)ds &= M_t^{i,k} - X_0^i M_t^k - X_0^k M_t^i \\
&+ \int_0^t X_s^i dB_s^k - X_t^i B_t^k + \int_0^t X_s^k dB_s^i - X_t^k B_t^i + B_t^i B_t^k \\
&= M_t^{i,k} - X_0^i M_t^k - X_0^k M_t^i \\
&+ \int_0^t (X_s^i - X_t^i)b_i(s, X_s)ds + \int_0^t (X_s^k - X_t^k)b_k(s, X_s)ds + \int_0^t b_i(s, X_s)ds \int_0^t b_k(s, X_s)ds
\end{aligned}$$

This yields

$$\forall t \geq 0: \quad \langle M^i, M^k \rangle_t = \int_0^t a_{ik}(s, X_s)ds \quad \mathbb{P} - a.s.$$

Because the cross variation $\langle M^i, M^j \rangle_t$ is $\mathbb{P}$-a.s. an absolutely continuous function there exists with Theorem 1.5 and extension $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ of $(C([0, \infty)^d), \mathcal{B}(C([0, \infty)^d)), \mathbb{P})$ endowed with the filtration $(\mathcal{F}_t)_{t\geq 0}$ which

satisfies the usual conditions and a matrix valued process $\rho := (\rho_{ij}(t))_{t\geq 0}$ and a Brownian motion $\tilde{W}$ such that

$$\forall t \geq 0, i, j = 1, ..., d : \tilde{P}(\int_0^t \rho_{ij}^2(s)ds < \infty) = 1$$

and

$$\forall t \geq 0, i = 1, ..., d : M_t^i = \sum_{j=1}^d \int_0^t \rho_{ij}(s)d\tilde{W}_s^j \quad \tilde{\mathbb{P}} - a.s.$$

Rewriting the definition of M_t^i yields

$$X_t^i = M_t^i + X_0^i + \int_0^t b_i(s, X_s)ds = X_0^i + \int_0^t b_i(s, X_s)ds + \sum_{j=1}^d \int_0^t \rho_{ij}(s)d\tilde{W}_s^j, \quad t \geq 0.$$

We now want to construct another Brownian motion on $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t\geq 0}, \tilde{\mathbb{P}})$ such that

$$\int_0^t \rho_{ij}(s)d\tilde{W}_s^i = \int_0^t \sigma(s, X_s)dW_s^i, \quad t \geq 0, i, j = 1, ..., d \quad \tilde{\mathbb{P}} - a.s.$$

There exists

$$R \in D := \{(\rho, \sigma) \mid \rho \text{ and } \sigma \text{ are } (d \times d) \text{ matrices: } \quad \rho\rho^T = \sigma\sigma^T\},$$

such that

$$\sigma = \rho R(\rho, \sigma) \quad \text{and} \quad R(\rho, \sigma)R^T(\rho, \sigma) = Id.$$

So R is an orthogonal matrix. Define

$$W_t := \int_0^t R^T(\rho(s), \sigma(s, X_s))d\tilde{W}_s, \quad t \geq 0$$

Because W^i is a continuous local martingale and

$$\begin{aligned}
\langle W^i, W^j\rangle_t &= \sum_{k=1}^d \sum_{l=1}^d \langle \int_0^\cdot R^T(\rho(s), \sigma(s, X_s))_{ik}d\tilde{W}_s^k, \int_0^\cdot R^T(\rho(s), \sigma(s, X_s))_{jl}d\tilde{W}_s^l\rangle_t \\
&\overset{???}{=} \sum_{k=1}^d \sum_{l=1}^d \int_0^t R^T(\rho(s), \sigma(s, X_s))_{ik}R^T(\rho(s), \sigma(s, X_s))_{jl}d\langle\tilde{W}^i, \tilde{W}^l\rangle_s \\
&= \sum_{k=1}^d \sum_{l=1}^d \int_0^t R^T(\rho(s), \sigma(s, X_s))_{ik}R^T(\rho(s), \sigma(s, X_s))_{jl}\delta_{jl}ds \\
&= t
\end{aligned}$$

it is a Brownian motion. Because

$$\forall t \geq 0, y \in C([0, \infty)^d) : \quad a_{ik}(t, y) := \sum_{j=1}^r \sigma_{ij}(t, y)\sigma_{kj}(t, y),$$

it follows that

$$\begin{aligned}
\int_0^t a_{ik}(s, X_s)ds = \langle M^i, M^k\rangle_t &= \sum_{j=1}^d \sum_{l=1}^d \int_0^t \rho_{ij}(s)\rho_{kl}(s)d\langle\tilde{W}^i, \tilde{W}^j\rangle_s \\
&= \sum_{j=1}^d \sum_{l=1}^d \int_0^t \rho_{ij}(s)\rho_{kl}(s)\delta_{jl}ds \\
&= \sum_{j=1}^d \int_0^t \rho_{ij}(s)\rho_{kj}(s)ds \quad \tilde{\mathbb{P}} - a.s.
\end{aligned}$$

It follows that

$$\forall i,k=1,...,d: \quad \tilde{\mathbb{P}}(\sum_{j=1}^{d}\rho_{ij}(t)\rho_{kj}(t)=a_{ik}(t,X_s),a.e.t\geq 0)=1,$$

which finally yields with (4.2) that

$$\forall t\geq 0, i=1,...,d, j=1,...,r: \quad \tilde{\mathbb{P}}(\int_0^t \sigma_{ij}^2(s,X_s)ds<\infty)=1.$$

Thus,

$$\forall t\geq 0, i=1,...,d, j=1,...,r: \quad \tilde{\mathbb{P}}(\int_0^t |b_i(s,X_s)|+\sigma_{ij}^2(s,X_s)ds<\infty)=1$$

and

$$\forall t\geq 0, i=1,...,d: \quad X_t^i=X_0^i+\int_0^t b_i(x,X_s)ds+\sum_{j=1}^{r}\int_0^t \sigma_{ij}(s,X_s)dW_s^j.$$

So $((\tilde{\Omega},\tilde{\mathcal{F}},\tilde{\mathbb{P}}),(X,W),(\mathcal{F}_t)_{t\geq 0})$ is a weak solution. □

Corollary 4.6

The existence of a solution $\mathbb{P}$ to the local martingale problem associated with $(\mathcal{A}'_t)_{t\geq 0}$ is equivalent to the existence of a weak solution $((\tilde{\Omega},\tilde{\mathcal{F}},\tilde{\mathbb{P}}),(X,W),(\mathcal{F}_t)_{t\geq 0})$ to the functional SDE.
Both solutions are related by

$$\mathbb{P}=\tilde{\mathbb{P}}\circ X^{-1},$$

i.e. X induces a probability measure $\mathbb{P}$ on $(C([0,\infty)^d)),\mathcal{B}(C([0,\infty)^d)))$.

Corollary 4.7

The solution $\mathbb{P}$ to the local martingale problem with fixed but arbitrary initial distribution

$$\forall A\in\mathcal{B}(\mathbb{R}^d): \quad \mathbb{P}(X\in C([0,\infty)^d),X\in A)=\mu(A),$$

being unique is equivalent to the uniqueness in sense of law for the SDE

$$dX_t=b(t,X)dt+\sigma(t,X)dW_t, \quad t\geq 0.$$

Given progressively measurable functionals $b_i,\sigma_{ij}:[0,\infty)\times C([0,\infty)^d)\to\mathbb{R}$, for $i=1,...,d, j=1,...,r$, $(\mathcal{A}'_t)_{t\geq 0}$ the associated family of differential operators and μ a probability measure on $\mathcal{B}(\mathbb{R}^d)$ consider the conditions

A) There exists a weak solution to the functional SDE with initial distribution μ.

B) There exists $\mathbb{P}$ solution to the local martingale problem associated with $(\mathcal{A}'_t)_{t\geq 0}$ with

$$\forall A\in\mathcal{B}(\mathbb{R}^d): \quad \mathbb{P}(X_0\in A)=\mu(A).$$

C) There exists $\mathbb{P}$ a solution to the martingale problem associated with $(\mathcal{A}'_t)_{t\geq 0}$ with

$$\forall A\in\mathcal{B}(\mathbb{R}^d): \quad \mathbb{P}(X_0\in A)=\mu(A).$$

Definition 4.8

A probability measure $\mathbb{P}$ on $(C([0,\infty)^d), \mathcal{B}(C([0,\infty)^d)))$ under which

$$M_t^f := f(y(t)) - f(y(0)) - \int_0^t (\mathcal{A}'_s f)(y) ds,$$

is a continuous $\mathcal{F}_t$-martingale, for every $f \in C_0^2(\mathbb{R}^d)$ is called solution to the martingale problem associated with $(\mathcal{A}'_t)_{t\geq 0}$.

Proposition 4.9

A) and B) are equivalent. C) implies A and B. Further, if either

- For every $T \in (0,\infty)$

$$\exists K_T, \forall t \in [0,T], y \in C([0,\infty)^d): \quad \|\sigma(t,y)\| \leq K_T$$

- For every σ_{ij} exists $\tilde{\sigma}_{ij} : [0,\infty) \times \mathbb{R}^d \to \mathbb{R}$ Borel measurable and bounded on compact sets such that

$$\forall t \geq 0, y \in C([0,\infty)^d): \quad \sigma_{ij}(t,y) = \tilde{\sigma}_{ij}(t, y(t)),$$

then it holds that A) implies C).

Proof. The equivalence of A) and B) has been already established. Let We show that C) implies A). Let C) be true. In the local martignale problem $f \in C^2(\mathbb{R}^d)$. So if $\mathbb{P}$ is solution to the martignale problem, it does not need to have compact support. Define the localizing sequence

$$S_k := \inf\{t \geq 0 \mid \|y(t)\| \geq k\},$$

i.e. $S_k \to \infty$, as $k \to \infty$. Because the canonical path process $X_t := y(t)$ posseses continuous paths, for every $t \leq S_k$:

$$y(t) \in \{x \in \mathbb{R}^d \mid \|x\| \leq k\} =: A.$$

Choose $g_k \in C_0^2(\mathbb{R}^d)$ such that for every $x \in A$ holds that $g_k(x) = f(x)$. Then, for every $t \geq 0$:

$$\begin{aligned} M^f_{t\wedge S_k} &= f(y(t\wedge S_k)) - f(y(0)) - \int_0^{t\wedge S_k} (\mathcal{A}f)(s, y(s)) ds \\ &= g_k(y(t\wedge S_k)) - g_k(y(0)) - \int_0^{t\wedge S_k} (\mathcal{A}g_k)(s, y(s)) ds = M^{g_k}_{t\wedge S_k}. \end{aligned}$$

As C) is ture, M^{g_k} is a martignale for every $k \geq 0$. Doobs optimal stopping implies that $(M^{g_k}_{t\wedge S_k})_{t\geq 0}$ is a martignale. So $(M^f_{t\wedge S_k})_{t\geq 0}$ is a martingale and thus M^f a continuous local martingale. □

SECTION 5

MARKOV PROCESSES