

Contents

1	Thurs Sept 10: Syllabus	5
1.1	Instructor Information	5
1.2	Course Information	5
1.3	Method of Evaluation	6
1.4	Additional Policies	7
1.5	Additional Supports	7
1.6	Tentative course schedule	8
1.7	Handing in your work	8
2	Friday Sept 11: What is a population?	9
2.1	Questions	9
2.2	Resources	9
3	Tues Sept 15: Exponential growth - discrete time	11
3.1	Required reading	11
3.2	Questions	11
4	Thurs Sept 17: Getting started with R	13
4.1	Required reading	13
4.2	Questions	13
5	Friday Sept 18: Protection Island 1	15
5.1	Questions	15
6	Tues Sept 22: Protection Island 2	19
6.1	Questions	19
7	Thurs Sept 24: Protection Island 3	23
7.1	Questions	23
8	Fri Sept 25: Doubling times	29
9	Tue Sept 29: Density dependence and logistic growth	31
9.1	Required reading	31
9.2	Questions	31

10 Thurs Oct 1: Solving the logistic growth equation using a computer	33
11 Fri Oct 2: Data and the logistic equation	35
11.1 Questions	35
12 Tues Oct 6: Discrete time models with density dependence using a computer	37
13 Thurs Oct 8: Solving the discrete time models with density dependence using a computer	39
14 Fri Oct 9: Analysis of discrete time models	41
15 Thurs Oct 15: Density-yield and density dependence in births versus deaths	43
16 Tues Oct 20: Balsam fir	45
17 Thurs Oct 22: Stage-structured models	47
18 Fri Oct 23: Stage-structured dynamics	49
19 Tues Oct 27: Yellow columbine	51
20 Thurs Oct 29-Fri Oct 30: Midterm [due Fri Nov 6 at 5pm]	53
21 Tues Nov 3: Evolutionary ecology	55
22 Thurs Nov 5: Evolutionary ecology	57
23 Fri Nov 6: Evolutionary ecology	59
24 Tues Nov 10: Evolutionary ecology	61
25 Thurs Nov 12: Evolutionary ecology	63
26 Fri Nov 13: Evolutionary ecology	65
27 Tues Nov 17: Evolutionary ecology	67
28 Thurs Nov 19: Evolutionary ecology	69
29 Fri Nov 20: Evolutionary ecology	71
30 Final project ideas	73
30.1 How to read scientific papers	74
30.2 Sources for datasets	74
30.3 Grading Rubric: Review	74

<i>CONTENTS</i>	3
30.4 Grading Rubric: Analysis	74

Chapter 1

Thurs Sept 10: Syllabus

1.1 Instructor Information

Instructor: Dr. Amy Hurford

Office: Teaching remotely

Email: ahurford@mun.ca

I will try to reply to emails within 24 hours (excluding evenings, weekends and holidays). I am always available during the lecture times. Please email to request a meeting for a different time. Please check my schedule and suggest a time I am free that works for you.

1.2 Course Information

TR 12.00-12.50pm

F 1-1.50pm

WebEx links for lecture times are on the course Brightspace under Announcements.

Course description:

Population and Evolutionary Ecology is an introduction to the theory and principles of evolutionary ecology and population dynamics. Pre-requisites: BIOL 2600; at least one of BIOL 2010, 2122 or 2210.

Course format:

The course has been re-designed for online delivery. Specifically, no exams that require invigilation are part of the grading scheme because these are challenging to deliver remotely. Pre-recorded lectures limit my ability to interact with

students. Therefore, I have elected to dedicate all lecture time to interacting with students. For each class there is a list of questions you are required to answer and hand-in. Prior to some classes there may be a *Required Reading*, that if completed will allow you to answer the day's assignment questions. Prior to the day of class you should complete the *Required Reading*. In addition, I can most effectively help you if you have read over the questions ahead of time.

Course expectations:

Any students that are disruptive, violating university policies, or acting in a potentially unsafe way will be warned and asked to leave.

Learning goals:

I consider your completed assignments to be a portfolio of your knowledge in population and evolutionary ecology. You will also get some exposure to coding in R. It takes time to become proficient in a programming language, but the time you will spend coding in this class will help you towards becoming more proficient. The course content emphasizes a deeper understanding of fewer concepts. You have the opportunity to further explore a topic of interest to you for the final project.

Required Text and Resources:

The course materials are online at <https://ahurford.github.io/BIOL-3295-Fall-2020/>. In addition you will need a computer to install R and RStudio. This will be covered on Thursday Sept 17 (see Chapter 4). Class announcements and WebEx links will be provided on the course BrightSpace and your assignments are to be submitted to BrightSpace.

1.3 Method of Evaluation

- 27 assignments (equal weighting) - 50%
- Midterm (due Fri Nov 6 at 5pm) - 15%
- Final Project (due Monday Dec 14 at 9am) - 35%

You should aim to complete each assignment before the next class, but assignments will be accepted, without penalty, up to a week later.

Late assignments, labs, and missed midterms, and final exams will be accommodated as described by University Regulation 6.7.3 and 6.7.5 (see <https://www.mun.ca/regoff/calendar/sectionNo=REGS-0474> for Regulations).

1.4 Additional Policies

1.4.1 Accommodation of students with disabilities

Memorial University of Newfoundland is committed to supporting inclusive education based on the principles of equity, accessibility and collaboration. Accommodations are provided within the scope of the University Policies for the Accommodations for Students with Disabilities see www.mun.ca/policy/site/policy.php?id=239. Students who may need an academic accommodation are asked to initiate the request with the Glenn Roy Blundon Centre at the earliest opportunity (see www.mun.ca/blundon for more information).

1.4.2 Academic misconduct

Students are expected to adhere to those principles, which constitute proper academic conduct. A student has the responsibility to know which actions, as described under Academic Offences in the University Regulations, could be construed as dishonest or improper. Students found guilty of an academic offence may be subject to a number of penalties commensurate with the offence including reprimand, reduction of grade, probation, suspension or expulsion from the University. For more information regarding this policy, students should refer to University Regulation 6.12.

1.4.3 Equity and Diversity

A safe learning environment will be provided for all students regardless of race, colour, nationality, ethnic origin, social origin, religious creed, religion, age, disability, disfigurement, sex (including pregnancy), sexual orientation, gender identity, gender expression, marital status, family status, source of income or political opinion.

You should not photograph or record myself, teaching assistants, or other students in the class without first obtaining permission. Accommodation will be made for students with special needs.

The sound should be turned off on phones and computers during class.

1.5 Additional Supports

Resources for additional support can be found at:

- www.mun.ca/currentstudents/student/
- <https://munsu.ca/resource-centres/>

1.6 Tentative course schedule

The course schedule is found in the toolbar of the class materials, see <https://ahurford.github.io/BIOL-3295-Fall-2020/>.

The last day to drop the course without academic prejudice is Wednesday Nov. 4.

1.7 Handing in your work

1.7.1 Making figures to hand-in

The graphs you hand in need to have descriptive axeses and a figure caption. You may put these elements together using a word processing software such as *Microsoft Word*.

1.7.2 Writing R scripts to hand-in

To write your own R scripts follow the guidelines described in Chapter 7 Best Practices of *Quantitative training in Biology*. If you are asked to hand in your R script this means you need to submit an `.R` file on Brightspace.

Chapter 2

Friday Sept 11: What is a population?

For the questions below, submit your answers to Brightspace ideally before the next class. The deadline to submit your answers is Friday Sept 18.

The *Resources* below are sufficient to answer all the questions, however, you are encouraged to find your own textbooks or peer-reviewed articles to answer the questions if you feel comfortable. You can search for textbooks using and the library catalogue and peer-reviewed articles using Web of Science.

2.1 Questions

1. Give a definition of a population from a textbook or peer-reviewed publication. Provide the citation. [2 marks]
2. Find a peer-reviewed paper where a population is studied. Write 1 paragraph discussing how a population is defined for the study and how this compares to your definition of a population given in Question 1. [5 marks]

2.2 Resources

Vandermeer, J.H., Goldberg, D.E., 2013. Population Ecology: First Principles (Second Edition). Princeton University Press, Princeton, United States. [Link](#)

The Princeton Guide to Ecology, edited by Simon A. Levin, et al., Princeton University Press, 2009. ProQuest Ebook Central, [Link](#)

Sacchi, R., Gentili, A., Razzetti, E., Barbieri, F., 2002. Effects of building features on density and flock distribution of feral pigeons *Columba livia* var. *domestica* in an urban environment. *Can. J. Zool.* 80, 48-54. [Link](#)

Chapter 3

Tues Sept 15: Exponential growth - discrete time

Please submit your answers before the next class. You have until Tues Sept 22 to submit your answers.

3.1 Required reading

Vandermeer, J.H., Goldberg, D.E., 2013. Population Ecology: First Principles (Second Edition). Princeton University Press, Princeton, United States. p1-3. [Link](#)

3.2 Questions

1. Suppose $\lambda = 5$ in equation (3) (see the required reading). Explain in 1-2 sentences the meaning of $\lambda = 5$. [1 mark]
2. Suppose the number of lilypads during week 7 is 150. Let $\lambda = 5$, and assume that the units of t are weeks. Use equation (3) to calculate the number of lilypads in week 8. Show your calculations. [2 marks]
3. Use your answer to question 2. to calculate the number of lilypads in week 9. Show your calculations. [2 marks]
4. Equation (4) for the required reading assumes that $N_0 = 1$, however, this formula can be generalized such that

$$N_t = N_0 \lambda^t$$

where N_t is the population size at time, t . Define time such that $t = 0$ is week 7 and t is then the number of weeks since week 7. Use the equation above to answer question 3 and confirm that the answer is the same (i.e., find the population size for week 9, when $\lambda = 5$ and the population size for week 7 is 150). Show your calculations. [2 marks]

5. Use the formula from question 4 to find the population size for week 15, where $N_0 = 1$ and $\lambda = 5$ (the same as question 4). Show your calculations [2 marks]
6. It is important to note that all mathematical formulas should have the same units on both sides of the equals sign, and for each term that is added or subtracted. The units of the population size, N_t , at time, t , are number. The geometric growth rate, λ is unitless. Choose an equation from the required reading and give the units for each of the terms to show that both sides of the equals have the same units. For example, for the equation that appears in question 4, we have:

$$\begin{aligned} N_t &= N_0 \lambda^t \\ (\text{number}) &= (\text{number})(\text{unitless})^{\text{weeks}} \\ (\text{number}) &= (\text{number}) \end{aligned}$$

Note that:

$$\begin{aligned} (\text{unitless}) \times (\text{quantity with units}) &= (\text{quantity with units}) \\ (\text{unitless})^{(\text{quantity})} &= (\text{unitless}) \end{aligned}$$

[2 marks]

7. Although not stated in the reading, $\lambda = 1 + b - d$ where b is the per capita birth rate over one time step (i.e. one week for this example), and d is the fraction of the lilypad population that dies over one time step. The number 1 is considered unitless, what must the units of b and d be? [1 mark]
8. In the reading, $\lambda = 2$. Given that $\lambda = 1 + b - d$, what are some possible values of b and d . [1 mark]
9. [True or False] For discrete time exponential growth (as per the reading), the change in population size from one week to the next depends not so much on the per capita birth rate, but on the difference between the per capita birth rate and the per capita death rate. [1 mark]

Chapter 4

Thurs Sept 17: Getting started with R

You need to have the **R** and **RStudio** softwares installed before you can proceed with the next class. Your answer to the question is due Thursday Sept 24.

4.1 Required reading

You are required to read and complete all the exercises in Chapters 1 *Introduction*, 3 *R and RStudio*, and 4 *Finding your way around RStudio* of:

Quantitative skills for biology <https://ahurford.github.io/quant-guide-all-courses/>

When you are finished you should have **R** and **RStudio** installed on your computer, or you should be familiar with running **RStudio Cloud**.

4.2 Questions

1. Write 1 paragraph describing your experience completing the the exercises. [5 marks. You will receive full marks for attempting the question and submitting an answer.]

Chapter 5

Friday Sept 18: Protection Island 1

The information below is taken from the following source: Newcomb, HR. 1940. Ring-necked pheasant studies on Protection Island in the Strait of Juan de Fuca, Washington. MS thesis. Oregon State University.

- a. Pheasant chicks are born during the summer.
- b. In May 1937, 10 pheasants were introduced to the island. Before the next breeding season there were 35 pheasants.
- c. November 10, 1938 a census estimated 110 pheasants.
- d. October 13, 1939 a census estimated 400 pheasants.

5.1 Questions

1. Read and complete all the exercises in Chapters 6.3 *Variables and assignment* to 6.10 *R packages* and 11 *Making graphs in R* of *Quantitative skills for biology*

HAND IN

- Answer all questions marked HAND IN in the reading

[5 marks]

2. To make a graph of the data listed in b.-d., we need to learn how to work with dates. We will consider two possible approaches:
 - i. Use a built-in R function to convert dates to a format that can be plotted (this question); and

- ii. Convert the dates to number of days since a reference date. Now the dates are numbers and these values can be plotted on the x-axis of a graph (question 3).

In this question, we will proceed with option i. The function we will use is `as.Date()`. You can learn how to use this function using an internet search or by typing the following into your **Console**:

```
?as.Date
```

These files can be difficult to understand (see R Help files. A good way to proceed is to experiment with the function in the **Console**. Try these:

```
as.Date(2012-01-31, format = %Y-%m-%d)
as.Date("2012-01-31", format = "%Y-%m-%d")
```

Note that only the second command is error-free. The first command fails because the date argument for the `as.Date()` function must be a character string, i.e., must be enclosed in `"` (see `?character`).

It is also possible to omit the format argument and just code: `as.Date("2012-01-31")`. The help file notes that when the format argument is not specified, that formats will be tried one by one and an error will be returned if none work. It is advisable to specify the format, as allowing the function to infer the format could introduce errors.

Chapter 6.9 Data structures describes how to make a vector (note a vector is a list of numbers rather than just a single number). We need to make a vector of the dates so that we can make our plot. For example,

```
x <- as.Date(c("2012-01-31", "2012-03-05", "2013-01-11"), format = "%Y-%m-%d")
```

Having completed Chapter 11 Making graphs in R, and having learned how to work with dates, you should now be able to write an R script to make plot using the information in b.-d. above.

HAND IN

- A graph and figure caption, which has dates on the x-axis and the pheasant population size on the y-axis drawing from the information provided in b.-d. You will need to guess the date of ‘before the breeding season’ as stated in b. and you should disclose the value of this guess in the figure caption. See 1.7.1 for more information. [10 marks]
- An R Script that produces the figure described above. See 1.7.2 for more information. [5 marks]

[10 marks]

3. Next, we try approach ii. to make the graph. Under approach ii. we will work with the dates by converting them to the number of days since a

reference date. To do this we will use the `julian()` function, which is part of the `chron` package.

Read Section 4.4 of Quantitative skills for biology regarding installing packages. Install the package `chron` using either the **Install** button on the **Packages** tab, or by using the command `install.packages("chron")` in the **Console** window. Note that the package is only available for use once you check the box on the **Packages** tab or by running the following command in the **Console**:

```
require("chron")
```

After the `chron` package is loaded, we can then query the `julian` function,

```
?julian
```

or use an internet search to better understand how to use it. As the help files can be difficult to understand, another approach to is to try out the function. Try the following:

```
julian(1,1,1970)
julian(1,2,1970)
julian(2,1,1970)
julian(1,1,1971)
julian(1,1,1969)
```

Which argument position of the `julian()` function corresponds to the month? Note also that by default the origin (the origin is when the returned value is 0) is set to January 1, 1970. Experiment by running the following lines of code:

```
julian(1,1,2000)
julian(1,1,2000, origin = c(1,1,1970))
julian(1,1,2000, origin = c(1,1,2000))
```

Finally, we need to make our figure. Recall that the `plot` function requires vectors of equal length for the x- and y-axes. Make a vector of the days since a reference date as follows:

```
ref.day = c(1,1,2000)
x = c(julian(1,1,2000, origin = ref.day), julian(1,1,2002, origin = ref.day))
```

If you run into problems you can query the value of `x` in your console, and you can use `length(x)` to check the length of `x`.

HAND IN

- As for question 2. you need to hand in a graph with descriptive axes and with a figure caption. The y-axis on your graph is population size and the x-axis will be created using the `julian()` function. Be sure to label the x-axis differently than you did in question 2. [10 marks]
- You also need to produce an R Script that makes the figure described above.

See 1.7.2 for more information. Name this file *protection-island-q3.R* [5 marks]

4. For the discrete time exponential (geometric) growth model, ideally the census points should be a fixed amount of time apart, i.e., a day, a week, or a year.

HAND IN

- Provide 1-3 sentences discussing the timing of the census points for Protection Island.

Chapter 6

Tues Sept 22: Protection Island 2

The information below is taken from the following source: Newcomb, HR. 1940. Ring-necked pheasant studies on Protection Island in the Strait of Juan de Fuca, Washington. MS thesis. Oregon State University.

- a. Pheasant chicks are born during the summer.
- b. In May 1937, 10 pheasants were introduced to the island. Before the next breeding season there were 35 pheasants.
- c. November 10, 1938 a census estimated 110 pheasants.
- d. October 13, 1939 a census estimated 400 pheasants.

6.1 Questions

1. Read and complete all the exercises in Chapters 6.3 *Variables and assignment* to 6.10 *R packages* and 11 *Making graphs in R of Quantitative skills for biology*

HAND IN

- Answer all questions marked HAND IN in the reading

[5 marks]

2. To make a graph of the data listed in b.-d., we need to learn how to work with dates. We will consider two possible approaches:
 - i. Use a built-in R function to convert dates to a format that can be plotted (this question); and

- ii. Convert the dates to number of days since a reference date. Now the dates are numbers and these values can be plotted on the x-axis of a graph (question 3).

In this question, we will proceed with option i. The function we will use is `as.Date()`. You can learn how to use this function using an internet search or by typing the following into your **Console**:

```
?as.Date
```

These files can be difficult to understand (see R Help files. A good way to proceed is to experiment with the function in the **Console**. Try these:

```
as.Date(2012-01-31, format = %Y-%m-%d)
as.Date("2012-01-31", format = "%Y-%m-%d")
```

Note that only the second command is error-free. The first command fails because the date argument for the `as.Date()` function must be a character string, i.e., must be enclosed in `"` (see `?character`).

It is also possible to omit the format argument and just code: `as.Date("2012-01-31")`. The help file notes that when the format argument is not specified, that formats will be tried one by one and an error will be returned if none work. It is advisable to specify the format, as allowing the function to infer the format could introduce errors.

Chapter 6.9 Data structures describes how to make a vector (note a vector is a list of numbers rather than just a single number). We need to make a vector of the dates so that we can make our plot. For example,

```
x <- as.Date(c("2012-01-31", "2012-03-05", "2013-01-11"), format = "%Y-%m-%d")
```

Having completed Chapter 11 Making graphs in R, and having learned how to work with dates, you should now be able to write an R script to make plot using the information in b.-d. above.

HAND IN

- A graph and figure caption, which has dates on the x-axis and the pheasant population size on the y-axis drawing from the information provided in b.-d. You will need to guess the date of ‘before the breeding season’ as stated in b. and you should disclose the value of this guess in the figure caption. See 1.7.1 for more information. [10 marks]
- An R Script that produces the figure described above. See 1.7.2 for more information. [5 marks]

[10 marks]

3. Next, we try approach ii. to make the graph. Under approach ii. we will work with the dates by converting them to the number of days since a

reference date. To do this we will use the `julian()` function, which is part of the `chron` package.

Read Section 4.4 of Quantitative skills for biology regarding installing packages. Install the package `chron` using either the **Install** button on the **Packages** tab, or by using the command `install.packages("chron")` in the **Console** window. Note that the package is only available for use once you check the box on the **Packages** tab or by running the following command in the **Console**:

```
require("chron")
```

After the `chron` package is loaded, we can then query the `julian` function,

```
?julian
```

or use an internet search to better understand how to use it. As the help files can be difficult to understand, another approach to is to try out the function. Try the following:

```
julian(1,1,1970)
julian(1,2,1970)
julian(2,1,1970)
julian(1,1,1971)
julian(1,1,1969)
```

Which argument position of the `julian()` function corresponds to the month? Note also that by default the origin (the origin is when the returned value is 0) is set to January 1, 1970. Experiment by running the following lines of code:

```
julian(1,1,2000)
julian(1,1,2000, origin = c(1,1,1970))
julian(1,1,2000, origin = c(1,1,2000))
```

Finally, we need to make our figure. Recall that the `plot` function requires vectors of equal length for the x- and y-axes. Make a vector of the days since a reference date as follows:

```
ref.day = c(1,1,2000)
x = c(julian(1,1,2000, origin = ref.day), julian(1,1,2002, origin = ref.day))
```

If you run into problems you can query the value of `x` in your console, and you can use `length(x)` to check the length of `x`.

HAND IN

- As for question 2. you need to hand in a graph with descriptive axes and with a figure caption. The y-axis on your graph is population size and the x-axis will be created using the `julian()` function. Be sure to label the x-axis differently than you did in question 2. [10 marks]
- You also need to produce an R Script that makes the figure described above.

See 1.7.2 for more information. Name this file *protection-island-q3.R* [5 marks]

4. For the discrete time exponential (geometric) growth model, ideally the census points should be a fixed amount of time apart, i.e., a day, a week, or a year.

HAND IN

- Provide 1-3 sentences discussing the timing of the census points for Protection Island.

Chapter 7

Thurs Sept 24: Protection Island 3

Here is some additional information also taken from: Newcomb, HR. 1940. Ring-necked pheasant studies on Protection Island in the Strait of Juan de Fuca, Washington. MS thesis. Oregon State University.

- a. Pheasant chicks are born during the summer.
- b. In May 1937, 10 pheasants were introduced to the island. Before the next breeding season there were 35.
- c. November 10, 1938 a census estimated 110 pheasants.
- d. October 13, 1939 a census estimated 400 pheasants.
- e. Between the 1938 and 1939 censuses, Newcomb observed that 17 adult birds died.
- f. During the 1938 nesting season: 5.86 eggs/nest. 83.57% of eggs hatched.
- g. During the 1939 nesting season: 8.73 eggs/nest. 64.58% hatched.
- h. During the 1939 nesting season: Average number of chicks per clutch was 6.93.¹
- i. You can assume the sex ratio is 50:50 male to female. Pheasants are a sexually reproducing species.

¹ Note that g. and h. appear to be contradictory.

7.1 Questions

1. Let d be the fraction of population that dies each year. What is d for the ring-tailed pheasant population on Protection Island? Write down any assumptions you have made.

2. b is the per capita number of births each year. What is the value of b ? Write down any assumptions you have made.
3. Recall that $\lambda = 1 + b - d$. What is the value of λ ? Is this population expected to grow over time?
4. Let's assume that the pheasant population on Protection Island grows geometrically (i.e. exponentially) where the geometric growth rate, λ , is the value that you estimated in 5. Let's predict the population size each May beginning with May 1937. Let $N_0 = 10$ and let t be the number of years since May 1937. Recall that when a population grows geometrically,

$$N_t = N_0 \lambda^t$$

You can use R to do this calculation as follows (you should use your value of λ from question 5):

```
t <-1
N0 <-10
lambda <-3
N0*lambda^t
```

where since $t = 1$ the result of `N0*lambda^t` is N_{t+1} , with $t = 1$, such that `N0*lambda^t` is the value of $N_{t+1} = N_2$: the population size two years after May 1937. You can change the value of t and repeat the calculation. Unless you have cleared your workshop it won't be necessary to re-input $N_0 = 10$ and $\lambda = 3$. As such, you can calculate N_3 with the following commands:

```
t <-2
N0*lambda^t
```

HAND IN

- Use R to predict the value of the pheasant population size every year up until May 1940.
7. The approach to calculating the pheasant population size in Question 6 is not very organized. In this question, we will learn how to make a data frame, use a for loop, and use the function `rbind()`.

Read Data structures in *Quantitative training for biology*.

Create a one row dataframe called `df`:

```
df <- data.frame(time = 0, popn.size = 10)
```

Query `df` in your Console to see the data frame you have created. We would like to add successive values of the population size that we calculate to the data frame. To do this we use the `rbind()` function, which binds rows together.


```
new.result <- data.frame(time = 1, popn.size = 20)
df <- rbind(df, new.result)
```

Here the `rbind()` function takes the `df` dataframe and adds the `new.result` data frame as a new row onto the bottom. Note that the code above *overwrites* the value of `df`: that is, `new.result` is added to the bottom of the `df` dataframe (containing only one row), and the result is called `df` (which now has two rows), and the old dataframe `df` (with one row) is overwritten. As such, each time you run the command `df <- rbind(df, new.result)` another row is added to `df`. Try the following:

```
new.result <- data.frame(time = 1, popn.size = 20)
df <- rbind(df, new.result)
df <- rbind(df, new.result)
df <- rbind(df, new.result)
```

If you query the value of `df` you can see that the several rows, all with identical values have been added because we have run the command `df <- rbind(df, new.result)` multiple times while the value of `new.result` is unchanged. Now let's change the value of `new.result` between each time we run the `df <- rbind(df, new.result)` command.

```
new.result <- data.frame(time = 1, popn.size = 20)
df <- rbind(df, new.result)
new.result <- data.frame(time = 2, popn.size = 30)
df <- rbind(df, new.result)
```

Finally, when we do calculations for a sequence of values, it is easier to code this using a `for` loop.

```
lambda <- 1.2
NO <- 10
df <- data.frame(time = 0, popn.size = 10)
for(t in seq(1,4,1)){
  val <- NO*lambda^t
  new.result <- data.frame(time = t, popn.size = val)
  df <- rbind(df, new.result)
}
```

To understand the above code, after copy and pasting it into your **Console**, query the value of `df`: you should see predicted population sizes up until 4 years after May 1937. Now, let's try to understand `seq(1,4,1)`. Let's learn about the `seq()` function by trying it out in the **Console**. What is the result of each of these?


```
seq(-10,10)
seq(-10,5,0.1)
```

The `for` loop works by beginning with `t` equal to the first value of the sequence

and stepping through each value until the final value. The code is written so that quantities that depend on `t` are inside the `for` loop (i.e., enclosed with in the `{}` and those that do not depend on `t` are outside the `for` loop). Note that `val` changes for different values of `t`, `new.result` changes for different values of `t` (because `new.result` has `time = t` and `popn.size = val`, where `val` depends on `t`). Finally, `df` also depends on `t`, because `new.result` depends on `t`. In contrast, `NO` and `lambda` do not change with `t`, so it is more efficient to place the allocated values for these parameters outside of the loop.


We can also plot the results of our calculations:

```
lambda <- 2
NO <- 10
df <- data.frame(time = 0, popn.size = 10)
for(t in seq(1,4,1)){
  val <- NO*lambda^t
  new.result <- data.frame(time = t, popn.size = val)
  df <- rbind(df, new.result)
}
plot(df$time, df$popn.size, typ = "l", xlab = "years since May 1937", ylab = "Population size")
```



If you already have an existing plot you can add new lines using `lines()`. For example,

```
plot(seq(1,4), c(1,3,4,2), ylab = "y-axis", xlab = "x-axis")
lines(seq(1,4), seq(1,4))
```



HAND IN

- Write an R script that builds on the file you have previously make *protection-island-q3.R*. Use the `lines()` command to add the predicted population size assuming geometric growth using the commands described in this section. If you have written the code correctly the result should look something like this:

```
ref.day = c(1,1,2000)
#x = c(julian(1,1,2000, origin = ref.day), julian(1,1,2002, origin = ref.day))
```

8. The geometric growth model is called ‘discrete time’. This equation is appropriate for populations that have regular, pulse reproduction, for example, pheasants that reproduce once per year in the summer. Can you think of examples of other species that reproduce like this?
9. Can you think of any species that reproduce continuously throughout the year? These species are better modelled with a continuous time model.

Chapter 8

Fri Sept 25: Doubling times

[DRAFT]

Chapter 9

Tue Sept 29: Density dependence and logistic growth

[DRAFT]

9.1 Required reading

Vandermeer, J.H., Goldberg, D.E., 2013. Population Ecology: First Principles (Second Edition). Princeton University Press, Princeton, United States. p9-17.
[Link](#)

9.2 Questions

1. What is the equation for continuous time logistic growth? Define all the symbols in the equation.
2. What does dN/dt mean?
3. Equilibrium values

Chapter 10

Thurs Oct 1: Solving the logistic growth equation using a computer

Numerical solutions to CT logistic growth

Chapter 11

Fri Oct 2: Data and the logistic equation

Density dependence + data

11.1 Questions

1. Question 1.9 on p12 of Vandermeer and Gordon.
2. Question 1.10 on p12 of Vandermeer and Gordon.

Chapter 12

**Tues Oct 6: Discrete time
models with density
dependence using a
computer**

Chapter 13

**Thurs Oct 8: Solving the
discrete time models with
density dependence using a
computer**

Chapter 14

Fri Oct 9: Analysis of discrete time models

Chapter 15

**Thurs Oct 15: Density-yield
and density dependence in
births versus deaths**

Chapter 16

Tues Oct 20: Balsam fir

Chapter 17

Thurs Oct 22: Stage-structured models

- The idea of stage-structured models.
- Multiplying matrices.
- Eigenvalues of 2×2 matrix

Chapter 18

Fri Oct 23: Stage-structured dynamics

- Eigenvalues and eigenvectors
- Diagrams

Chapter 19

**Tues Oct 27: Yellow
columbine**

Chapter 20

**Thurs Oct 29-Fri Oct 30:
Midterm [due Fri Nov 6 at
5pm]**

What is chaos? Do real populations exhibit chaos? Discuss the evidence for and against. Provide citations.

Chapter 21

**Tues Nov 3: Evolutionary
ecology**

Chapter 22

Thurs Nov 5: Evolutionary
ecology

Chapter 23

Fri Nov 6: Evolutionary ecology

Chapter 24

**Tues Nov 10: Evolutionary
ecology**

Chapter 25

Thurs Nov 12:
Evolutionary ecology

Chapter 26

**Fri Nov 13: Evolutionary
ecology**

Chapter 27

Tues Nov 17: Evolutionary
ecology

Chapter 28

Thurs Nov 19:
Evolutionary ecology

Chapter 29

**Fri Nov 20: Evolutionary
ecology**

Chapter 30

Final project ideas

[DRAFT]

For your final project you may choose either:

1. Review paper: Read and synthesize information on a new topic in population and evolutionary ecology that we have not considered in class.

or,

2. Analysis: Download and visualize a dataset relevant to population and evolutionary ecology. Discuss your graphs in the context of a principle in population and evolutionary ecology.

Use Dias et al. 1996 *Sources and sinks in population biology* as a guide for what a final project that is a *Review* should aspire to look like.

If you choose to do a final project that is an *Analysis* you should use some of the exercises we have completed for class as a guide (for example, the Doubling time, Protection Island, Yellow columbine, or balsam fir analyses). Note that you will need to write your final project as a report, rather than a series of questions.

Some examples of final project topics are:

Review paper: - Population dynamics in warming environments. - Population dynamics in seasonal environments. - Spatial population dynamics. - Metapopulations.

Analysis:

30.1 How to read scientific papers

Please consult *How to Read a Scientific Paper* (2014) for the recommended approach to ‘reading’ journal articles. Steps 1,2, and 4 are good advice, but step 3 may not be relevant for some readings.

30.2 Sources for datasets

30.3 Grading Rubric: Review

30.4 Grading Rubric: Analysis