

# CANNIBALISM AMONG THE SUPERMASSIVE BLACK HOLES

CHARLES ZIVANCEV<sup>1</sup>, JEREMIAH P. OSTRICKER<sup>1,3</sup>, ANDREAS H.W. KÜPPER<sup>1,2</sup>

*Draft version June 1, 2018*

Abstract

*Subject headings:* Galaxy: kinematics and dynamics

## 1. INTRODUCTION

What is the ultimate fate of the supermassive black holes (SMBH) resident at the centers of all massive galaxies? For the galaxies themselves we know that mergers are frequent from the observed (ADD VAN DOKKUM PAPER) evolution of the size and mass of these systems. But assuming that all galaxy mergers lead to SMBH mergers overpredicts the observations of gravitational waves from the pulsar timing arrays (PTAs) (Kulier et al. (2015), McWilliams et al. (2014), Sesana et al. (2008), Sesana (2013), Sesana et al. (2018), Sesana et al. (2009), NATARJAN). But we do not often see multiple SMBHs at the centers of massive ellipticals. So, what does happen? What we argue in this paper is that as a result of the N-body interactions among the infalling black holes some are ejected, some remain in extended orbits and some (the few most massive ones) do in fact merge but we do not exceed the PTA limits, and the orbiting ones should be detectable via their ability to accrete gas and emit radiation.

A majority of all galaxies with masses  $> 10^8 M_{\odot}$  at  $z = 0$  are believed to host super-massive black holes (SMBHs) in their cores. In a  $\Lambda$ CDM universe, galaxies at the high end of the galaxy-mass function are merger products of numerous such massive galaxies. Even if only a small fraction of merging galaxies hosted an SMBH in their core at the time of merging, it is reasonable to assume that, during a Hubble time of continuous growth, these galaxies will have substantial phases in which they contain two or more SMBHs at the same time.

Here we look at the merger history of XX exemplary galaxies across the galaxy mass spectrum extracted from a cosmological simulation of hierarchical structure formation. We investigate how, after merging with incoming galaxies, SMBHs diffuse into the cores of the hosts and interact with the resident black hole. We show that gravitational interactions of multiple SMBHs are most probable in high-mass galaxies with  $10^{12} M_{\odot} < M < 10^{13} M_{\odot}$ . Galaxies with lower masses have too few mergers with SMBH hosting galaxies. Galaxies with higher masses are more extended, making dynamical friction processes less efficient and hence failing to drive SMBHs into the host galaxy core. Several papers (Inayoshi et al. (2018), McWilliams et al. (2014)) have shown that if all galaxy mergers lead to black hole mergers, the resulting PTA signal would be roughly twice the current limits. One paper (Ryu et al. (2018)) has indicated that dynamical

interactions among multiple black holes, which eject a non-negligible fraction of the mass, may solve this problem. The present paper also addresses this purported dynamical solution.

This paper is organized as follows: in Section 2 we are going to describe the cosmological simulation from which we use the merger history to set up our idealized numerical simulations. We present the few-body integration code, AR-CHAIN that we used for our simulations of SMBH dynamics, and the modifications we made to this code in order to deal with a host galaxy's gravitational potential. In Section 3, we show the results of our four exemplary simulations of galaxies growing with time and acquiring new SMBHs. We analyze how the SMBHs are driven into the core of their new host galaxies and how interaction with the host black hole leads to near-ejections or mergers. The final Section 4 contains a discussion of the result and our conclusions.

## 2. METHODS

### 2.1. Overview of simulation

Our simulations focused on elliptical galaxies with central SMBHs. The galaxies were given a background potential based on the Ostriker-Stone profile (Stone & Ostriker (2015)), which is a three-parameter potential-density pair, whose quantities such as density, potential, and binding energy can be written in closed form. The galaxies were evolved from  $4 > z > 0$ . Orbiting black holes were periodically introduced into the "host" galaxy and their dynamical interaction with the background potential and central SMBH were followed, as described further below.

For the numerical simulations presented here, we used a modified version of the algorithmic chain integrator AR-CHAIN developed by Mikkola & Merritt (2006). It uses algorithmic chain regularization for high-precision integration of few-body dynamics, and is capable of handling velocity-dependent forces efficiently. It includes relativistic post-Newtonian terms up to order PN2.5 (Mikkola & Merritt 2008).

### 2.2. Merger tree

Our merger tree data came from two sources:

- Galactic merger tree data from simulations of Lackner et al. (2012) (hereafter Lackner12), which was centered on galaxies at each redshift slice, providing information on the galaxies' stellar mass, dark matter mass, central BH, and any orbiting black holes within the galaxy.
- SMBH evolution simulations of Kulier et al. (2015) (hereafter Kulier15), which used the large-scale hy-

Correspondence to: csz2104@columbia.edu

<sup>1</sup> Department of Astronomy, Columbia University, 550 West 120th Street, New York, NY 10027, USA

<sup>2</sup> Hubble Fellow

<sup>3</sup> Princeton University, New Jersey, USA

drodynamical galaxy simulations outlined in Cen (2011a,b, 2012a,b, 2013). The data was centered around SMBHs at each redshift slice from  $4 > z > 0$ , giving information for each SMBH's seed mass, accreted and seed mass, its host galaxy (categorically defined), the stellar mass in its host galaxy, and time after  $z=4$  at which the BH entered the host galaxy (if it is not the central BH).

SHOULD I INCLUDE PLOTS AND/OR HISTOGRAMS OF THE DISTRIBUTION OF GALAXY MASSES, STELLAR MASSES, ETC AT  $Z=0$ ?

The merger tree data from Lackner12 and Kulier15 provided 1,830 galaxies in total. However, not all the galaxies were suitable for our simulations. We placed further requirements as follows:

- The galaxies had to exist through the entire simulation ( $4 > z > 0$ ). If they merged with other galaxies, they had to have been the "surviving" galaxy at each merger.
- They had to have accumulated orbiting black holes by  $z = 0$ .

The Kulier15 merger tree collected black holes entering galaxies during the simulation and noted the time after  $z=4$  at which they entered their host galaxy. Nothing further was done with them, so each galaxy had a "running list" of other galaxies/BHs orbiting them at  $z=0$ . Clean up this paragraph.

After finding our subject galaxies, we curve-fit their total masses as a function of time using a 7<sup>th</sup>-order polynomial fit, so as to be able to use their masses in the AR-Chain code. Additionally, we curve-fit each galaxy's accreted+seed central black hole mass to exponential functions of time.

The following cosmological parameters were used in the simulations of both Lackner12 and Kulier15:  $\Omega_M = 0.28$ ,  $\Omega_b = 0.046$ ,  $\Omega_\Lambda = 0.72$ ,  $\sigma_8 = 0.82$ ,  $H_0 = 100h^{-1} \text{Mpc}^{-1} = 70 \text{km s}^{-1} \text{Mpc}^{-1}$ , and  $n = 0.96$ .

### 2.2.1. Galactic Stellar Mass Adjustment

Lackner12 noted that the efficiency of star formation in their simulations, defined as  $f_* = M_*/M_{DM}(\Omega_{DM}/\Omega_b)$ , was approximately 0.6. Compared with the expected range of  $0.10 \lesssim f_* \lesssim 0.15$  that they referenced from Leauthaud et al. (2012), their stellar masses were a factor of roughly 4 times greater. In order to adjust our stellar masses to levels more in line with current observations, we used observational data from Kravtsov et al. (2018), in particular Figure 11.

## 2.3. AR-Chain code

Summary of the code and the modifications we made.

### 2.3.1. Galaxy background potential

For our simulations, the Stone-Ostriker profile (Stone & Ostriker (2015)), which is a three-parameter potential-density pair, whose quantities such as density, potential, and binding energy can be written in closed form. It is essentially an analytic form of a finite, cored isothermal mass distribution:

$$\rho(r) = \frac{\rho_c}{(1 + r^2/r_c^2)(1 + r^2/r_h^2)} \quad (1)$$




FIG. 1.— Original and rescaled stellar masses

Here  $\rho_c$  is the central density,  $r_c$  is the core radius, and  $r_h$  is the outer halo radius.

In order to parametrize this profile,  $r_c$  was given an arbitrary initial value of 100 pc. To calculate  $r_h$ , we began by using the stellar mass to calculate the velocity dispersion at the galaxy's effective radius. In Kulier12, the galaxy's effective radius  $R_e$  and velocity dispersion  $\sigma(R_e)$  at the effective radius are:

$$R_e = 2.5 \text{kpc} \left( \frac{M_*}{10^{11} M_\odot} \right)^{0.73} (1+z)^{-0.98} \quad (2)$$

$$\sigma(R_e) = 190 \text{km/s} \left( \frac{M_*}{10^{11} M_\odot} \right)^{0.2} (1+z)^{0.47} \quad (3)$$

We can then equate the value for  $\sigma(R_e)$  obtained from Equation 3 to the analytic expressions for  $\sigma(R_e)$  in the Stone-Ostriker profile (Eqns 9 and A1-A4). The only unknown is  $r_h$ , which we can solve for using a simple recursive Newton method. Whether  $\sigma_{near}$  or  $\sigma_{far}$  is used from Stone-Ostriker is determined by whether  $R_e$  is less than or greater than  $\sqrt{r_c r_h}$ .

The central density,  $\rho_c$ , can be found from Equation 5 in Stone-Ostriker:

$$M_{tot} = \frac{2\pi^2 r_c^2 r_h^2 \rho_c}{r_h + r_c}, \quad (4)$$

where  $M_{tot}$  is the total dark matter mass.

At each iteration in the code,  $r_h$  is updated according to the stellar mass and Equations 2 and 3. The core radius,  $r_c$ , is recalculated only if an orbiting black hole gets within  $r_c$ . The work done by diffusion, as described in Section 2.3.2 is calculated, the total potential energy is updated, and  $r_c$  is solved for from Equation 8 in Stone-Ostriker.

### 2.3.2. Phase-space diffusion

Weak encounters with background stars will let the SMBHs diffuse through phase space while they are orbiting within the gravitational potential of the galaxy. The diffusion can be expressed as change in velocity of an SMBH by  $\Delta \vec{v}$  per unit time. We can split this change into a component along the direction of motion of the

SMBH, and one perpendicular to that. Following [Binney & Tremaine \(2008\)](#), the diffusion coefficients can be expressed as

$$D[\Delta v_{\parallel}] = -\frac{4\pi G^2 \rho(r) M_{\bullet} \ln \Lambda}{\sigma^2} f(\chi), \quad (5)$$

$$D[(\Delta v_{\parallel})^2] = \frac{4\sqrt{2}\pi G^2 \rho(r) M_{\bullet} \ln \Lambda}{\sigma} \frac{f(\chi)}{\chi}, \quad (6)$$

$$D[(\Delta \vec{v}_{\perp})^2] = \frac{4\sqrt{2}\pi G^2 \rho(r) M_{\bullet} \ln \Lambda}{\sigma} \left[ \frac{\text{erf}(\chi) - f(\chi)}{\chi} \right] \quad (7)$$

where  $\Delta v_{\parallel} \equiv \Delta \vec{v} \cdot \vec{v}/v$  is the velocity change in direction of motion, and  $\Delta \vec{v}_{\perp} \equiv \Delta \vec{v} - \Delta v_{\parallel} \cdot \vec{v}/v$  is the velocity change perpendicular to the direction of motion. Here,  $M_{\bullet}$  is the mass of the black hole, and  $\chi = \frac{v}{\sqrt{2}\sigma(r)}$ . The function  $f(\chi)$  is given by

$$f(\chi) = \frac{1}{2\chi^2} \left( \text{erf}(\chi) - \frac{2\chi}{\sqrt{\pi}} \exp(-\chi^2) \right). \quad (8)$$

We approximate the factor  $\Lambda$  in the Coulomb logarithm as

$$\Lambda = \left( \frac{M_{NSC}}{M_{\bullet}} \right) \left( \frac{r}{r_h} \right). \quad (9)$$

We can identify Equation 5 as the dynamical friction term. The second term introduces a variance of the friction term, and even allows the SBHs to be accelerated when the velocity of a SBH gets sufficiently small. The third term introduces a change in velocity perpendicular to the direction of motion of the SBH. It is a randomly oriented vector, and hence causes the SBHs to execute a random walk in phase space. The last two terms will establish that the SBHs are ultimately in energy equipartition with the background stars. The velocity changes  $\Delta v_{\parallel}$  and  $\Delta \vec{v}_{\perp}$  per unit time  $\Delta t$  can be computed with the above equations. Both changes are normally distributed, where the mean,  $\mu$ , and the variance,  $\Sigma$ , of the distributions are given by

$$\mu_{\parallel} = D[\Delta v_{\parallel}] \Delta t, \quad (10)$$

$$\Sigma_{\parallel} = D[(\Delta v_{\parallel})^2] \Delta t, \quad (11)$$

$$\mu_{\perp} = 0, \quad (12)$$

$$\Sigma_{\perp} = D[(\Delta \vec{v}_{\perp})^2] \Delta t. \quad (13)$$

We compute the diffusion coefficients for each black hole at each time step, and modify its velocity on a Monte Carlo basis. For each time step we draw a random orientation before adding the perpendicular velocity change to the respective SBH. Hence, the SBH's modified velocity,  $\vec{v}_f$ , is computed using

$$\vec{v}_f = \vec{v}_0 + \Delta v_{\parallel} \hat{v}_{\parallel} + \Delta v_{\perp} \hat{v}_{\perp}, \quad (14)$$

$$\Delta v_{\parallel} = \mathcal{N}(\mu_{\parallel}, \Sigma_{\parallel}), \quad (15)$$

$$\Delta v_{\perp} = \mathcal{N}(\mu_{\perp}, \Sigma_{\perp}). \quad (16)$$

The change of energy,  $dE_{BH}$ , of the orbiting black hole due to phase-space diffusion is given back to the stellar background potential, with  $dE = -dE_{BH}$ . As a consequence of this energy transfer, inspiralling black holes will cause an expansion of the NSC. For this purpose we calculate the change in potential energy,  $dW$ , of the

stellar system using

$$E = T + W = \frac{1}{2}W, \quad (17)$$

$$dW = -2 dE_{BH}, \quad (18)$$

where we made use of the virial theorem  $2T + W = 0$ . With this change in potential energy we can calculate a new radius for the stellar background potential at each integration step. For the Plummer sphere the new scale radius can be calculated as

$$a_{new} = a \left( 1 + \frac{32a dW}{3\pi G M_{NSC}^2} \right)^{-1}. \quad (19)$$

### 2.3.3. Gravitational wave recoils

The code AR-CHAIN includes post-Newtonian terms up to order 2.5. The SMBHs can therefore merge via gravitational wave emission. We include gravitational wave recoils following the prescription outlined in [Kulier15](#), which is based on the fitting formula by [Lousto et al. \(2012\)](#). To save computational time, we assume that a merger will be inevitable when the separation between two SBHs gets smaller than 1.0 Schwarzschild radius. At the moment of the merger, we assume that the spin vectors of the two SBHs are randomly aligned. This results in kick velocities of up to several thousand  $\text{km s}^{-1}$ , with a median kick of  $\approx 290 \text{ km s}^{-1}$ . Since our simulations focus on NSC with relatively low escape velocities, this implies that a majority of the merging SBHs escape from the NSCs.

Black holes can also eject each other via strong three-body interactions. We remove SBHs from the simulations once they move beyond 1 kpc from the NSC, assuming that it will take them more than a Hubble time to find their way back into the center of the host galaxy.

### 2.4. Simulation setup

Note to discuss: Injection, merging, escape, effective radius, set of galaxies

Using the merger tree data, orbiting black holes were injected into its host galaxy at redshift  $z$ , at a distance from the galactic center of  $R_e$  (Eqn. 2). Their initial velocity was arbitrarily chosen to be circular,  $v_c = \sqrt{M(R_e)/R_e}$ , with  $v_x$ ,  $v_y$ , and  $v_z$  randomly chosen.

Note: we only inject the BHs into the simulation that have a  $t_{fric}$  smaller than 100 times the Hubble time.

### 2.5. Description of Simulations

## 3. RESULTS

### 3.1. Mergers

Note to work on: A: two mergers early on, and then two more at around 10-12 Gyr B: 4 mergers across the final 8 Gyr C: two mergers in the final Gyrs D: ?

### 3.2. Ejections

No complete ejections, but kicks to higher energy orbits. A: one in a three-body encounter B: 7-ish kicks, some in nearly 4-body encounters C: none D: ?

TABLE 1  
GALAXY CHARACTERISTICS

Galaxy	$M_{gal} [M_{\odot}]$		$M_* [M_{\odot}]$		$M_{BH} [M_{\odot}]$	
	Init	Final	Init	Final	Init	Final
A	$7.42 \times 10^{11}$	$1.09 \times 10^{14}$	$1.22 \times 10^{11}$	$1.75 \times 10^{13}$	$4.80 \times 10^6$	$2.19 \times 10^9$
B	$1.11 \times 10^{12}$	$3.41 \times 10^{13}$	1.80311	$5.58 \times 10^{12}$	$1.94 \times 10^8$	$1.46 \times 10^9$
C	$7.23 \times 10^{10}$	$3.48 \times 10^{12}$	$1.18 \times 10^{10}$	$5.68 \times 10^{11}$	$9.47 \times 10^4$	$2.70 \times 10^8$
D	$3.92 \times 10^{10}$	$1.34 \times 10^{12}$	$6.47 \times 10^9$	$2.20 \times 10^{11}$	$9.95 \times 10^5$	$8.32 \times 10^7$

TABLE 2  
SMBH STATISTICS

Galaxy	Infalling SMBHs	SMBHs with $t_{fric} < 100 t_H$	Mergers
A	206	19	4
B	29	12	4
C	3	3	2
D	4	4	2




FIG. 2.— default

### 3.3. Stalling SMBH binaries

A: two periods of BH binary in center for  $> 6$  Gyr, up to 6 Gyr  
 B: two short periods of BHBs  
 C: 2 classic BHB periods of about 1 Gyr each  
 D: ?

### 3.4. Stalling SMBHs in the halo

A: 95% never make it into the central kpc  
 B: still about 90% are in the halo  
 C:  
 D: ?

## 4. CONCLUSIONS

What do we want to say?

## ACKNOWLEDGEMENTS




FIG. 3.— default




FIG. 4.— default

The authors would like to thank Andrea Kulier and Claire Lackner for providing merger tree data. AHWK acknowledges support by NASA through Hubble Fellowship grant HST-HF-51323.01-A awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555.

## REFERENCES

- Binney, J., & Tremaine, S. 2008, *Galactic Dynamics: Second Edition* (Princeton University Press) [2.3.2](#)
- Cen, R. 2011a, *ApJ*, 741, 99 [2.2](#)
- . 2011b, *ApJ*, 742, L33 [2.2](#)
- . 2012a, *ApJ*, 753, 17 [2.2](#)
- . 2012b, *ApJ*, 748, 121 [2.2](#)
- . 2013, *ApJ*, 770, 139 [2.2](#)
- Inayoshi, K., Li, M., & Haiman, Z. 2018, ArXiv e-prints, arXiv:1804.03143 [1](#)
- Kravtsov, A. V., Vikhlinin, A. A., & Meshcheryakov, A. V. 2018, *Astronomy Letters*, 44, 8 [2.2.1](#)
- Kulier, A., Ostriker, J. P., Natarajan, P., Lackner, C. N., & Cen, R. 2015, *ApJ*, 799, 178 [1](#), [2.2](#)
- Lackner, C. N., Cen, R., Ostriker, J. P., & Joung, M. R. 2012, *MNRAS*, 425, 641 [2.2](#)




FIG. 5.— default

Leauthaud, A., George, M. R., Behroozi, P. S., et al. 2012, *ApJ*, 746, 95 [2.2.1](#)  
 Lousto, C. O., Zlochower, Y., Dotti, M., & Volonteri, M. 2012, *Phys. Rev. D*, 85, 084015 [2.3.3](#)  
 McWilliams, S. T., Ostriker, J. P., & Pretorius, F. 2014, *ApJ*, 789, 156 [1](#)  
 Mikkola, S., & Merritt, D. 2006, *MNRAS*, 372, 219 [2.1](#)  
 —. 2008, *AJ*, 135, 2398 [2.1](#)

Ryu, T., Perna, R., Haiman, Z., Ostriker, J. P., & Stone, N. C. 2018, *MNRAS*, 473, 3410 [1](#)  
 Sesana, A. 2013, *MNRAS*, 433, L1  
 Sesana, A., Haiman, Z., Kocsis, B., & Kelley, L. Z. 2018, *ApJ*, 856, 42  
 Sesana, A., Vecchio, A., & Colacino, C. N. 2008, *MNRAS*, 390, 192  
 Sesana, A., Vecchio, A., & Volonteri, M. 2009, *MNRAS*, 394, 2255  
 Stone, N. C., & Ostriker, J. P. 2015, *ApJ*, 806, L28 [2.1](#), [2.3.1](#)