

Problem 1

Let v_1, \dots, v_n be mutually orthogonal vectors in an inner product space V . Show that

$$\left\| \sum_{k=1}^n v_k \right\|^2 = \sum_{k=1}^n \|v_k\|^2.$$

Proof:

$$\begin{aligned} \left\| \sum_{k=1}^n v_k \right\|^2 &= \left\langle \sum_{k=1}^n v_k, \sum_{k=1}^n v_k \right\rangle \\ &= \sum_{i=1}^n \left\langle \sum_{k=1}^n v_k, v_i \right\rangle \\ &= \sum_{i=1}^n \langle v_i, v_i \rangle && \text{since for } i \neq j, \langle v_i, v_j \rangle = 0 \\ &= \sum_{i=1}^n \|v_i\|^2 \end{aligned}$$

Problem 2

Let V be an inner product space and fix $w \neq 0$ in V . We define the one-dimensional projection

$$P_w : V \rightarrow V; P_w(v) := \frac{\langle v, w \rangle}{\langle w, w \rangle} w.$$

- (i) Prove that $v - P_w(v) \perp P_w(v)$.
- (ii) Show that $P_w : V \rightarrow V$ is a linear operator with $\|P_w\|_{\text{op}} = 1$.
- (iii) Show that $P_w \circ P_w = P_w$.

Problem 3

Let V be an inner product space. Prove the reverse Cauchy-Schwarz Inequality which states

$$v, w \in V, \text{ and } |\langle v, w \rangle| = \|v\| \|w\| \Rightarrow v = \alpha w.$$

Problem 4

Let V be an inner product space. Then, for any $v, w \in V$, show that

$$\|v + w\|^2 + \|v - w\|^2 = 2\|v\|^2 + 2\|w\|^2$$

Problem 5

Let $\lambda = (\lambda_k)_k$ belong to ℓ_∞ . Show that the map

$$D_\lambda : \ell_2 \rightarrow \ell_2; D_\lambda((\xi_k)_k) = (\lambda_k \xi_k)_k$$

is well-defined, linear, and bounded with $\|D_\lambda\|_{\text{op}} = \|\lambda\|_\infty$

Problem 6

Consider the vector space $C([0, 2\pi])$ equipped with

$$\langle f, g \rangle := \frac{1}{2\pi} \int_0^{2\pi} f(t) \overline{g(t)} dt.$$

- (i) Show that this pairing defines an inner product on $C([0, 2\pi])$.

Proof: We will show that $\langle f, g \rangle$ satisfies the axioms of the inner product.

Addition:

$$\begin{aligned}\langle f_1 + f_2, g \rangle &= \frac{1}{2\pi} \int_0^{2\pi} (f_1(t) + f_2(t)) \overline{g(t)} dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} (f_1(t) \overline{g(t)} + f_2(t) \overline{g(t)}) dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} f_1(t) \overline{g(t)} dt + \frac{1}{2\pi} \int_0^{2\pi} f_2(t) \overline{g(t)} dt \\ &= \langle f_1, g \rangle + \langle f_2, g \rangle.\end{aligned}$$

Scalar Multiplication:

$$\begin{aligned}\langle \alpha f, g \rangle &= \frac{1}{2\pi} \int_0^{2\pi} (\alpha f(t)) \overline{g(t)} dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} \alpha (f(t) \overline{g(t)}) dt \\ &= \alpha \left(\frac{1}{2\pi} \int_0^{2\pi} f(t) \overline{g(t)} dt \right) \\ &= \alpha \langle f, g \rangle.\end{aligned}$$

Conjugation:

$$\begin{aligned}\overline{\langle g, f \rangle} &= \frac{1}{2\pi} \int_0^{2\pi} \overline{g(t) \overline{f(t)}} dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(t) \overline{g(t)} dt \\ &= \langle f, g \rangle.\end{aligned}$$

Positive Definition:

$$\begin{aligned}\langle f, f \rangle &= \frac{1}{2\pi} \int_0^{2\pi} f(t) \overline{f(t)} dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} |f(t)|^2 dt \\ &\geq 0.\end{aligned}$$

For $\langle f, f \rangle = 0$, we have that the integral equals zero — since f is continuous, it means that if $|f(t)|^2 > 0$ for some $t_0 \in [0, 2\pi]$, then $|f(t)|^2 \neq 0$ on some interval $[t_0 - \delta, t_0 + \delta]$, meaning the integral can only equal zero if f is $\mathbb{0}_f$ on $[0, 2\pi]$.

(ii) For $n \in \mathbb{Z}$, set $e_n(t) = \cos(nt) + i \sin(nt)$. Show that the family $\{e_n\}_{n \in \mathbb{Z}}$ is orthonormal.

Proof: We will show that $\{e_n\}_{n \in \mathbb{Z}}$ is orthonormal by showing that $\langle e_n, e_n \rangle = 1$ and $\langle e_n, e_m \rangle = 0$ for $m \neq n$.

$$\begin{aligned}\langle e_n, e_n \rangle &= \frac{1}{2\pi} \int_0^{2\pi} (\cos(nt) + i \sin(nt))(\cos(nt) - i \sin(nt)) dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} (\cos^2(nt) + \sin^2(nt)) dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} dt \\ &= 1 \\ \langle e_n, e_m \rangle &= \frac{1}{2\pi} \int_0^{2\pi} (\cos(nt) + i \sin(nt))(\cos(mt) - i \sin(mt)) dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} (\cos(mt) \cos(nt) + i \sin(nt) \cos(mt) - i \sin(mt) \cos(nt) + \sin(nt) \sin(mt)) dt \\ &= \frac{1}{2\pi} \left(\int_0^{2\pi} (\cos(mt) \cos(nt) + \sin(nt) \sin(mt)) dt + i \int_0^{2\pi} (\sin(nt) \cos(mt) - \sin(mt) \cos(nt)) dt \right) \\ &= 0.\end{aligned}$$

Problem 7

Let V be any normed space, $p \in [1, \infty]$, and suppose $T : \ell_p^n \rightarrow V$ is linear. Show that T is bounded.

Problem 8

Let $\mathbb{P}[0, 1] = \{\sum_0^n a_k x^k \mid a_k \in \mathbb{C}\} \subseteq C([0, 1])$ denote the linear subspace of all polynomial functions equipped with the uniform norm $\|\cdot\|_u$ inherited from $C([0, 1])$. We define the map

$$D : \mathbb{P}[0, 1] \rightarrow \mathbb{P}[0, 1]$$

$$D(p(x)) = p'(x).$$

Show that D is unbounded.

Problem 9

Let V be an infinite-dimensional normed space. Show that there is a linear functional $\varphi : V \rightarrow \mathbb{F}$ that is unbounded.

Problem 10

Let $a, b \in \mathbb{M}_n$. Show the following properties of the operator norm.

$$(i) \quad \|a\|_{\text{op}} = \sup \left\{ |\langle a\xi, \eta \rangle| \mid \xi, \eta \in B_{\ell_2^n} \right\}$$

$$(ii) \quad \|a^*\|_{\text{op}} = \|a\|_{\text{op}}$$

$$(iii) \quad \|ab\|_{\text{op}} \leq \|a\|_{\text{op}} \|b\|_{\text{op}}$$

$$(iv) \quad \|a^*a\|_{\text{op}} = \|a\|_{\text{op}}^2$$