

# Homeomorphism between the Open Ball Topology on $\mathbb{R}^2$ and the Open Rectangle Topology on $\mathbb{R}^2$

Avinash Iyer

Occidental College

Word Count: 832.

## Abstract

In this paper, I introduce the concept of metric spaces (including the definition of distance metrics, open sets, and continuity), topologies on sets, and homeomorphisms. With this knowledge, we can prove that the Open Rectangle Topology, where  $\bigcup_{i=1}^n (a_i, b_i) \times (c_i, d_i) \subseteq \mathbb{R}^2$  is defined to be open, and the Open Ball Topology, where  $\bigcup_{i=1}^n V_{\epsilon_i}((x, y)) \subseteq \mathbb{R}^2$  is defined to be open, are homeomorphic.

## Metric on $\mathbb{R}$ and $\mathbb{R}^2$

Consider a set  $X$ , with  $a, b \in X$ . It's often useful to consider the idea of a distance between  $a$  and  $b$ ,  $d(a, b)$ . This distance function must map every pair of points to some positive number.

$$d : X \times X \rightarrow \mathbb{R}^+$$

Additionally, it has to have the following properties:

- Zero distance property:  $d(a, a) = 0$  and if  $d(a, b) = 0$ , then  $a = b$ .
- Commutativity:  $d(a, b) = d(b, a)$ .

- Triangle Inequality:  $d(x, z) = d(x, y) + d(y, z)$ .

This defines a **distance metric** on  $X$ . For instance, for  $x, y \in \mathbb{R}^n$ , we define the **Euclidean metric** as follows:

$$d(x, y) = \left( \sum_{i=1}^n (x_i - y_i)^2 \right)^{1/2}$$

In  $\mathbb{R}^1 = \mathbb{R}$ , this is equivalent to  $|x - y|$ .

## Open Balls and Open Sets

In  $\mathbb{R}^n$  with the Euclidean metric, we define the **open ball** of radius  $\varepsilon > 0$  about the point  $x$  as follows:

$$V_\varepsilon = \{p \mid d(x, p) < \varepsilon\}$$

In  $\mathbb{R}$ , this equivalent to the open interval  $(x - \varepsilon, x + \varepsilon)$ .

In  $A \subseteq X$ , if  $\forall x \in A$ ,  $\exists \varepsilon > 0$  such that  $V_\varepsilon(x) \subseteq A$ , then  $A$  is an **open set**. If  $A = \emptyset$  or  $A = X$ , then  $A$  is open. Every open ball is an open set.

**Proposition:** Let  $A$  and  $B$  be open sets in a metric space  $X$ . Then,  $A \cap B$  and  $A \cup B$  are open sets in  $X$ .

**Proof:** Let  $a \in A$  and  $b \in B$ . Then,  $\exists \varepsilon_a > 0$  such that  $V_{\varepsilon_a}(a) \subseteq A$ , meaning that  $V_{\varepsilon_a}(a) \subseteq A \cup B$ , and similarly for  $b$  and  $\varepsilon_b$ .

If  $A = \emptyset$ , then  $A \cap B = \emptyset$ , meaning  $A \cap B$  is open. Otherwise, suppose  $A \cap B \neq \emptyset$ . Let  $x \in A \cap B$ , meaning  $x \in A$  and  $x \in B$ . So,  $\exists \varepsilon_1 > 0$  such that  $V_{\varepsilon_1}(x) \subseteq A$ .

Specifically, define  $\varepsilon_1$  to be the maximum such value. Similarly,  $\exists \varepsilon_2 > 0$  such that  $V_{\varepsilon_2}(x) \in B$ , where  $\varepsilon_2$  is the maximum such value. Let  $\varepsilon = \min\{\varepsilon_1, \varepsilon_2\}$ . Then,  $V_\varepsilon \subseteq V_{\varepsilon_1}$  and  $V_\varepsilon \subseteq V_{\varepsilon_2}$ , meaning  $V_\varepsilon \subseteq A$  and  $V_\varepsilon \subseteq B$ , so  $V_\varepsilon \subseteq A \cap B$ .

## Continuity

Consider a function in a metric space  $f : X \rightarrow Y$ . Just as with functions in the real numbers, we can consider the continuity of  $f$ .<sup>1</sup> Specifically,  $f$  is continuous if  $\forall x_1, x_2 \in X$

$$(\forall \varepsilon > 0) (\exists \delta > 0) \text{ such that } d(x_1, x_2) < \delta \Rightarrow d(f(x_1), f(x_2)) < \varepsilon.$$

Alternatively, this can be phrased as

$$(\forall \varepsilon > 0) (\exists \delta > 0) \text{ such that } \forall a, a \in V_\delta(x) \Rightarrow f(a) \in V_\varepsilon(f(x)).$$

**Proposition:** If  $f : X \rightarrow Y$  is continuous, then  $\forall x \in X, f(V_\delta(x)) \subseteq V_\varepsilon(f(x))$  for some  $\varepsilon, \delta > 0$ .

**Proof:** Let  $\varepsilon > 0$  and  $f : X \rightarrow Y$  be continuous. For  $x, a \in X$ , define  $\delta > 0$  such that  $d(x, a) < \delta \Rightarrow d(f(x), f(a)) < \varepsilon$ , which exists by definition of continuity.

Then,  $a \in V_\delta(x)$ , meaning  $f(a) \in f(V_\delta(x))$ , and  $f(a) \in V_\varepsilon(f(x))$  as defined earlier.

Therefore,  $f(V_\delta(x)) \subseteq V_\varepsilon(f(x))$ .

Using this result, we can prove an important fact of continuity.

**Proposition:**  $f : X \rightarrow Y$  is continuous if and only if the preimage of every open set in  $Y$  is open in  $X$ .

**Proof:** ( $\Rightarrow$ ) Let  $f$  be a continuous function. If  $B \subseteq Y$  is open, then  $\forall y \in B$ ,  $\exists \epsilon > 0$  such that  $V_\epsilon(y) \subseteq B$ . Since  $x \in Y$ ,  $\exists x \in X$  such that  $f(x) = y$ , meaning  $V_\epsilon(f(x)) \subseteq B$ .

By our previous result, we know  $\exists \delta > 0$  such that  $f(V_\delta(x)) \subseteq V_\epsilon(f(x)) \subseteq B$ , meaning  $V_\delta(x) \subseteq f^{-1}(B)$ . Therefore,  $f^{-1}(B)$  is open.

( $\Leftarrow$ ) Let  $f : X \rightarrow Y$  be defined such that for every open  $B \subseteq Y$ ,  $f^{-1}(B)$  is open in  $X$ . Let  $B$  be open in  $Y$ , and  $x \in f^{-1}(B)$ , meaning  $f(x) \in B$ .

Since  $B$  is open in  $Y$  and  $f^{-1}(B)$  is open in  $X$ , we know  $\exists \epsilon > 0$  such that  $V_\epsilon(f(x)) \subseteq B$ .

By definition, every open ball is an open set, meaning  $f^{-1}(V_\epsilon(f(x)))$  is open in  $X$ . Similarly, since  $x \in f^{-1}(B)$ ,  $\exists \delta > 0$  such that  $V_\delta(x) \subseteq f^{-1}(V_\epsilon(f(x)))$ .

Therefore,  $\forall \epsilon > 0$ , we have a suitable  $\delta$  defined as above such that  $\forall a$  such that  $a \in V_\delta(x)$ ,  $f(a) \in V_\epsilon(f(x))$ .

## Topology Induced by Metric

We can generalize the concept of open sets and continuity using the properties above to sets without a metric. For this purpose, on a set  $L$ , we define a **topology** on  $L$ ,  $\mathcal{T}(L)$ , to be a set of open sets defined as follows:

- $L, \emptyset \in \mathcal{T}(L)$
- if  $A, B \in \mathcal{T}(L)$ , then  $A \cap B$  and  $A \cup B \in \mathcal{T}(L)$

For a metric space  $X$ , the topology **induced** by the metric is the set of open sets in  $X$ . Similar

to continuity in a metric space, we define continuity on a function between topological spaces by the following:

The function  $f : X \rightarrow Y$  between topological spaces  $X$  and  $Y$  is continuous if and only if the preimage of every open set in  $Y$  is open in  $X$ .

## Homeomorphism

Recall that a given topological space  $X$  has two properties:

- Elements:  $X = \{x_1, x_2, \dots\}$
- Open Sets:  $A_1, A_2, A_3, \dots \subseteq X$

Consider two topological spaces,  $X$  and  $Y$ . If we want to consider an idea that  $X$  and  $Y$  are “equal,” we must find a correspondence such that both the open sets and the elements correspond to each other one to one.

Let  $f : X \rightarrow Y$  be defined as follows:

- $f$  is a bijection.
- $f$  is continuous.
- $f^{-1}$  is continuous.

Then we know that each open set in  $X$  corresponds to one, and only one, open set in  $Y$ , and each element in  $X$  corresponds to one, and only one, element in  $Y$ . The function  $f$  is a **homeomorphism**, and if such an  $f$  exists, then  $X$  is **homeomorphic** to  $Y$ .

## Two Topologies on $\mathbb{R}^2$

Consider two topologies,  $\mathcal{T}_R$  and  $\mathcal{T}_V$ , defining open sets in  $\mathbb{R}^2$ .

- The “Open Rectangle Topology” on  $\mathbb{R}^2$  is defined by a union of Cartesian products of open intervals.

$$\bigcup_{i=1}^n (a_i, b_i) \times (c_i, d_i) \in \mathcal{T}_R$$

- The “Open Ball Topology” on  $\mathbb{R}^2$  is defined by a union of open balls.

$$\bigcup_{i=1}^n V_{\epsilon_i}(x) \in \mathcal{T}_V$$

**Proposition:**  $\mathcal{T}_V(\mathbb{R}^2)$  and  $\mathcal{T}_R(\mathbb{R}^2)$  are homeomorphic.

**Proof:** Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be defined as  $f(x) = x$ . We will show that every open set in  $\mathcal{T}_R$  is open in  $\mathcal{T}_V$ , and vice versa.

Consider an open set  $A \in \mathcal{T}_R$  — we will show that this open set is open in  $\mathcal{T}_V$ .

Let  $(a, b) \times (c, d) \subseteq A$  be an open rectangle, and  $x = (x_a, y_a) \in (a, b) \times (c, d)$ .

We can always find  $\epsilon = \min \{|x_a - a|, |x_a - b|, |y_a - c|, |y_a - d|\}$ . Therefore, this open rectangle is open in  $\mathcal{T}_V$ , and since every open set in  $\mathcal{T}_R$  is a union of open rectangles, and each open rectangle is open in  $\mathcal{T}_V$ , the union of these open rectangles is open in  $\mathcal{T}_V$ .

Consider an open set  $B \in \mathcal{T}_V$ . Let  $V_\epsilon(x) \subseteq B$  be an open ball centered at  $x = (x_x, x_y)$ .

Let  $a = (a_x, a_y) \in V_\epsilon(x)$  be a point in  $V_\epsilon(x)$ .

Since  $V_\varepsilon(x)$  is open,  $\exists \delta$  such that  $V_\delta(a) \subseteq V_\varepsilon(x)$ . Define the open rectangle to be  $A = (a_x - \delta/\sqrt{2}, a_x + \delta/\sqrt{2}) \times (a_y - \delta/\sqrt{2}, a_y + \delta/\sqrt{2})$ . Then, this open rectangle is a subset of  $V_\delta(a) \subseteq V_\varepsilon(x)$ , meaning  $V_\varepsilon(x)$  is open in  $\mathcal{T}_R$ . Since  $\mathcal{T}_V$  is a union of open balls, and the union of these open sets is open, the union of open balls is open in  $\mathcal{T}_R$ .

## Acknowledgments

I would like to thank professor Ramin Naimi, whose Math 362 class was indispensable in developing the background necessary to show this result.

## References

- [1] Steven R Lay. *Analysis*. 4th ed. Upper Saddle River, NJ: Pearson, 2004.