

These are some notes from my Algebra I class. We use the textbook *Abstract Algebra* by Dummit and Foote, and will cover rings, groups, and modules.

## Contents

<b>PIDs, UFDs and All That</b>	<b>1</b>
Preliminaries . . . . .	1
Chinese Remainder Theorem . . . . .	2
Field of Fractions and Localization . . . . .	3
Unique Factorization Domains . . . . .	5
Euclidean Domains . . . . .	8
Unique Factorization in Polynomial Rings . . . . .	9
<b>Modules</b>	<b>14</b>
Basic Definitions . . . . .	14
Some Special $R$ -Modules . . . . .	15
Free Modules and Direct Sums . . . . .	16
Noetherian Properties . . . . .	18
A Taste of Homological Algebra . . . . .	20
<b>Groups</b>	<b>23</b>
Canonical Examples of Group Actions . . . . .	25
The Sylow Theorems . . . . .	27
Semidirect Products . . . . .	28
Conjugacy in $S_n$ and $A_n$ . . . . .	30
Solvable Groups . . . . .	35
Nilpotent Groups . . . . .	37
Free Groups . . . . .	39

## PIDs, UFDs and All That

We always assume here that  $R$  is commutative and unital.

### Preliminaries

**Definition:** If  $a_1, \dots, a_n \in R$ , then the *ideal generated by*  $a_1, \dots, a_n$  is given by

$$(a_1, \dots, a_n) := \bigcap \{I \mid a_1, \dots, a_n \in I, I \text{ is an ideal in } R\}.$$

An ideal is called *principal* if  $I = (a)$  for some  $a \in I$ . We may write  $I = a \cdot R$  in this case. A ring where every ideal is principal is called a *principal ideal domain*.

**Definition:** If  $I$  and  $J$  are ideals in  $R$ , then  $IJ$  is given by

$$IJ = \left\{ \sum_{i=1}^n x_i y_i \mid x_i \in I, y_i \in J, n \in \mathbb{N} \right\}.$$

**Theorem** (Isomorphism Theorems for Rings):

**First Isomorphism Theorem:** Let  $\varphi: R \rightarrow S$  be a ring homomorphism. Then,  $\bar{\varphi}: R/\ker(\varphi) \rightarrow \text{im}(\varphi)$  is an isomorphism given by  $\bar{\varphi}(a + \ker(\varphi)) = \varphi(a)$ .

**Second Isomorphism Theorem:** Let  $R$  be a ring,  $S \subseteq R$  a subring, and let  $I \subseteq R$  be an ideal. Then,

- (i)  $I + S$  is a subring of  $R$ ;

- (ii)  $I$  is an ideal of  $I + S$ ;
- (iii)  $I \cap S$  is an ideal of  $S$ ;
- (iv)  $S/I \cap S \cong I + S/I$ .

**Third Isomorphism Theorem:** Let  $R$  be a ring,  $I, J$  ideals of  $R$  with  $I \subseteq J$ . Then,  $J/I$  is an ideal of  $R/I$ , and we have  $(R/I)/(J/I) \cong R/J$ .

**Fourth Isomorphism Theorem:** If  $R$  is a ring and  $I$  is an ideal, then there is a one-to-one correspondence between subrings of  $R/I$  and subrings of  $R$  containing  $I$ .

**Definition:** Let  $M$  be an ideal in  $R$ .

- (i) We say  $M$  is prime if  $M \neq R$  and, for any  $ab \in M$ , we have either  $a \in M$  or  $b \in M$ .
- (ii) We say  $M$  is maximal if  $M \neq R$  and if  $M \subseteq I \subseteq R$  where  $I$  is an ideal, then either  $I = M$  or  $I = R$ .

**Theorem:** Let  $M$  be an ideal in  $R$ .

- (i)  $M$  is prime if and only if  $R/M$  is an integral domain.
- (ii)  $M$  is maximal if and only if  $R/M$  is a field.

*Proof.*

- (i) Let  $M$  be maximal, with  $a + M \in R/M$ ,  $a + M \neq 0 + M$ . Then,  $a \notin M$ , so that the ideal  $(a) + M$  strictly contains  $M$ . Therefore,  $1 + M \in (a) + M$ , meaning there is some  $r + M$  such that  $(r + M)(a + M) = 1 + M$ . Thus, an inverse exists.

Now, if  $R/M$  is a field, and  $M \subsetneq I \subseteq R$ , then  $I/M$  is an ideal of  $R/M$ , and since  $I \supsetneq M$ , we have  $I/M \neq 0 + M$ . Since  $R/M$  is a field, its only ideals are either  $0 + M$  and  $R/M$ , so  $I/M = R/M$ , meaning  $I = R$ .

- (ii) We have  $P \subseteq R$  is prime if and only if  $ab \in P$  implies  $a \in P$  or  $b \in P$ . Yet, means that  $ab + P = 0 + P$  if and only if  $a = 0 + P$  or  $b = 0 + P$ .

□

## Chinese Remainder Theorem

**Definition:** We say two ideals  $I$  and  $J$  are *coprime* if  $I + J = R$ , or that there exist  $x \in I$  and  $y \in J$  such that  $x + y = 1$ .

**Theorem** (Chinese Remainder Theorem): Let  $I_1, \dots, I_n$  be pairwise coprime ideals of  $R$ . Then, for any  $a_1, \dots, a_n \in R$ , there exists  $x \in R$  with  $x \equiv a_i$  modulo  $I_i$  for all  $i$ . In other words, there a solution to the system of congruences given by

$$\begin{aligned} x + I_1 &= a_1 + I_1 \\ x + I_2 &= a_2 + I_2 \\ &\vdots \\ x + I_n &= a_n + I_n. \end{aligned}$$

*Proof.* It suffices to construct elements  $y_1, \dots, y_n$  such that  $y_i \equiv 1$  modulo  $I_i$  and 0 otherwise. Then, we will be able to set  $x = \sum_i a_i y_i$  as our desired solution.

We construct  $y_1$  as follows. From our assumption,  $I_1 + I_j = R$  for all  $j \geq 2$ , so for each  $j \geq 2$ , there exists

$u_j \in I_1$  and  $v_j \in I_j$  such that  $u_j + v_j = 1$ . Taking the product, we find that

$$\prod_{j=2}^n (u_j + v_j) = 1$$

$$= \underbrace{v_2 \cdots v_n}_{=: y_1} + \cdots + \underbrace{u_2 \cdots u_n}_{=: x_1}.$$

We verify that  $y_1$  does the job, which we can see by the fact that  $y_1 \equiv 0$  modulo  $I_j$  for  $j \neq 1$ , as  $v_2 \cdots v_j \in I_2 \cdots I_j \subseteq I_j$  for each  $j \geq 2$ . Similarly, each summand in  $x_1$  contains at least one  $u_j$ , so  $x_1 \equiv 0$  modulo  $I_1$ .

The rest of the  $y_i$  follow analogously. □

We can restate the Chinese Remainder Theorem in a variety of ways.

**Theorem** (Chinese Remainder Theorem, Alternative Versions): Let  $I_1, \dots, I_n$  be pairwise coprime ideals.

(i) There exists a surjective homomorphism

$$\begin{aligned} \varphi: R &\rightarrow R/I_1 \times \cdots \times R/I_n \\ r &\mapsto (r + I_1, \dots, r + I_n). \end{aligned}$$

This homomorphism induces an isomorphism

$$\overline{\varphi}: R/(I_1 \cap \cdots \cap I_n) \rightarrow R/I_1 \times \cdots \times R/I_n.$$

(ii) If  $I_1, \dots, I_n$  are pairwise coprime, then

$$R/I_1 \cdots I_n \cong R/I_1 \times \cdots \times R/I_n$$

are isomorphic.

**Example:** We observe that if  $R = \mathbb{Z}$ , and  $p_1, \dots, p_r$  are distinct primes with  $\ell_1, \dots, \ell_r$  positive integers, then

$$\mathbb{Z}/p_1^{\ell_1} \cdots p_r^{\ell_r} \mathbb{Z} \cong \mathbb{Z}/p_1^{\ell_1} \mathbb{Z} \times \cdots \times \mathbb{Z}/p_r^{\ell_r} \mathbb{Z}.$$

**Example** (Polynomial Interpolation): If we let

$$p_i(x) = x - \alpha_i,$$

where  $\alpha_i \in \mathbb{F}$ , we observe that there is a surjective evaluation homomorphism

$$\text{ev}: \frac{\mathbb{F}[x]}{(p_i(x))} \rightarrow \mathbb{F},$$

given by  $f(x) \mapsto f(\alpha_i)$ . In particular, if  $\alpha_1, \dots, \alpha_r$  are distinct, then

$$\frac{\mathbb{F}[x]}{(p_1(x), \dots, p_r(x))} \cong \mathbb{F} \times \cdots \times \mathbb{F},$$

so that, for all  $\beta_1, \dots, \beta_r \in \mathbb{F}$ , there is some  $f(x) \in \mathbb{F}[x]$  such that  $f(\alpha_i) = \beta_i$  for  $i = 1, \dots, r$ .

## Field of Fractions and Localization

Given a ring  $R$ , how can we find maximal ideals in  $R$ ? More specifically, given a commutative ring  $R$  with 1, and prime ideal  $P \subseteq R$ , we want to construct a new ring  $R_P$  with unique maximal ideal  $P$ .

Toward this end, we start by reviewing a useful construction known as the field of fractions.

**Definition:** Let  $R$  be an integral domain. We define the field  $K = \text{frac}(R)$  to be the unique field with an injection

$$\begin{aligned}\iota: R &\hookrightarrow K \\ 1_R &\mapsto 1_K,\end{aligned}$$

satisfying the following universal property.

Given any embedding into a field,  $\sigma: R \hookrightarrow L$ , such that  $1_R \mapsto 1_L$ , there is a unique extension  $\tilde{\sigma}: K \rightarrow L$  such that the following diagram commutes.

$$\begin{array}{ccc} R & \xrightarrow{\iota} & K \\ & \searrow \sigma & \downarrow \tilde{\sigma} \\ & & L \end{array}$$

In order to construct  $K$ , we let  $S \subseteq R \times R$  be defined by

$$S = \{(a, b) \mid b \neq 0\}.$$

We impose an equivalence relation on  $S$  by saying  $(a, b) \sim (c, d)$  if and only if  $ad - bc = 0$ . Clearly, this relation is reflexive and symmetric. To see that it is transitive, we let  $(a, b) \sim (c, d)$ , and  $(c, d) \sim (e, f)$ , meaning  $ad - bc = 0$  and  $cf - de = 0$ . Multiplying the first equation by  $f$  and the second equation by  $b$ , then subtracting, we get  $adf - bde = 0$ , meaning  $d(af - be) = 0$ . Since  $R$  admits no zero divisors, this means that  $af - be = 0$ , so the relation is transitive.

We write  $[(a, b)] = \frac{a}{b}$  for  $K$ , with operations

$$\begin{aligned}\frac{a}{b} + \frac{c}{d} &= \frac{ad + bc}{bd} \\ \frac{a}{b} \cdot \frac{c}{d} &= \frac{ac}{bd}.\end{aligned}$$

These operations are well-defined and do satisfy the universal property. Verifying this is a pain, but it can be done.

Now, we may extend this to all unital commutative rings, not just integral domains.

**Definition:** Let  $R$  be a unital commutative ring, and let  $S \subseteq R$ . We say  $S$  is *multiplicative* if

- $1 \in S$ ;
- $0 \notin S$ ;
- for any  $x, y \in S$ ,  $xy \in S$ .

**Example:**

- (i) If  $R$  is an integral domain, then  $R \setminus \{0\}$  is multiplicative.
- (ii) If  $z \in R$  is such that  $z$  is not nilpotent, then  $S = \{z^n \mid n \geq 0\}$  is multiplicative.
- (iii) If  $P$  is a prime ideal, then  $S = R \setminus P$  is multiplicative.

We will use (iii) to construct a ring with a unique maximal ideal. First, though, we construct a ring of fractions using multiplicative sets.

**Definition:** Let  $R$  be a unital commutative ring, and let  $S \subseteq R$  be multiplicative. We construct a ring

$S^{-1}R$  by taking an equivalence relation on  $R \times S$  as follows:

$$(a, s) \sim (b, t) \Leftrightarrow \exists s' \in S \text{ such that } s'(at - bs) = 0.$$

We write

$$S^{-1}R = \{[(a, s)] \mid a \in R, s \in S\},$$

and denote

$$[(a, s)] = \frac{a}{s}.$$

This becomes a ring under the operations

$$\begin{aligned} \frac{a}{s} + \frac{b}{t} &= \frac{at + bs}{st} \\ \frac{a}{s} \cdot \frac{b}{t} &= \frac{ab}{st}. \end{aligned}$$

We call  $S^{-1}R$  the *localization of  $R$  with respect to  $S$* .

We can see some basic properties of the localization.

**Proposition:** Let  $R$  be a unital commutative ring,  $S \subseteq R$  multiplicative, and let  $S^{-1}R$  be the corresponding localization.

- The additive identity in  $S^{-1}R$  is  $\frac{0}{1}$ .
- The additive inverse of  $\frac{a}{s}$  in  $S^{-1}R$  is  $\frac{-a}{s}$ .
- For all  $a \in R$  and all  $s, s' \in S$ , we have  $\frac{as'}{ss'} = \frac{a}{s}$ .
- Every element of the form  $\frac{s}{t}$  where both  $s, t \in S$  is invertible, with corresponding inverse  $\frac{t}{s}$ .
- The map  $\iota_S: R \rightarrow S^{-1}R$  given by  $r \mapsto \frac{r}{1}$  is an injective ring homomorphism such that  $\iota_S(S) \subseteq (S^{-1}R)^\times$ , where  $(S^{-1}R)^\times$  denotes the group of invertible elements in  $S^{-1}R$ .

## Unique Factorization Domains

**Definition:** A ring  $R$  is called *Noetherian* if, for any ascending chain of ideals  $I_1 \subseteq I_2 \subseteq \dots$ , there is some index  $N$  such that for all  $m \geq N$ ,  $I_m = I_N$ .

**Proposition:** The following are equivalent:

- $R$  is Noetherian;
- every ideal in  $R$  is finitely generated.

*Proof.* Let  $R$  be Noetherian. Suppose toward contradiction that there exists  $I$  that is not finitely generated. Then,  $I$  is nonzero, so there is  $\alpha_1 \in I$  such that  $I_1 = (\alpha_1)$  is nonzero. Since  $I$  is not finitely generated,  $I \neq I_1$ , so there is  $\alpha_2 \in I \setminus I_1$ , so that  $I_2 = (\alpha_1, \alpha_2)$  is such that  $I_1 \subsetneq I_2$ . Inductively, we generate  $I_n = (\alpha_1, \dots, \alpha_n)$  such that  $I_{n-1} \subsetneq I_n$ , implying that we have a strictly ascending chain of ideals, which is a contradiction.

Suppose every ideal in  $R$  is finitely generated. Let  $I_1 \subseteq I_2 \subseteq \dots$  be an ascending chain of ideals, and set  $I = \bigcup I_n$  be their union. By assumption,  $I$  is finitely generated, so we have  $I = (\alpha_1, \dots, \alpha_N)$  for some  $\alpha_1, \dots, \alpha_N \in R$ . Yet, since  $I$  is the union of all these ideals, there is some  $M$  such that  $\alpha_1, \dots, \alpha_N \in I_M$ , meaning the chain stabilizes.  $\square$

**Corollary:** If  $R$  is a principal ideal domain, then  $R$  is Noetherian.

**Definition:** Let  $R$  be an integral domain.

- (i) Two elements  $a, b \in R$  are called *associated* if  $a = bu$  for some unit (invertible) element  $u \in R$ . Equivalently,  $a$  and  $b$  are associated if  $(a) = (b)$
- (ii) An element  $a \in R$  is called *irreducible* if
  - $a$  is not a unit element;
  - whenever  $a = bc$  for some  $b, c \in R$ , then one of  $b$  or  $c$  is a unit.
- (iii) An element  $a$  is called *prime* if  $a \neq 0$ ,  $a \notin R^\times$ , and  $(a)$  is prime. Equivalently,  $a$  is prime if, whenever  $a|bc$ , it follows that  $a|b$  or  $a|c$ , where divisibility in  $R$  is defined traditionally (i.e., there exists  $z \in R$  such that  $az = b$ ).

**Note:** Prime elements are irreducible, but not necessarily vice versa.

The question then arises: when are irreducibles prime?

**Definition:** We say  $a \in R$  with  $a \neq 0$ ,  $a \notin R^\times$  has a *unique factorization* into irreducibles if

- (i) we may write  $a = up_1 \cdots p_r$ , where  $u$  is a unit and  $p_1, \dots, p_r$  are irreducible;
- (ii) for any other such factorization

$$\begin{aligned} a &= u \prod_{i=1}^r p_i \\ &= v \prod_{j=1}^s q_j, \end{aligned}$$

where  $p_i, q_j$  are irreducible and  $u, v$  are units, we have

- $r = s$ ;
- upon permutation of factors,  $p_i$  and  $q_i$  are associated.

We call  $R$  a *unique factorization domain* if, for any  $a \in R$  with  $a \neq 0$ ,  $a \notin R^\times$ ,  $a$  has unique factorization into irreducibles.

**Proposition:** If  $R$  a Noetherian ring, then every  $a \in R$  with  $a \neq 0$  and  $a \notin R^\times$  admits a factorization into irreducibles.

*Proof.* First, we show that every such  $a$  has an irreducible factor or divisor. If  $a$  is itself irreducible, then we are done. Else, there are  $b, c \in R$  with  $a = bc$  and neither  $a$  nor  $b$  a unit. In particular, this means that  $(a) \subsetneq (b)$ . Inductively, if  $b$  is not irreducible, then we may find  $b_2, c_2$  such that  $b = b_2c_2$ , meaning that  $(b) \subsetneq (b_2)$ , and so on and so forth.

This gives a chain of ideals

$$(a) \subsetneq (b) \subsetneq (b_2) \subsetneq \cdots$$

that eventually stabilizes, meaning that there is some  $b_N$  such that  $b_N$  is irreducible.

Now, we may show that  $a$  admits a factorization. If  $a = bc$  with  $b$  irreducible (as we showed previously), then if  $c$  is not irreducible, we may take  $c = b_1c_1$  and create this same chain of ideals

$$(c) \subsetneq (c_1) \subsetneq (c_2) \subsetneq \cdots$$

using the Noetherian condition to end up at an irreducible or a unit. □

The main issue facing general Noetherian rings is that the uniqueness of the factorization may go awry.

**Example:** For instance, in the ring  $R = \mathbb{Z}[\sqrt{-5}]$ , there is not unique factorization. For instance, we may write

$$\begin{aligned} 6 &= (2)(3) \\ &= (1 + \sqrt{-5})(1 + \sqrt{-5}), \end{aligned}$$

where we may see that all of these are irreducible as follows. Define a norm on  $\mathbb{Z}[\sqrt{-5}] \subseteq \mathbb{C}$  by  $N(a + b\sqrt{-5}) = a^2 + 5b^2$ , where this norm is multiplicative as it is inherited from  $\mathbb{C}$ .

**Lemma:** If  $N$  is a norm on the ring  $R = \mathbb{Z}[\sqrt{-D}]$ , where  $D$  is a square-free positive integer, then  $u \in R$  is an invertible (or unit) element if and only if  $N(u) = 1$ .

*Proof of Lemma.* If  $v \in R$  is such that  $uv = 1$ , then  $N(uv) = N(u)N(v) = 1$ , meaning that both  $N(u)$  and  $N(v)$  are 1.

Meanwhile, if  $N(u) = 1$ , then  $1 = u\bar{u}$ , meaning that  $\bar{u} = u^{-1}$ . □

We may show that 2 is irreducible relatively quickly. Observe that if there were a factorization of  $2 = ab$  into irreducibles, then  $4 = N(a)N(b)$  would hold, with neither  $N(a)$  nor  $N(b)$  being equal to 1. This would mean that  $N(a) = 2$  for some  $a = x + y\sqrt{-5}$ , or that  $x^2 + 5y^2 = 2$ . Yet, reducing modulo 5, this implies that  $x^2 \equiv 2$  modulo 5, yet the only squares in  $\mathbb{Z}/5\mathbb{Z}$  are 1 and 4.

Given a factorization, there is a simple way to classify the uniqueness of the factorization.

**Proposition:** Let  $a \in R$  be such that  $a \neq 0$  and  $a \notin R^\times$ . If  $a$  admits a factorization

$$a = up_1 \cdots p_r,$$

with  $p_1, \dots, p_n$  prime, then this factorization is unique (up to associates).

*Proof.* Suppose  $a$  admits another factorization,

$$a = vq_1 \cdots q_s,$$

where  $q_1, \dots, q_s$  are irreducible and  $v$  is a unit. Then, we have

$$up_1 \cdots p_r = vq_1 \cdots q_s,$$

meaning that  $p_1$  divides  $vq_1 \cdots q_s$ . Since  $p_1$  is prime,  $p_1 | q_j$  for some  $j$ , meaning that  $q_j = v_1 p_1$  for some  $v_1 \in R$ . Yet, since  $q_j$  is irreducible, it follows that  $v_1$  is a unit. By permuting elements, we may say that  $p_1$  and  $q_1$  are associated, so we have

$$up_1 \cdots p_r = vv_1 p_1 q_2 \cdots q_s.$$

Now, since  $R$  is a domain, it admits the cancellation property, so we may then write

$$up_2 \cdots p_r = vv_1 q_2 \cdots q_s.$$

Proceeding in this fashion, we observe first that  $r \leq s$ , as else, we would have  $p_i$  dividing a unit for  $R$ , which is not allowed. Thus, we find

$$u = vv_1 \cdots v_r q_{r+1} \cdots q_s.$$

Similarly, this means there cannot be any more  $q_j$ , or else the  $q_j$  would be a unit. Thus, these are the same factorizations (up to associates). □

**Theorem:** If a domain  $R$  is a principal ideal domain, then  $R$  is a unique factorization domain.

*Proof.* First, we show that if  $a \in R$  is irreducible, then  $a$  is prime.

Observe that  $(a)$  is then contained in a maximal ideal  $M$ , where  $M = (p)$  for some  $p \in R$  with  $p$  not a

unit. Since  $M$  is maximal,  $M$  is prime, so that  $p$  is prime, and  $(a) \subseteq (p)$ . Observe then that  $a = pu$  for some  $u \in R$ ; since  $a$  is irreducible and  $p$  is not a unit, it must be the case that  $u$  is a unit. Thus,  $(a) = (p)$ , so that  $a$  is prime.

Now, since  $R$  is a principal ideal domain, every element in  $R$  admits a factorization into irreducibles, and all irreducibles are prime. Therefore, the factorization is unique by the above lemma.  $\square$

## Euclidean Domains

**Definition:** An integral domain  $R$  is called a *Euclidean Domain* if there exists  $N: R \setminus \{0\} \rightarrow \mathbb{Z}_{\geq 0}$  such that for all  $a, b \in R$ , with  $b \neq 0$ , there exist  $q, r \in R$  such that

- $a = qb + r$ ;
- either  $r = 0$  or  $N(r) < N(b)$ .

**Example:**

- Any field admits the vacuous norm,  $N(k) = 0$  for all  $k \in F \setminus \{0\}$ .
- The ring  $R = \mathbb{Z}$  is Euclidean with the norm  $N(n) = |n|$ .
- The ring  $R = \mathbb{F}[x]$ , where  $\mathbb{F}$  is a field, is Euclidean with norm  $N: \mathbb{F}[x] \setminus \{0\} \rightarrow \mathbb{N}$  given by  $N(f) = \deg(f)$ .

**Theorem:** If  $R$  is Euclidean, then  $R$  is a principal ideal domain.

*Proof.* Let  $I \subseteq R$  be an ideal. If  $I = \{0\}$ , then  $I$  is principal and we are done.

Else, suppose  $I \neq 0$ . There exists  $\alpha \in I$  with  $\alpha \neq 0$ , so that  $N(\alpha)$  is well-defined. Let  $b \in I$  be such that  $N(b)$  is minimal for all possible elements of  $I$ .

We claim that  $I = (b)$ . Let  $a \in I$  be arbitrary, and perform Euclidean division on  $a$  by  $b$ , yielding

$$a = qb + r,$$

where  $r = 0$  or  $N(r) < N(b)$ .

If  $r \neq 0$ , then  $N(r) < N(b)$ , but  $r = a - bq \in I$ , which would contradict minimality of  $N(b)$ , so that  $r = 0$ , and thus  $a = bq \in (b)$ .  $\square$

**Theorem:** The Gaussian integers,  $\mathbb{Z}[i]$ , are Euclidean with norm

$$N(a + bi) = a^2 + b^2.$$

*Proof.* Observe that  $N$  is multiplicative. If we let  $\alpha = a + bi$  and  $\beta = c + di$  with  $\alpha, \beta \neq 0$ , we want to show that there exist  $\gamma$  and  $\delta$  such that  $\alpha = \beta\gamma + \delta$  and  $\delta = 0$  or  $N(\delta) < N(\beta)$ .

Consider  $\frac{\alpha}{\beta} \in \mathbb{C}$ , so that

$$\begin{aligned} \frac{\alpha}{\beta} &= \frac{(a + bi)(c - di)}{c^2 + d^2} \\ &= \frac{(a + bi)(c - di)}{N(\beta)} \\ &=: x + yi, \end{aligned}$$

so that  $\frac{\alpha}{\beta} \in \mathbb{Q}[i]$ .

Now, we can find  $x_0, y_0 \in \mathbb{Z}$  such that  $|x - x_0| \leq \frac{1}{2}$  and  $|y - y_0| \leq \frac{1}{2}$ . Setting  $\delta = x_0 + y_0 i$ , we have that  $\delta = \alpha - \beta\gamma \in \mathbb{Z}[i]$ . We claim that if  $\delta \neq 0$ , then  $N(\delta) < N(\beta)$ .



Observe that since  $N$  is multiplicative, this condition is equivalent to  $N\left(\frac{\delta}{\beta}\right) < 1$ . We observe that

$$\begin{aligned} N\left(\frac{\delta}{\beta}\right) &= N\left(\frac{\alpha - \beta\gamma}{\beta}\right) \\ &= N\left(\frac{\alpha}{\beta} - \gamma\right) \\ &= (x - x_0)^2 + (y - y_0)^2 \\ &\leq \frac{1}{2} \\ &< 1. \end{aligned}$$

□

**Remark:** While the remainder in Euclidean division for  $\mathbb{Z}$  and  $\mathbb{F}[x]$  is unique, this is not the case for general Euclidean domains. For instance, if we want to divide  $a = 1 + i$  by  $b = 2$  in  $\mathbb{Z}[i]$  with our previously specified norm, we find that

$$\begin{aligned} 1 + i &= 2 \cdot 0 + (1 + i) \\ &= 2 \cdot 1 + (-1 + i), \end{aligned}$$

both of which satisfy the conditions for Euclidean division.

Now, in any PID (really, any UFD), we can talk about a greatest common divisor. In a principal ideal domain, the GCD for  $a, b \in R$  is given by the unique (up to associates) element  $d$  such that

$$(a, b) = (d).$$

Meanwhile, greatest common divisors in a UFD are slightly more complicated. If we have two elements  $a, b \in R$  with prime factorizations

$$\begin{aligned} a &= up_1^{v_1} p_2^{v_2} \cdots p_n^{v_n} \\ b &= vp_1^{w_1} p_2^{w_2} \cdots p_n^{w_n}, \end{aligned}$$

then the greatest common divisor is given by

$$\gcd(a, b) = \prod_{i=1}^n p_i^{\min(v_i, w_i)}.$$

This is defined up to associates, similar to how the factorization of any element is defined up to associates.

## Unique Factorization in Polynomial Rings

Our goal is to prove that if  $R$  is a UFD, then  $R[x]$  is a UFD.

We do this by first discussing irreducibility in  $R[x]$ , including a full characterization of irreducible elements.

**Definition:** Assume  $R$  is a unique factorization domain, and let  $0 \neq f(x) \in R[x]$ . Writing

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

we define the *content* of  $f$ , written  $c(f)$ , to be

$$c(f) = \gcd(a_0, a_1, \dots, a_n).$$

**Proposition** (Gauss's Lemma): Let  $R$  be a UFD, and let  $f(x), g(x) \in R[x]$  be nonzero polynomials. Then,

$$c(fg) = c(f) c(g).$$

*Proof.* For any nonzero polynomial  $h \in R[x]$ , we may write

$$h(x) = c(h)z(x),$$

where  $c(z) = 1$ , simply by factoring. Thus, writing

$$\begin{aligned} f(x) &= c(f)u(x) \\ g(x) &= c(g)v(x), \end{aligned}$$

where  $c(u) = c(v) = 1$ , hence

$$\begin{aligned} c(fg) &= c(c(f) c(g)uv) \\ &= c(f) c(g) c(uv). \end{aligned}$$

We want to show that  $c(u(x)v(x)) = 1$  (up to associates).

Suppose not. Since  $c(uv)$  is nonzero and (assumed to be) not a unit, we may find a prime  $p$  such that  $p \mid c(uv)$ . That is, we may find  $p$  such that  $p$  divides all coefficients of  $u(x)v(x)$ .

Consider now the reduction homomorphism

$$\pi: R[x] \rightarrow (R/(p))[x],$$

where we reduce all coefficients modulo  $(p)$ . Since  $p$  is prime,  $(p)$  is prime, so that  $R/(p)$  is an integral domain, meaning that  $(R/(p))[x]$  is an integral domain.

Since  $c(u) = c(v) = 1$ , it follows that  $\pi(u(x)) \neq 0$  and  $\pi(v(x)) \neq 0$ , as at least one coefficient in  $u(x)$  or  $v(x)$  is not divisible by  $p$ . Thus, in  $(R/(p))[x]$  is a domain, it follows that  $\pi(u(x))\pi(v(x)) \neq 0$ . Yet, since  $\pi$  is a homomorphism, it follows that  $0 = \pi(u(x)v(x)) = \pi(u(x))\pi(v(x))$ , since we assumed that  $p$  divides all the coefficients of  $u(x)v(x)$ .  $\square$

**Corollary** (Gauss's Lemma, Redux): Let  $R$  be a UFD, and let  $F = \text{frac}(R)$ . Let  $f(x) \in R[x]$ , and assume  $f(x)$  is reducible in  $F[x]$ . Then,  $f(x)$  is reducible in  $R[x]$ .

*Proof.* Let  $f(x)$  be reducible in  $F[x]$ , so that  $f(x) = g(x)h(x)$ , where  $g(x)$  and  $h(x)$  are nonconstant polynomials in  $F[x]$ .

By factoring, we have

$$\begin{aligned} g(x) &= \frac{a}{b}u(x) \\ h(x) &= \frac{c}{d}v(x), \end{aligned}$$

where  $a, b, c, d \in R \setminus \{0\}$ ,  $u(x), v(x) \in R[x]$ , and  $c(u) = c(v) = 1$ .

Substituting this information into the expression for  $f(x)$ , we have

$$\begin{aligned} f(x) &= \frac{ac}{bd}u(x)v(x) \\ bdf(x) &= acu(x)v(x), \end{aligned}$$

so that

$$bdc(f) = ac c(u) c(v).$$

meaning

$$bdc(f) = ac.$$

In particular, this means that  $\frac{ac}{bd}$  is a valid representative for  $c(f)$ , so that  $\frac{ac}{bd} \in R$ . Therefore,

$$f(x) = \left(\frac{ac}{bd}u(x)\right)v(x),$$

both nonconstant and in  $R[x]$ , meaning  $f(x)$  has a nontrivial factorization in  $R[x]$ , and thus  $f$  is reducible.  $\square$

**Corollary** (Classification of Irreducibles): Let  $R$  be a UFD, let  $F = \text{frac}(R)$ , and let  $f(x) \neq 0 \in R[x]$ .

- (i) If  $f(x)$  is constant, then  $f$  is irreducible in  $R[x]$  if and only if  $f(x)$  is irreducible in  $R$ .
- (ii) If  $f(x)$  is not constant, then  $f$  is irreducible in  $R[x]$  if and only if  $c(f) = 1$  and  $f(x)$  is irreducible in  $F[x]$ .

*Proof.*

- (i) Observe that  $R[x]$  and  $R$  have the same units (since  $R$  is an integral domain, and so admits no nilpotent elements), meaning that the product of two nonzero polynomials is a constant if and only if the polynomials themselves are constant.
- (ii) Let  $f$  be nonconstant. If  $f$  is irreducible in  $R[x]$ , then we may write

$$f(x) = c(f)u(x),$$

where  $u(x)$  is nonconstant and has  $c(u) = 1$ . Yet, since  $f$  is irreducible, it also follows that  $c(f) = 1$ . Additionally,  $f$  is irreducible in  $F[x]$  by the contrapositive of Gauss's Lemma.

If  $f$  is irreducible in  $F[x]$ , and has content 1, then for any factorization

$$f(x) = g(x)h(x),$$

where  $g(x), h(x) \in F[x]$ , either  $g$  or  $h$  must be a constant. Now, since  $f$  is contained in  $R[x]$ , we may take a common denominator to yield

$$f(x) = au(x),$$

where  $u(x)$  is nonconstant and has content 1, with  $a \in R$ . Since  $f$  has content 1, it follows that  $a$  is a unit element, meaning that any factorization of  $f$  must contain a unit, so that  $f$  is irreducible in  $R[x]$ .  $\square$

**Theorem:** If  $R$  is a UFD, then  $R[x]$  is a UFD.

*Proof.* Let  $F = \text{frac}(R)$ , and let  $f(x) \in R[x]$  be a nonzero, non-unit element. If  $f(x) \in R$ , then  $f$  is a product of irreducibles in  $R$  by part (i) of the classification, meaning the product is automatically unique up to permutation and associates as  $R$  is a UFD.

Now, if  $f$  is nonconstant, then  $f(x) \in F[x]$  is nonzero and non-unit, as the units in  $F[x]$  are the elements of  $F$ . Since  $F[x]$  is a principal ideal domain (as  $F[x]$  is a Euclidean domain, following from the division algorithm),  $F[x]$  is a UFD, so we may write

$$f(x) = \prod_{i=1}^n g_i(x),$$

where the  $g_i(x)$  are irreducible in  $F[x]$ . Writing

$$g_i(x) = \frac{a_i}{b_i}u_i(x),$$

where the  $u_i(x) \in R[x]$  with  $c(u_i) = 1$  for each  $i$ , we have

$$\prod_{i=1}^n \frac{a_i}{b_i} \in R,$$

as  $f(x) \in R[x]$ , so we may write

$$f(x) = \prod_{i=1}^n \frac{a_i}{b_i} \prod_{i=1}^n u_i(x).$$

Each of the  $u_i(x)$  are irreducible in  $R[x]$  by the classification, and the product  $\prod_{i=1}^n \frac{a_i}{b_i} \in R$  is either a unit or a product of irreducibles. This gives the existence of such a factorization for  $f$ .

To see uniqueness, if

$$\begin{aligned} f(x) &= \left( \prod_{i=1}^k a_i \right) \left( \prod_{i=1}^m p_i(x) \right) \\ &= \left( \prod_{j=1}^\ell b_j \right) \left( \prod_{j=1}^m q_j(x) \right) \end{aligned}$$

are factorizations where  $a_i, b_j$  are irreducible in  $R$ , and  $p_i, q_j$  are nonconstant and irreducible with content 1, then we may take the content of both sides, yielding

$$\begin{aligned} c \left( \left( \prod_{i=1}^k a_i \right) \left( \prod_{i=1}^m p_i(x) \right) \right) &= \prod_{i=1}^k a_i \\ c \left( \left( \prod_{j=1}^\ell b_j \right) \left( \prod_{j=1}^m q_j(x) \right) \right) &= \prod_{j=1}^\ell b_j. \end{aligned}$$

Since contents are only well-defined up to associates, the most we can say is that

$$\prod_{i=1}^k a_i = u \prod_{j=1}^\ell b_j,$$

where  $u \in R^\times$ . Since there is at least one  $q_j$ , we may replace  $q_1$  by  $uq_1$ , then divide, so that we find

$$\prod_{i=1}^k a_i = \prod_{j=1}^\ell b_j.$$

Since both of these are products of irreducibles in  $R$ , it follows that  $k = \ell$  and, after permutation of factors,  $b_i = u_i a_i$  for some  $u_i \in R^\times$ . Additionally, we also have the equality

$$\prod_{i=1}^m p_i = \prod_{j=1}^n q_j.$$

Since all of these factors are irreducible in  $F[x]$ , and  $F[x]$  is a PID, we find that  $n = m$  and, upon permutation of factors, we have  $q_i(x) = \gamma_i p_i(x)$  for some  $\gamma_i \in F \setminus \{0\}$ . Write

$$\gamma_i = \frac{c_i}{d_i},$$

where  $c_i, d_i \in R \setminus \{0\}$ , so that

$$d_i q_i(x) = c_i p_i(x).$$

Taking the content of both sides, we then get that  $v_i d_i = c_i$  for some  $v_i \in R^\times$ , so that  $\gamma_i = v_i \in R^\times$ , meaning that  $p_i$  and  $q_i$  are associates in  $R[x]$ .  $\square$

Unique factorization in polynomial rings having the rigidity laid out in the classification theorem makes for very useful criteria to understand irreducibility.

**Theorem** (Eisenstein's Criterion): Let  $R$  be a UFD, and let  $p \in R$  be a prime element. If we write  $f(x) \in R[x]$  as

$$f(x) = \sum_{i=0}^n a_i x^i,$$

then if

- $a_0 \neq 0$ ;
- $p \nmid a_n$ ;
- $p \mid a_i$  for  $0 \leq i \leq n-1$ ;
- and  $p^2 \nmid a_0$ ,

then  $f(x)$  is irreducible in  $F[x]$ . If, in addition,  $c(f) = 1$ , then  $f$  is irreducible in  $R[x]$ .

**Remark:** This is the more general formulation of the case when  $R = \mathbb{Z}$  and  $f$  is monic that we see in undergrad abstract algebra.

*Proof.* Suppose toward contradiction that  $f$  is reducible in  $F[x]$ , where we may write

$$f(x) = g(x)h(x)$$

with  $g(x), h(x) \in R[x]$  nonconstant as in the proof of Gauss's Lemma. The reduction map  $\pi: R[x] \rightarrow (R/(p))[x]$  is a homomorphism, so that

$$\bar{f}(x) = \bar{g}(x)\bar{h}(x).$$

Thus, by our assumptions, we have

$$\bar{f}(x) = \bar{a}_n x^n,$$

with  $\bar{a}_n \neq \bar{0}$ . Since the degree of  $\bar{f}$  remains the same upon reduction, it follows that  $\bar{g}$  and  $\bar{h}$  have the same degrees as they had originally.

Observe that in a domain, the product of the highest-degree terms is the highest-degree term of the product, and similarly for the lowest-degree terms. Therefore, we must have  $\bar{g}(x)$  and  $\bar{h}(x)$  are monomials, as their product is a monomial. Writing

$$\begin{aligned}\bar{g}(x) &= \beta x^k \\ \bar{h}(x) &= \gamma x^\ell,\end{aligned}$$

with  $\gamma, \beta \in R$  and  $k = \deg(g)$ ,  $\ell = \deg(h)$ , we then get

$$\begin{aligned}g(x) &= bx^k + pu(x) \\ h(x) &= cx^\ell + pv(x),\end{aligned}$$

where  $k, \ell > 0$  and  $u(x), v(x) \in R[x]$ . Then,

$$f(x) = (bx^k + pu(x))(cx^\ell + pv(x)),$$

whence the constant term of this product is divisible by  $p^2$ . □

## Modules

For this section, a ring  $R$  may not be commutative nor unital.

### Basic Definitions

**Definition:** Let  $R$  be a ring. A *left  $R$ -module* is a set  $M$  with operations

$$\begin{aligned} +: M \times M &\rightarrow M \\ (m, n) &\mapsto m + n \\ \cdot: R \times M &\rightarrow M \\ (r, m) &\mapsto r \cdot m, \end{aligned}$$

satisfying the following axioms:

- (M0)  $(M, +)$  is an abelian group;
- (M1)  $(r + s) \cdot m = r \cdot m + s \cdot m$  for all  $r, s \in R$  and  $m \in M$ ;
- (M2)  $(rs) \cdot m = r \cdot (s \cdot m)$  for all  $r, s \in R$  and  $m \in M$ ;
- (M3)  $r \cdot (m + n) = r \cdot m + r \cdot n$  for all  $r \in R$  and  $m, n \in M$ ;
- (M4) if  $R$  is unital, then  $1 \cdot m = m$  for all  $m \in M$ .

A *submodule*  $N \leq M$  of an  $R$ -module  $M$  is an abelian subgroup such that  $r \cdot n \in N$  for all  $r \in R$  and  $n \in N$ .

**Definition:** If  $N \leq M$  is a submodule, then the *quotient module*  $M/N$  is formed by taking equivalence classes of the form  $m + N$ , where  $m + N = k + N$  if  $m - k \in N$ .

**Definition:** An  $R$ -module homomorphism between  $M$  and  $N$  is a map  $\varphi: M \rightarrow N$  such that  $\varphi$  is  $R$ -linear, in the sense that

$$\varphi(r \cdot m + s \cdot k) = r \cdot \varphi(m) + s \cdot \varphi(k).$$

The set of all homomorphisms between  $R$ -modules  $M$  and  $N$  is denoted  $\text{hom}_R(M, N)$ , and forms an  $R$ -module itself under pointwise operations.

The set of all  $R$ -module *endomorphisms* is denoted

$$\text{end}_R(M) := \text{hom}_R(M, M),$$

and forms a ring under pointwise operations and composition.

The space of  $R$ -module *automorphisms* is denoted

$$\text{aut}_R(M) := (\text{end}_R(M))^\times.$$

Modules admit the most “natural” form of the isomorphism theorems, as we only need to concern ourselves with submodules, rather than encountering issues like normal subgroups or ideals.

**Theorem** (Isomorphism Theorems for Modules):

**First Isomorphism Theorem:** If  $\varphi: M \rightarrow N$  is a homomorphism of  $R$ -modules, there is an induced isomorphism

$$\bar{\varphi}: M/\ker(\varphi) \rightarrow \text{im}(\varphi),$$

given by  $\bar{\varphi}(m + \ker(\varphi)) = \varphi(m)$ .

**Second Isomorphism Theorem:** If  $A$  and  $B$  are submodules of  $M$ , then there is an isomorphism

$$\frac{A+B}{A} \cong \frac{A}{A \cap B},$$

where

$$A+B = \{m+n \mid m \in A, n \in B\}.$$

**Third Isomorphism Theorem:** If  $A, B \leq M$  are submodules with  $A \subseteq B$ , then there is an isomorphism

$$M/B \cong \frac{M/A}{B/A}.$$

**Fourth Isomorphism Theorem:** If  $B \leq M$  is a submodule, then there is a one to one correspondence between the set of submodules of  $M/B$  and the set of submodules of  $M$  containing  $B$ .

## Some Special $R$ -Modules

There are three special cases of  $R$ -modules that we will discuss here. The first one is pretty straightforward, while the other two are a bit more complex and will enable us to understand some particularly deep results later down the line.

**Example:** If  $F$  is a field, then the  $F$ -modules are precisely the vector spaces over  $F$ . This is because  $F$ -vector spaces and  $F$ -modules have the exact same axioms.

**Example:** We claim that there is a one to one correspondence between  $\mathbb{Z}$ -modules and abelian groups.

One direction follows from applying a “forgetful functor” on  $M$ , taking  $M \mapsto (M, +)$ , simply discarding the  $\mathbb{Z}$ -module structure of  $M$ . In fact, this can apply to all  $R$ -modules.

In the reverse direction, if  $(M, +)$  is an abelian group, then we can specify a compatible action by  $\mathbb{Z}$  onto  $M$  by taking

$$n \cdot a = \begin{cases} \underbrace{a + \cdots + a}_{n \text{ times}} & n > 0 \\ 0 & n = 0 \\ \underbrace{-a - \cdots - a}_{-n \text{ times}} & n < 0 \end{cases}.$$

The last example is the most intriguing. In fact, as we will see towards the end, it ties directly to the Jordan Canonical Form, as we will see once we discuss the structure of finitely generated ideals over a principal ideal domain.

**Example:** We want to understand the  $F[x]$  modules, where  $F$  is a field.

Now, first, observe that since constants are elements of  $F[x]$ , it immediately follows that if  $V$  is a  $F[x]$  module, then  $V$  admits a compatible structure with respect to  $F$ , meaning that  $V$  is in fact a vector space.

Now, observe that the action of  $p(x) \in F[x]$  on  $v \in V$  is fully determined by  $x$ , as

$$x^n \cdot v = x \cdot (x \cdot (\cdots x \cdot v)).$$

If we consider a single linear transformation  $T: V \rightarrow V$ , then by defining  $T^n = T \circ \cdots \circ T$ , we observe that for any  $v \in V$ , the map

$$p(T)(v) = (a_n T^n + a_{n-1} T^{n-1} + \cdots + a_1 T + a_0)v$$

is an action on  $v \in V$ ; we may then consider the pair  $(V, T)$  to be the corresponding  $F[x]$ -module. The reverse direction follows from defining  $T: V \rightarrow V$  by  $Tv = x \cdot v$ .

Observe then that the  $F[x]$ -submodules of a  $F[x]$ -module  $V$  are precisely the  $T$ -invariant subspaces of  $V$ .

## Free Modules and Direct Sums

**Definition:** Let  $R$  be a ring,  $M$  an  $R$ -module,  $X \subseteq M$ . Then, we call

$$R \cdot X = \{r \cdot x \mid r \in R, x \in X\}$$

the submodule *generated by*  $X$ . We also write  $\langle X \rangle$ .

**Definition:**

- We say  $X \subseteq M$  is  *$R$ -linearly independent* if

$$a_1 \cdot x_1 + \cdots + a_n \cdot x_n = 0$$

for any  $x_1, \dots, x_n \in X$  and  $a_1, \dots, a_n \in R$  implies that  $a_1, \dots, a_n = 0$ .

- A subset  $X$  of  $M$  is called an  *$R$ -basis* if  $X$  is  $R$ -linearly independent and  $\langle X \rangle = M$ .
- We say  $M$  is a *free  $R$ -module* if  $M$  admits a basis.

**Theorem:** Every  $\mathbb{F}$ -vector space  $V$  has a basis. Furthermore, the following hold:

- if  $X$  is a generating set for  $V$ , then  $X$  contains a basis;
- if  $X$  is a linearly independent subset of  $V$ , then  $X$  can be extended to a basis.

**Example:** This does not always hold if we are not dealing with vector spaces. For instance,  $\mathbb{Z}$  is a free  $\mathbb{Z}$ -module, but  $\{2\}$  is a  $\mathbb{Z}$ -linearly independent subset that cannot be extended to a basis for  $\mathbb{Z}$ . This follows from the fact that  $\{2, n\}$  for any  $n \neq 2$  is  $\mathbb{Z}$ -dependent.

Similarly,  $\{2, 3\}$  is a generating set for  $\mathbb{Z}$  as  $\gcd(2, 3) = 1$ , yet  $X$  does not contain any  $\mathbb{Z}$ -bases.

**Definition** (External Direct Sum):

- Let  $M_1, \dots, M_r$  be  $R$ -modules. The *external direct sum*  $M_1 \oplus \cdots \oplus M_n$  is a module with coordinate-wise operations consisting of elements  $(m_1, \dots, m_r)$  with  $m_i \in M_i$ .
- If  $\{M_i\}_{i \in I}$  is an indexed family of  $R$ -modules, then the external direct sum of  $\{M_i\}_{i \in I}$  is defined as

$$\bigoplus_{i \in I} M_i := \left\{ f: I \rightarrow \prod_{i \in I} M_i \mid f(i) \in M_i, f \text{ is finitely supported} \right\}.$$

**Theorem:** Let  $M$  be a free  $R$ -module,  $\Sigma$  a cardinal number. The following are equivalent

- $M$  has a basis with cardinality  $\Sigma$ ;
- $M \cong \bigoplus_{i \in \Sigma} R$  as  $R$ -modules.

*Proof.* Let  $M$  have an  $R$ -basis indexed by  $\Sigma$ , written  $\{m_i\}_{i \in \Sigma}$ . Then, for every  $y \in M$ , we may write

$$y = \sum_{i \in \Sigma} r_i \cdot m_i,$$

where  $r_i = 0$  for all but finitely many such  $i \in \Sigma$ .



Now, consider the map

$$\begin{aligned}\varphi: M &\rightarrow \bigoplus_{i \in \Sigma} R \\ \sum_{i \in \Sigma} r_i \cdot m_i &\mapsto \{f_y: \Sigma \rightarrow R \mid f_y(i) = r_i\}.\end{aligned}$$

Since the expression is unique, it follows that  $\varphi$  is well-defined, and is an  $R$ -module homomorphism that is injective by the definition of a basis. Furthermore, we can define an inverse for  $\varphi$  by defining

$$\begin{aligned}\psi: \bigoplus_{i \in \Sigma} R &\rightarrow M \\ f &\mapsto \sum_{i \in \Sigma} f(i) \cdot m_i.\end{aligned}$$

Therefore,  $M \cong \bigoplus_{i \in \Sigma} R$ .

Now, if  $M \cong \bigoplus_{i \in \Sigma} R$ , then letting

$$X = \{e_i \mid e_i: \Sigma \rightarrow R, e_i(i) = 1, e_i(j) = 0 \text{ for all } j \neq i\},$$

we claim that  $X$  is an  $R$ -basis of  $\bigoplus_{i \in \Sigma} R$ . Toward this end, we only need to verify that  $X$  spans  $R$ . If  $f \in \bigoplus_{i \in \Sigma} R$ , then  $f$  is a finitely supported function from  $\Sigma$  to  $R$ , so we may write

$$f = \sum_{k=1}^n f(i_k) \cdot e_{i_k}.$$

□

**Corollary:** If  $M$  and  $N$  are free modules with bases of the same cardinality, then  $M \cong N$ .

We now turn our focus towards other ways to build up a large family of modules.

**Definition:** If  $\{M_i\}_{i \in I}$  is a collection of  $R$ -modules, then the *direct product*,

$$M = \prod_{i \in I} M_i$$

is the Cartesian product of the  $M_i$  with coordinatewise operations.

**Theorem** (Universal Property of Products): Let  $\{M_i\}_{i \in I}$  be a family of  $R$ -modules, and let

$$M = \prod_{i \in I} M_i$$

be the direct product.

(i) We have that  $M$  admits a family of projection homomorphisms

$$\begin{aligned}\pi_j: M &\rightarrow M_j \\ (m_i)_{i \in I} &\mapsto m_j.\end{aligned}$$

(ii) Given an  $R$ -module  $N$  with  $R$ -module homomorphisms  $f_j: N \rightarrow M_j$ , there exists a unique  $R$ -module homomorphism  $f: N \rightarrow M$  such that  $\pi_j \circ f = f_j$ .

$$\begin{array}{ccc}
 N & \xrightarrow{f} & \prod_{i \in I} M_i \\
 & \searrow f_j & \downarrow \pi_j \\
 & & M_j
 \end{array}$$

**Theorem** (Universal Property of Direct Sum): Let  $\{M_i\}_{i \in I}$  be a family of  $R$ -modules, and let  $L = \bigoplus_{i \in I} M_i$  be the direct sum.

Then, for an  $R$ -module  $N$  and a family of  $R$ -module homomorphisms  $g_j: M_j \rightarrow N$ , there exists a unique  $R$ -module homomorphism  $g: L \rightarrow N$  such that  $g \circ \iota_j = g_j$ , where  $\iota_j$  is the inclusion of  $M_j$  into  $L$ , given by  $m_j \mapsto (0, 0, \dots, m_j, 0, 0, \dots)$ .

$$\begin{array}{ccc}
 M_j & \xrightarrow{f} & \bigoplus_{i \in I} M_i \\
 & \searrow f_j & \downarrow \pi_j \\
 & & N
 \end{array}$$

**Proposition:** Let  $\{M_i\}_{i \in I}$  be a family of  $R$ -submodules of a fixed  $R$ -module  $M$ . The following are equivalent:

- (i) the sum  $\sum_{i \in I} M_i$  is a direct sum;
- (ii) for all  $j \in I$  and all  $i_1, \dots, i_n \in I \setminus \{j\}$ ,

$$M_j \cap \left( \sum_{k=1}^n M_{i_k} \right) = \{0\}.$$

**Theorem** (Universal Property of Free Modules): Let  $M$  be a free  $R$ -module with basis  $X$ . Then, for any  $R$ -module  $N$  and set map  $f: X \rightarrow N$ , there exists a unique  $R$ -module homomorphism  $\tilde{f}: M \rightarrow N$  such that  $\tilde{f}|_X = f$ .

$$\begin{array}{ccc}
 M & & \\
 \uparrow \iota & \searrow \tilde{f} & \\
 X & \xrightarrow{f} & N
 \end{array}$$

**Corollary:** Every  $R$ -module is a quotient of a free module.

*Proof.* Let  $M$  be an  $R$ -module, and let  $F = R[M]$  be the free module generated by the elements of  $M$ . Then, it follows that  $\text{id}: M \rightarrow M$  is a bijective set map, which then extends to a surjective module homomorphism  $q: F \rightarrow M$ . By the first isomorphism theorem, it follows that  $M \cong F/\ker(q)$ .  $\square$

**Definition:** We call the module  $\ker(q)$  the *module of relations* for  $M$ . Generally speaking, we seek to understand the minimal such module.

## Noetherian Properties

For this subsection, we fix a unital, commutative ring  $R$ . Recall that  $R$  is Noetherian if and only if every ascending chain of ideals stabilizes, if and only if every ideal is finitely generated.

We can give a similar description for Noetherian *modules*. We will use this in the proof of a much-celebrated theorem related to polynomial rings.

**Proposition:** Let  $M$  be an  $R$ -module. The following are equivalent:

- (i) Every submodule of  $M$  is finitely generated;

- (ii) every ascending chain of submodules stabilizes;
- (iii) every nonempty set of submodules has a maximal element.

**Proposition:** Let  $M$  be a Noetherian  $R$ -module, with  $N \leq M$ . Then, both  $N$  and  $M/N$  are Noetherian.

*Proof.* Let  $N' \leq N$ ; then,  $N' \leq M$ , whence  $N'$  is finitely generated, so  $N$  is Noetherian.

Meanwhile, recall that all  $R$ -submodules of  $M/N$  correspond to modules of the form  $N'/N$ , where  $N \leq N' \leq M$ , by the Fourth Isomorphism Theorem. Therefore,  $N'$  is finitely generated, so  $N'/N$  is finitely generated.  $\square$

**Proposition:** Suppose  $N \leq M$  and  $M/N$  are Noetherian. Then, so too is  $M$ .

*Proof.* Let  $K \leq M$  be an  $R$ -submodule. We will show that  $K$  is finitely generated.

Consider  $K \cap N \leq N$ . Since  $N$  is Noetherian,  $K \cap N$  is finitely generated, so it admits generators  $x_1, \dots, x_n$ . Similarly, since  $\frac{K+N}{N} \leq M/N$  is a submodule of the Noetherian module  $M/N$ , so there are generators  $y_1 + N, \dots, y_m + N$  of  $\frac{K+N}{N}$ .

In particular, since each of the  $y_i$  can be written as  $k_i + n_i$ , with  $y_i + N = k_i + N$ , we may say without loss of generality that each of the  $y_i$  are elements of  $K$ .

We claim now that  $\langle x_1, \dots, x_n, y_1, \dots, y_m \rangle = K$ . We call this generating set  $X$ . In particular, we observe already that  $\langle X \rangle \leq K$ , so we only need to show the other inclusion.

Let  $z \in K$ . Then,  $z \in K + N$ , meaning  $z + N \in \frac{K+N}{N}$ . Therefore, we have  $r_1, \dots, r_m$  such that  $z + N = r_1 \cdot y_1 + N + \dots + r_m \cdot y_m + N$ . In particular, this means that  $(z - \sum_{i=1}^m r_i \cdot y_i) \in N \cap K$ . Since  $\langle x_1, \dots, x_n \rangle = K \cap N$ , we have  $s_1, \dots, s_n \in R$  such that

$$z - (r_1 \cdot y_1 + \dots + r_m \cdot y_m) = s_1 \cdot x_1 + \dots + s_n \cdot x_n.$$

Thus,  $\langle X \rangle = K$ .  $\square$

**Theorem:** If  $R$  is a Noetherian ring, then any finitely generated  $R$ -module is Noetherian.

*Proof.* We start by proving that every finitely generated free  $R$ -module is Noetherian. Then, if  $M$  is an arbitrary finitely generated module, then  $M$  is a quotient of a finitely generated free module, and the Noetherian property is inherited under quotients.

We prove by induction on the rank. If  $F$  is free of rank  $n$ , then  $F \cong R^n$ . If  $n = 1$ , then  $F \cong R$  as  $R$ -modules. Since the submodules of  $R$  are exactly the ideals,  $F$  is Noetherian as an  $R$ -module as every ideal of  $R$  is finitely generated.

Inductively, if  $F \cong R^n$ , then the submodule  $R \cong R \times \{0\}^{n-1}$  is Noetherian, and

$$R^{n-1} \cong \frac{R^n}{R \times \{0\}^{n-1}}$$

is Noetherian by the inductive hypothesis. Thus,  $R^n$  is Noetherian by the previous proposition.  $\square$

**Theorem** (Hilbert Basis Theorem): If  $R$  is a Noetherian ring, then  $R[x]$  is Noetherian.

*Proof.* Let  $I \subseteq R[x]$  be a nonzero ideal. Define the set

$$J = \{a \in R \mid \text{there exists } f(x) \in I \text{ with leading coefficient } a\}.$$

We claim that  $J$  is an ideal of  $R$ . Towards this end, let  $a \in J$  and  $r \in R$ , with  $a \neq 0$ . Then, there exists  $f(x) \in I$  with  $f(x) = ax^n + O(x^{n-1})$ . We observe that  $rf(x) \in I$ , whence  $rax^n + O(x^{n-1}) \in I$ , so that  $ra \in J$ .

Now, let  $a, b \in J$ , and let  $f(x) = ax^n + O(x^{n-1})$  and  $g(x) = bx^m + O(x^{m-1})$ , with  $f(x), g(x) \in I$ . Without

loss of generality,  $n \geq m$ , so we may multiply  $x^{n-m}g(x) \in I$  to take  $x^{n-m}g(x) = bx^n + O(x^{n-1})$ , so that  $f(x) - x^{n-m}g(x) \in I$  and thus  $a - b \in J$ .

Now, since  $R$  is Noetherian, and  $J \subseteq R$  is an ideal, we have  $J = (a_1, \dots, a_n)$  for some  $a_1, \dots, a_n \in R$ . We may find corresponding elements of  $I$ , which we write  $f_i(x) = a_i x^{d_i} + O(x^{d_i-1})$ .

Set  $d = \max\{d_i \mid 1 \leq i \leq r\}$ . Consider the  $R$ -module

$$\begin{aligned} M &= (R + Rx + \dots + Rx^{d-1}) \cap I \\ &= \{\text{polynomials in } I \text{ with degree less than } d-1\} \cup \{0\}. \end{aligned}$$

Note that  $M$  is finitely generated as an  $R$ -module, as  $M \leq R + Rx + \dots + Rx^{d-1}$  and the latter is Noetherian as an  $R$ -module as it is a finitely generated module over a Noetherian ring.

We may write  $M = (g_1(x), \dots, g_s(x))$  as a result. We claim that

$$I = (f_1(x), \dots, f_r(x), g_1(x), \dots, g_s(x)).$$

Write  $I_0 = (f_1(x), \dots, f_r(x))$ . We claim that if  $f(x) \in I$  is such that  $\deg(f) > d-1$ , then there exists  $h(x) \in I_0$  such that  $\deg(f-h) < \deg(f)$ . This will allow us to take  $p(x) \in I$  and decompose it into constituent parts by repeatedly applying the claim to yield an element  $h \in I_0$  with  $\deg(p-h) \leq d-1$ , or  $p(x) - h(x) = 0$ , which yields an element of  $I_0$  and an element of  $M$ .

Now, let  $f(x) \in I$  have degree  $n \geq d$ . In particular,  $n \geq d_i$  for all  $i = 1, \dots, r$ . Writing

$$f(x) = cx^n + O(x^{n-1}),$$

we then have a leading coefficient  $c = \ell_1 a_1 + \dots + \ell_r a_r$  with  $\ell_1, \dots, \ell_r \in R$ . We take

$$h(x) = \ell_1 x^{n-d_1} f_1(x) + \dots + \ell_r x^{n-d_r} f_r(x),$$

and observe that  $h$  has leading term equal to  $cx^n$ . Thus,  $\deg(f-h) < \deg(f)$ . The rest follows from induction.  $\square$

The Hilbert Basis Theorem and the theorem on finitely generated modules over Noetherian rings together allow us to create a large family of Noetherian modules. For instance, if  $R$  is Noetherian, then

$$\left( \frac{R[x_1, \dots, x_n]}{I} \right) [y_1, \dots, y_m]$$

is Noetherian.

## A Taste of Homological Algebra

**Definition:** A sequence of  $R$ -modules homomorphisms

$$A \xrightarrow{f} B \xrightarrow{g} C$$

is called *exact* if  $\ker(g) = \text{im}(f)$ . More generally, a sequence of  $R$ -module homomorphisms

$$A_1 \xrightarrow{f_1} A_2 \xrightarrow{f_2} \dots \xrightarrow{f_n} A_n$$

is called exact if  $\ker(f_{i+1}) = \text{im}(f_i)$  for each  $i$ .

**Definition:** The special case of an exact sequence is one of the following form.

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

Here,  $f$  is injective,  $g$  is surjective, and  $\ker(g) = \text{im}(f)$ . We call such sequences *short exact* sequences.

**Example:**

(a) The sequence

$$0 \longrightarrow N \xrightarrow{\iota} M \xrightarrow{\pi} M/N \longrightarrow 0$$

is a short exact sequence.

(b) If  $M$  is an  $R$ -module, then  $M$  slots into a short exact sequence of  $R$ -modules given by the following.

$$0 \longrightarrow N \longrightarrow F \longrightarrow M \longrightarrow 0$$

Here,  $F$  is free and  $N$  is the module of relations.

If  $f: M \rightarrow N$  is an  $R$ -module homomorphism, and  $P$  is a fixed  $R$ -module, then there is a canonical induced homomorphism

$$\begin{aligned} f_*: \text{hom}(P, M) &\rightarrow \text{hom}(P, N) \\ [P \xrightarrow{\varphi} M] &\mapsto [P \xrightarrow{\varphi} M \xrightarrow{f} N] \end{aligned}$$

given by  $f_*(\varphi) = f \circ \varphi$ .

**Proposition:** Suppose we start with a short exact sequence of  $R$ -modules.

$$0 \longrightarrow L \xrightarrow{f} M \xrightarrow{g} N \longrightarrow 0$$

Then, the sequence

$$0 \longrightarrow \text{hom}(P, L) \xrightarrow{f_*} \text{hom}(P, M) \xrightarrow{g_*} \text{hom}(P, N)$$

is exact for all  $R$ -modules  $P$ .

**Remark:** The converse also holds.

*Proof.* Suppose the sequence of  $R$ -modules is exact. Then,  $f$  is injective,  $g$  is surjective, and  $\text{im}(f) = \ker(g)$ . We start by showing that  $f_*$  is injective.

Toward this end, let  $\varphi \in \ker(f_*)$ . Then, for all  $x \in P$ , we have  $f(\varphi(x)) = 0$ , whence  $\varphi(x) = 0$ , so  $\varphi$  is zero.

Next, we observe that  $\text{im}(f_*) \subseteq \ker(g_*)$ , as  $g_* \circ f_* = (g \circ f)_* = 0$  as the original sequence is exact. Now, if  $\varphi \in \ker(g_*)$ , then  $\varphi: P \rightarrow M$  is an  $R$ -module homomorphism such that  $g \circ \varphi = 0$ . If  $p \in P$ , then  $g(\varphi(p)) = 0$ , meaning that  $\varphi(p) \in \ker(g)$ . Since  $f$  is injective and  $\ker(g) = \text{im}(f)$ , we have that there is unique  $a \in L$  such that  $\varphi(p) = f(a)$ . Define  $\psi: P \rightarrow L$  by  $p \mapsto a$ . Observe that since  $f$  is a homomorphism, so too is  $\psi$ .  $\square$

Note that  $g_*$  may not be surjective. The cases for  $P$  where  $g_*$  is surjective are special to have their own definition.

**Definition:** An  $R$ -module  $P$  is called *projective* if, for all short exact sequences of  $R$ -modules

$$0 \longrightarrow L \xrightarrow{f} M \xrightarrow{g} N \xrightarrow{0} 0$$

the sequence

$$0 \longrightarrow \text{hom}(P, L) \xrightarrow{f_*} \text{hom}(P, M) \xrightarrow{g_*} \text{hom}(P, N) \xrightarrow{0} 0$$

is also exact.

Equivalently, this means that for all surjections  $M \xrightarrow{\pi} N \rightarrow 0$ , and all homomorphisms  $\varphi: P \rightarrow N$ , there is some  $\tilde{\varphi}: P \rightarrow M$  such that  $\pi \circ \tilde{\varphi} = \varphi$ .

$$\begin{array}{ccccc} & & P & & \\ & \swarrow \tilde{\varphi} & \downarrow \varphi & & \\ M & \xrightarrow{\pi} & N & \longrightarrow & 0 \end{array}$$

Our goal now is to determine exactly when an  $R$ -module is projective. As it turns out, there is a very tidy characterization.

**Lemma:** Every free module is projective.

*Proof.* Let  $F$  be a free  $R$ -module with basis  $X$ , and let  $M \xrightarrow{\pi} N \rightarrow 0$  be a surjection of  $R$ -modules. Suppose we have a homomorphism  $\varphi$  satisfying the following diagram.

$$\begin{array}{ccccc} & & F & & \\ & & \downarrow \varphi & & \\ M & \xrightarrow{\pi} & N & \longrightarrow & 0 \end{array}$$

Restricting  $\varphi$  to  $X$ , we see that since  $\pi$  is surjective, we may find  $m_x \in M$  such that  $\pi(m_x) = \varphi(x)$  for each  $x$ . Defining a set map  $\tilde{\varphi}: X \rightarrow M$  mapping  $x \mapsto m_x$ , we observe that there is then an  $R$ -module homomorphism  $\tilde{\varphi}: F \rightarrow M$  uniquely extending  $\tilde{\varphi}$ , by the universal property of free modules.  $\square$

**Theorem** (Characterization of Projective Modules): An  $R$ -module  $P$  is projective if and only if  $P$  is a direct summand of a free module. That is, if and only if there exists a free  $R$ -module  $F$  and an  $R$ -module  $Q$  such that

$$F \cong P \oplus Q$$

as an external direct sum.

We can show one direction right now.

*Proof of the Reverse Direction.* Suppose  $F = P \oplus Q$  as an internal direct sum for some  $R$ -module  $Q$  and a free  $R$ -module  $P$ . Then,  $F$  is equipped with a projection map  $\pi_P: F \rightarrow P$  taking  $p \oplus q$  to  $p$ , and an inclusion map  $\iota_P: P \hookrightarrow F$  mapping  $p \mapsto p \oplus 0$ . We then take the following diagram of homomorphisms, where  $\varphi: P \rightarrow N$  is the homomorphism we want to extend to  $g$ .

$$\begin{array}{ccccc} & & F & & \\ & \swarrow \tilde{\psi} & \downarrow \pi_P & \searrow \psi & \\ & & P & & \\ & & \downarrow \varphi & & \\ M & \xrightarrow{g} & N & \longrightarrow & 0 \end{array}$$

Now, we observe that by definition, we have  $g \circ \tilde{\psi} = \varphi \circ \pi_P$ , so we may define  $\tilde{\varphi}: P \rightarrow M$  by taking

$$\tilde{\varphi} = \tilde{\psi} \circ \iota_P.$$

This maps  $p \mapsto p \oplus 0 \mapsto \tilde{\psi}(p \oplus 0) = \varphi(p)$ .  $\square$

For the reverse direction, we need to be able to characterize when exactly a module is a direct sum of submodules.

**Definition:** Let  $f: M \rightarrow N$  be an  $R$ -module homomorphism. Then, we say  $s: N \rightarrow M$  is a *splitting* (or *section*) if  $f \circ s = \text{id}_N$ . In other words, a splitting is a right-inverse.

We observe automatically that if  $f$  has a splitting, then  $f$  is surjective and  $s$  is injective, by the equivalent characterization of surjective maps as those being with (necessarily injective) right-inverses.

**Lemma:** If  $f: M \rightarrow N$  admits a splitting  $s: N \rightarrow M$ , then  $M \cong s(N) \oplus \ker(f) \cong N \oplus \ker(f)$ .

*Proof.* First, we show that  $s(N) + \ker(f) = M$ . Letting  $m \in M$ , then we set  $m' = m - s(f(m))$ . Then,

$$\begin{aligned} f(m') &= f(m) - f(s(f(m))) \\ &= f(m) - (f \circ s)(f(m)) \\ &= f(m) - f(m) \\ &= 0, \end{aligned}$$

so  $m' \in \ker(f)$ , and  $M \subseteq s(N) + \ker(f)$ .

Next, we show that  $s(N) \cap \ker(f) = 0$ . If  $m \in s(N) \cap \ker(f)$ , then  $m = s(n)$  for some  $n \in N$ , and  $f(m) = f(s(n)) = n = 0$ , whence  $m = 0$ .  $\square$

**Definition:** We say the sequence

$$0 \longrightarrow L \xrightarrow{f} M \xrightarrow{g} N \longrightarrow 0$$

$\nwarrow \text{---} s \text{---}$

splits if  $g$  has a splitting, so by the lemma and exactness, we have  $M \cong L \oplus N$ .

**Lemma:** Let  $M \xrightarrow{g} P \rightarrow 0$  be a surjection of  $R$ -modules with  $P$  projective. Then,  $g$  admits a splitting.

*Proof.* Using the projectivity of  $P$  in the following diagram, we obtain our desired section  $s$  such that  $g \circ s = \text{id}$ .

$$\begin{array}{ccc} M & \xrightarrow{g} & P \longrightarrow 0 \\ & \nwarrow s & \uparrow \text{id} \\ & & P \end{array}$$

$\square$

Now, we can finally prove the forward direction of the characterization.

*Proof of the Forward Direction.* Let  $P$  be projective. From the universal property of  $R$ -modules, we know that there is a surjective  $R$ -module homomorphism  $F \xrightarrow{\pi} P \rightarrow 0$ . Thus,  $\pi$  splits, and we are done.  $\square$

## Groups

We begin our discussion of group theory with group actions.

**Definition:** Let  $X$  be a set, and let  $G$  be a group. A (left)  $G$ -action on  $X$  is a map

$$\begin{aligned} \mu: G \times X &\rightarrow X \\ (g, x) &\mapsto g \cdot x \end{aligned}$$

that satisfies

- $e \cdot x = x$  for all  $x \in X$
- $g \cdot (h \cdot x) = (gh) \cdot x$ .

We call  $X$  a  $G$ -set.

**Definition:** A group action induces a homomorphism

$$\rho: G \rightarrow \text{Sym}(X)$$

$$g \mapsto \pi_g,$$

where  $\pi_g(x) = g \cdot x$ . This is known as the *permutation representation* of the action.

**Definition:** We say the action of  $G$  on  $X$  is *faithful* if the induced map  $\rho$  is injective.

**Remark:** Any action  $\rho$  admits a faithful action by “modding out” by the kernel,

$$\ker(\rho) := \{g \in G \mid g \cdot x = x \text{ for all } x \in X\}.$$

Then, the action of  $G/\ker(\rho)$  on  $X$  is given by  $(g\ker(\rho)) \cdot x = g \cdot x$ .

**Definition:** For an element  $x \in X$ , we define the *stabilizer* of  $x$  to be

$$G_x = \{g \in G \mid g \cdot x = x\}.$$

**Definition:** For  $x \in X$ , the *orbit* of  $x$  is given by

$$G \cdot x = \{g \cdot x \mid g \in G\}.$$

**Definition:** An action is called *transitive* if there is one orbit for the action.

The orbits in fact partition  $X$  under the equivalence relation  $x \sim y$  whenever there is  $g \in G$  such that  $g \cdot x = y$ . The set of all distinct orbits is then denoted

$$\begin{aligned} X/G &= \{G \cdot x_i \mid i \in I\} \\ &= \bigsqcup_{i \in I} G \cdot x_i. \end{aligned}$$

**Lemma (Orbits and Stabilizers):** Let  $x \in X$  be fixed. There exists a bijection between the set of all left cosets  $G/G_x$  and the orbit  $G \cdot x$ . In particular, if  $X$  and  $G$  are finite, then  $|G \cdot x| = [G : G_x]$ .

*Proof.* Consider the map  $f_x: G \rightarrow G \cdot x$  given by  $g \mapsto g \cdot x$ . By definition of the orbit,  $f_x$  is necessarily surjective. For  $g \in G$  and  $h \in G_x$ , we observe that

$$\begin{aligned} f_x(gh) &= gh \cdot x \\ &= g \cdot (h \cdot x) \\ &= g \cdot x \\ &= f_x(g). \end{aligned}$$

Therefore, by the first isomorphism theorem,  $f_x$  induces a well-defined map  $\bar{f}_x: G/G_x \rightarrow G \cdot x$  that is surjective. We claim that  $\bar{f}_x$  is in fact injective.

Suppose  $f_x(g_1) = f_x(g_2)$ . Then,  $g_1 \cdot x = g_2 \cdot x$ , or that  $g_2^{-1}g_1 \cdot x = x$ , meaning that  $g_2^{-1}g_1 \in G_x$ , or that the cosets  $g_1G_x = g_2G_x$ .  $\square$

**Corollary:** Let  $G$  be a finite group acting on a finite set  $X$ , and let  $\{x_1, \dots, x_r\}$  be distinct representatives for  $X/G$ .

(i)

$$\begin{aligned} |X| &= \sum_{i=1}^r |G \cdot x_i| \\ &= \sum_{i=1}^r [G : G_{x_i}] \\ &= \sum_{i=1}^r \frac{|G|}{|G_{x_i}|}. \end{aligned}$$



(ii)

$$|G| = |G \cdot x| |G_x|$$

for all  $x \in X$ .

## Canonical Examples of Group Actions

**Example** (Symmetric Polynomials): Let  $R = F[x_1, \dots, x_n]$ . We let  $S_n$  act on  $R$  by permuting the indices, given by

$$\begin{aligned}\sigma \cdot x_i &= x_{\sigma(i)} \\ \sigma \cdot k &= k\end{aligned}$$

for all  $x_i$  and all  $k \in F$ . We claim that this is a faithful action.

If  $\sigma \in S_n$  is such that  $\sigma \in \ker(\rho)$ , then  $\sigma \cdot f(x_1, \dots, x_n) = f(x_1, \dots, x_n)$  for all  $f \in F[x_1, \dots, x_n]$ . In particular, this means that  $\sigma \cdot x_i = x_i$  for all  $i$ , and  $\sigma \cdot k = k$ , whence  $\sigma(i) = i$  for all  $i$ , so  $\sigma = e$ .

Therefore, we get the injective group homomorphism  $\rho: S_n \rightarrow \text{Sym}(R)$ . In this case, every  $\rho(\sigma): R \rightarrow R$  is in fact a ring automorphism. We say that  $S_n$  acts on  $R$  by ring automorphisms.

**Definition:** The fixed points of the action  $\rho: S_n \rightarrow \text{Sym}(R)$  are known as the *symmetric polynomials*.

**Definition** (Elementary Symmetric Polynomials): The elementary symmetric polynomials are defined as follows.

$$\begin{aligned}s_1 &= x_1 + \dots + x_n \\ s_2 &= \sum_{i < j} x_i x_j \\ s_3 &= \sum_{i < j < k} x_i x_j x_k \\ &\vdots \\ s_n &= x_1 \cdots x_n.\end{aligned}$$

The fundamental theorem of symmetric polynomials holds that every symmetric polynomial  $f \in F[x_1, \dots, x_n]$  has a unique expression as  $g(s_1, \dots, s_n)$ . In other words,

$$F[x_1, \dots, x_n]^{S_n} = F[s_1, \dots, s_n].$$

A version of this can in fact be proven using Galois theory, but we will prove it using a more “pedestrian” approach.

To see existence, we start by reducing to the homogeneous case, by observing that we may separate any  $f \in F[x_1, \dots, x_n]$  into polynomials with the property that

$$f(\lambda x_1, \dots, \lambda x_n) = \lambda^d f(x_1, \dots, x_n),$$

where  $d$  is the degree of the homogeneous polynomial. In particular, we may uniquely write

$$f = p_1 + \dots + p_r,$$

where each of the  $p_i$  is homogeneous with degree  $d_i$ , and  $d_1 > d_2 > \dots > d_r$ . If  $f$  is symmetric, then for any  $\sigma \in S_n$ ,  $\sigma \cdot p_i$  is also homogeneous of the same degree, with

$$\begin{aligned}f &= \sigma \cdot p_1 + \dots + \sigma \cdot p_r \\ &= p_1 + \dots + p_r\end{aligned}$$

by uniqueness, so that each of the homogeneous components of  $f$  is symmetric.

**Example:** Given a regular geometric shape (such as a square, circle, triangle, cube, etc.), we consider the group  $G$  consisting of all isometries  $S \rightarrow S$ .

For example, if  $S$  is a square, then the group of isometries is  $D_4$ . We claim that the group  $D_4$  has order 8, consisting of the set of combinations of the four pure rotations and four reflections about the symmetry axes of the square. To see this, let  $V = \{a, b, c, d\}$  be the set of vertices of the square, and let  $G$  act on  $V$ . We

observe that this action is transitive, so that  $|G \cdot a| = 4$ . Meanwhile the stabilizer of  $A$  consists of the identity map and the reflection about the diagonal axis of symmetry that passes through  $a$ . In particular, this means that  $|\text{stab}_G(a)| = 2$ , whence  $|G| = 8$ .

In the general case, we can define the group  $D_n$  to be the set of rotations  $r_1, \dots, r_n$  with an angle of  $\frac{2\pi k}{n}$  together with the reflections about the symmetry axes,  $s_1, \dots, s_n$ , subject to the relations that

$$\begin{aligned} r_1^n &= e \\ s_1^2 &= e \\ s_1 r_1 s_1^{-1} &= r_1^{-1}. \end{aligned}$$

**Example:** If  $G$  is a group, then  $G$  can in fact act on itself. The most canonical way of doing so is via left-multiplication, giving  $g \cdot h = gh$ .

Via the permutation representation, this induces a homomorphism

$$\rho: G \rightarrow \text{Sym}(G),$$

and since the action is faithful, this is injective. Immediately, we obtain *Cayley's Theorem*, which states that every group is isomorphic to a subgroup of a permutation group. This is known as the left-regular permutation representation of  $G$ .

We can extend this regular permutation to cosets. If  $H \leq G$ , then  $G$  acts on the coset space  $G/H$  by  $g \cdot aH = gaH$ . This action is transitive, and computing the stabilizer gives  $gaH = aH$  if and only if  $a^{-1}ga \in H$ , if and only if  $g \in aHa^{-1}$ . In particular, this means that  $g \in G_{aH}$  if and only if  $gaH$  is a conjugate of  $H$ .

Given the permutation representation homomorphism  $\rho: G \rightarrow \text{Sym}(G/H)$ , we see that

$$\ker(\rho) = \bigcap_{a \in G} aHa^{-1},$$

which is necessarily normal.

**Example:** The action of  $G$  on itself by conjugation, given by  $g \mapsto \iota_g$ , where

$$\iota_g(y) = gyg^{-1}$$

is an action of the group on itself by automorphisms. The image  $G \rightarrow \text{aut}(G)$  under this action is known as the *inner automorphism group* of  $G$ .

The orbit of an element  $x$  under conjugation is known as the *conjugacy class* of  $x$ , and will be written  $K_G(x)$ . The stabilizers are known as the *centralizer* of  $x$ , and are written as  $Z_G(x)$ . The kernel of the permutation representation for conjugation is the *center* of the group  $G$ , and is written  $Z(G)$ .

We observe that if  $G$  is a finite group with  $g_1, \dots, g_r$  representatives for the conjugacy classes, then

$$|G| = |Z(G)| + \sum_{i=1}^r [G : Z_G(g_i)].$$

It is possible to extend conjugation naturally to  $P(G)$ , by taking  $g \cdot A = gAg^{-1}$ . The stabilizers are denoted as  $N_G(T)$ . If  $H$  is an actual subgroup of  $G$ , then  $N_G(H)$  is known as the *normalizer* of  $H$  in  $G$ . Similarly, by applying orbit-stabilizer, we find that  $[G : N_G(H)]$  is the number of distinct subgroups conjugate to  $H$ .

## The Sylow Theorems

Let  $G$  be a finite group with  $|G| = p^m r$ , where  $p$  is a prime,  $m \geq 1$ , and  $p \nmid r$ .

**Definition:** A  $p$ -Sylow subgroup of  $G$  is a subgroup of order  $p^m$ . That is, a  $p$ -Sylow subgroup of  $G$  is a maximal  $p$ -subgroup of  $G$ .

There are three Sylow theorems. They are very useful.

**Theorem (Sylow Theorems):** Let  $G$  be a finite group with  $|G| = p^m r$ .

- (1) A  $p$ -Sylow subgroup exists.
- (2) If  $P$  and  $Q$  are  $p$ -Sylow subgroups, then there is  $g \in G$  such that  $Q = gPg^{-1}$ .
- (3) Let  $n_p$  denote the number of  $p$ -Sylow subgroups. Then, we have

$$\begin{aligned} n_p &\equiv 1 \pmod{p} \\ n_p &\mid r. \end{aligned}$$

## Semidirect Products

**Definition:** Let  $\{G_i\}_{i \in I}$  be a family of groups.

The direct product  $\prod_{i \in I} G_i$  is a group with coordinatewise operations and underlying set  $\{(g_i)_{i \in I} \mid g_i \in G_i\}$ .

It turns out that there is no analogue to the external direct sum for groups. This follows from the fact that if we have homomorphisms  $f_1: G_1 \rightarrow H$  and  $f_2: G_2 \rightarrow H$ , then we would need to glue the homomorphisms by

$$f(g_1, g_2) = f_1(g_1)f_2(g_2).$$

In particular, we get

$$\begin{aligned} f((g_1, g_2)(h_1, h_2)) &= f(g_1h_1, g_2h_2) \\ &= f_1(g_1)f_1(h_1)f_2(g_2)f_2(h_2) \\ &= f_1(g_1)f_2(g_2)f_1(h_1)f_2(h_2) \end{aligned}$$

if and only if  $f_1(h_1)$  and  $f_2(g_2)$  commute for all  $h_1 \in G_1$  and  $g_2 \in G_2$ .

We start by trying to find a scenario where we in fact can construct  $G \cong H \times N$  for some subgroups  $H$  and  $N$ .

**Lemma:** Let  $G$  be a group,  $H, N \leq G$ .

- (a) If both  $N$  and  $H$  are normal, and  $N \cap H = \{e\}$ , then  $nh = hn$  for all  $n \in N$  and  $h \in H$ .
- (b) If  $N$  is normal, then  $NH$  is a subgroup of  $G$ .

*Proof.*

- (a) We let  $[h, n] = hnh^{-1}n^{-1}$ . Since  $H \cap N = \{e\}$ , and the commutator is contained in both  $H$  and  $N$ , it follows that  $hnh^{-1}n^{-1} = e$ , whence  $hn = nh$ .
- (b) Let  $N$  be normal,  $h_1, h_2 \in H$ ,  $n_1, n_2 \in N$ . Then,

$$\begin{aligned} (n_1h_1)(n_2h_2) &= n_1(h_1n_2h_1^{-1})(h_1h_2) \\ &\in NH, \end{aligned}$$

so  $NH$  is closed under multiplication. Similarly,

$$\begin{aligned} (nh)^{-1} &= h^{-1}n^{-1} \\ &= (h^{-1}n^{-1}h)h^{-1} \\ &\in NH, \end{aligned}$$

so  $NH$  is a subgroup.

□

**Theorem:** Let  $G$  be a group,  $H, N$  normal, and suppose that

- (i)  $H \cap N = \{e\}$ ;
- (ii)  $G = NH$ .

Then,  $G \cong NH$ .

*Proof.* The map  $N \times H \rightarrow G$  given by  $(n, h) \mapsto nh$  is a group homomorphism since  $H$  and  $N$  commute elementwise. By (ii), the map is surjective, and by (i), the map is injective. □

An interesting nonexample is the case of  $S_3$ . We observe that the groups  $N = \langle (1, 2, 3) \rangle$  and  $H = \langle (1, 2) \rangle$  have  $N \cap H = \{e\}$ ,  $S_3 = NH$ , but  $N \cong \mathbb{Z}/3\mathbb{Z}$  and  $H \cong \mathbb{Z}/2\mathbb{Z}$ , meaning that  $S_3 \not\cong N \times H$ . Instead, we have a different construction.

**Definition:** Let  $N$  and  $H$  be groups. Suppose we have a group homomorphism

$$f: H \rightarrow \text{aut}(N).$$

The *semidirect product* with respect to  $f$  is given by

$$G = N \rtimes_f H,$$

with underlying set  $N \times H$  and operations given by

$$\begin{aligned} (n_1, h_1) \cdot (n_2, h_2) &= (n_1 f(h_1)(n_2), h_1 h_2) \\ (n, h)^{-1} &= (f(h^{-1})(n^{-1}), h^{-1}). \end{aligned}$$

The semidirect product in fact allows us another tool for classifying groups. This often appears in Galois theory.

To start, we can use the Sylow theorems to see that groups of order  $pq$  with  $p, q$  primes are not simple groups. In fact, there is a unique  $q$ -Sylow subgroup of  $G$ , which we call  $Q$ , which is then normal in  $G$ . Since  $Q$  is also prime, we have  $Q \cong \mathbb{Z}/q\mathbb{Z}$ .

We consider now the group homomorphism

$$\begin{aligned} f: G &\rightarrow \text{aut}(Q) \\ g &\mapsto \iota_g, \end{aligned}$$

where  $\iota_g$  is the inner automorphism given by  $\iota_g(x) = gxg^{-1}$ . Since  $Q$  is normal,  $f$  is well-defined. Furthermore, since  $Q$  is abelian,  $Q \leq \ker(f)$ , whence  $f$  induces a homomorphism  $\bar{f}: G/Q \rightarrow \text{aut}(Q)$ .

We have that  $\text{aut}(\mathbb{Z}/q\mathbb{Z}) \cong (\mathbb{Z}/q\mathbb{Z})^\times$ , which is cyclic of order  $q - 1$ . Meanwhile  $|G/Q| = p$ , which is prime, so we may express  $\bar{f}: \mathbb{Z}/p\mathbb{Z} \rightarrow (\mathbb{Z}/q\mathbb{Z})^\times$ .

First, if  $p \nmid q - 1$ , then Lagrange's Theorem implies that  $\bar{f}$  is trivial. In particular,  $|\ker(\bar{f})| = 1$ , in which case we would have  $\text{im}(\bar{f}) \leq (\mathbb{Z}/q\mathbb{Z})^\times$ , which cannot happen as  $p \nmid q - 1$ , meaning that we have  $|\ker(\bar{f})| = p$ , meaning  $\bar{f}$  is trivial.

If  $\bar{f}$  is trivial, then  $f$  is in fact trivial, so  $gxg^{-1} = x$  for all  $g \in G$ , meaning  $Q \leq Z(G)$ . This yields a surjection  $G/Q \rightarrow G/Z(G)$ , and since  $G/Q$  is cyclic, so too is  $G/Z(G)$ , whence  $G/Z(G)$  is cyclic, hence  $G$  is abelian. In particular, this gives

$$G \cong \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/q\mathbb{Z}.$$

Meanwhile, if  $p|q - 1$ , then there is a nontrivial homomorphism  $f: \mathbb{Z}/p\mathbb{Z} \rightarrow (\mathbb{Z}/q\mathbb{Z})^\times$ . There are then  $p - 1$  distinct nontrivial automorphisms, from which we may take  $Q \rtimes_f P$ .

We claim that the various choices for  $f$  yield isomorphic groups.

**Proposition:** Let  $P = \mathbb{Z}/p\mathbb{Z}$  and  $Q = \mathbb{Z}/q\mathbb{Z}$  with  $p, q$  prime and  $q \equiv 1$  modulo  $p$ . If

$$f, g: P \rightarrow \text{aut}(Q)$$

are nontrivial homomorphisms, then  $Q \rtimes_f P$  and  $Q \rtimes_g P$  are isomorphic.

*Proof.* Since  $f$  and  $g$  are both nontrivial, it follows that  $f$  and  $g$  are injective (again by Lagrange's Theorem), meaning that since  $\text{aut}(Q)$  is cyclic, there is a unique subgroup of order  $p$ , giving  $\text{im}(f) \cong \text{im}(g)$ .

Fix a generator for  $P$ , called  $P = \langle s \rangle$ . Since  $f(s)$  and  $g(s)$  are both generators for  $\text{im}(f)$  and  $\text{im}(g)$ , there is some  $\ell$  such that  $g(s) = f(s)^\ell$ . In particular, this means that  $g(v) = f(v)^\ell$  for all  $v \in P$ .

Define the homomorphism  $\varphi: Q \rtimes_g P \rightarrow Q \rtimes_f P$  by  $(x, y) \mapsto (x, y^\ell)$ . As a map of sets,  $\varphi$  is bijective since  $\gcd(\ell, p) = 1$ .

Now, we show that  $\varphi$  is a homomorphism.

$$\begin{aligned} \varphi((x_1, y_1)(x_2, y_2)) &= \varphi(x_1 g(y_1)(x_2), y_1 y_2) \\ &= (x_1 g(y_1)(x_2), y_1^\ell y_2^\ell) \\ &= (x_1 f(y_1^\ell)(x_2), y_1^\ell y_2^\ell) \\ &= (x_1, y_1^\ell)(x_2, y_2^\ell). \end{aligned}$$

□

## Conjugacy in $S_n$ and $A_n$

To understand conjugacy in  $S_n$  and  $A_n$ , we recall some properties of the symmetric group.

- (1) Elements of  $S_n$  are called permutations, and are written as

$$\sigma = \begin{pmatrix} 1 & 2 & \cdots & n \\ \sigma(1) & \sigma(2) & \cdots & \sigma(n) \end{pmatrix}.$$

- (2) An element of the form  $(i_1, i_2)$  is called a transposition, and an element of the form  $(i_1, \dots, i_k)$  is called a  $k$ -cycle.
- (3) Every  $\sigma \in S_n$  can be written as a product of disjoint cycles.
- (4) Disjoint cycles commute with each other.
- (5) The order of a  $k$ -cycle is  $k$ , and the order of

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_r,$$

with respective lengths  $k_1, \dots, k_r$  is  $o(\sigma) = \text{lcm}(k_1, \dots, k_r)$ .

- (6) Every  $\sigma \in S_n$  can be written as a product of transpositions.

**Definition:** A permutation is called *even* if it can be written as an expression

$$\sigma = \tau_1 \cdots \tau_{2k}$$

as an even number of transpositions. We define the *alternating group*,  $A_n$ , to be the set of even permutations.

**Theorem:** We have  $[S_n : A_n] = 2$ , and every transposition is odd.

*Proof.* Consider the function

$$\Delta = \prod_{1 \leq i < j \leq n} (x_i - x_j).$$

Letting  $S_n$  act on  $\mathbb{Z}[x_1, \dots, x_n]$  by permuting the indices, we note that  $\sigma \cdot f = \pm f$ . The sign of the permutation  $\sigma$  is defined to be  $k$  such that

$$\sigma(f) = \text{sgn}(\sigma)f$$

As a result, we find that transpositions are odd, and  $\text{sgn}$  is surjective, so  $S_n/A_n \cong \mathbb{Z}/2\mathbb{Z}$ .  $\square$

In order to understand conjugacy in  $S_n$  and  $A_n$ , we need to understand what the conjugacy class of a cycle is.

**Proposition:** Let  $g \in S_n$  be arbitrary, and let  $\sigma = (i_1, \dots, i_k)$  be a  $k$ -cycle. Then,

$$g\sigma g^{-1} = (g(i_1), \dots, g(i_k)).$$

*Proof.* Computing directly, we have

$$g\sigma g^{-1}(g(i_\ell)) = \begin{cases} g(i_{\ell+1}) & \ell < k \\ g(i_1) & \ell = k \end{cases}.$$

Now, if  $x \notin \{g(i_1), \dots, g(i_k)\}$ , then  $g^{-1}(x) \notin \{i_1, \dots, i_k\}$ , meaning that  $\sigma$  fixes  $g^{-1}(x)$ , so that  $g\sigma g^{-1}$  fixes  $x$ .  $\square$

Thus, we have that the conjugacy class of any  $k$ -cycle is the set of all  $k$ -cycles:

$$K_{S_n}(\sigma) = K_{S_n}((1, \dots, k)).$$

Computing the number of  $k$ -cycles is an exercise in combinatorics, meaning that we have

$$|K_{S_n}(\sigma)| = \frac{1}{k}(n(n-1)\cdots(n-(k-1))).$$

In general, we have that two permutations  $\sigma, \tau \in S_n$  are conjugate if and only if they have the same **cycle type** (partitions of  $n$  into cycle lengths).

Therefore, we have that the number of conjugacy classes in  $S_n$  is equal to the number of partitions of  $n$ .

Next, we compute centralizers for a  $k$ -cycle. If  $k = n$ , then we see that  $Z_{S_n}(\sigma) = \langle \sigma \rangle$ , as if  $\tau \in Z_{S_n}(\sigma)$ , then

$$\tau(i_1) = i_r$$

for some  $r = 1, \dots, n$ . Multiplying by  $\sigma^\ell$  for a suitable  $\ell$  yields  $\sigma^\ell \tau(i_1) = i_1$ , or that

$$(\sigma^\ell \tau) \sigma (\sigma^\ell \tau)^{-1} = \sigma,$$

whence  $\sigma^\ell \tau(i_j) = i_j$ , or that  $\tau \in \langle \sigma \rangle$ .

Now, if  $k \leq n$ , then we observe that for any  $\tau \in Z_{S_n}(\sigma)$ , we have that  $\tau|_{\{i_1, \dots, i_k\}}$  is equal to  $\sigma^m|_{\{i_1, \dots, i_k\}}$ , while there is no restriction for any  $\tau$  on  $\{1, \dots, n\} \setminus \{i_1, \dots, i_k\}$ . Thus, we have that

$$Z_{S_n}(\sigma) \cong \langle \sigma \rangle \times S_{n-k},$$

since the subgroups commute, so that

$$\begin{aligned} |Z_{S_n}(\sigma)| &= |\langle \sigma \rangle| \cdot |S_{n-k}| \\ &= k \cdot (n-k)!. \end{aligned}$$

Orbit-stabilizer yields our original fact that

$$|K_{S_n}(\sigma)| = \frac{1}{k}(n(n-1)\cdots(n-(k-1))).$$

If  $\pi = \sigma_1 \cdots \sigma_r$  has a cycle decomposition with pairwise distinct lengths  $k_1, \dots, k_r$ , then for any  $\rho \in Z_{S_n}(\pi)$ , we have

$$\begin{aligned} \rho\pi\rho^{-1} &= (\rho\sigma_1\rho^{-1}) \cdots (\rho\sigma_r\rho^{-1}) \\ &= \sigma_1 \cdots \sigma_r, \end{aligned}$$

whence

$$\begin{aligned} \rho\sigma_1\rho^{-1} &= \sigma_1 \\ &\vdots \\ \rho\sigma_r\rho^{-1} &= \sigma_r. \end{aligned}$$

In particular, this means that  $\rho \in Z_{S_n}(\sigma_i)$  for each  $i = 1, \dots, r$ . This gives the isomorphism

$$Z_{S_n}(\pi) \cong \prod_{i=1}^r \langle \sigma_i \rangle \times S_{n-(k_1+\cdots+k_r)}.$$

Finally, if  $\pi = \sigma_1 \cdots \sigma_\ell$  all with length  $k$ , we observe that an arbitrary permutation  $\tau \in Z_{S_n}(\pi)$  may simply permute the cycles  $\sigma_1, \dots, \sigma_\ell$  in addition to what has been seen earlier. In particular, we have

$$Z_{S_n}(\pi) \cong \left( \prod_{i=1}^{\ell} \sigma_i \rtimes_f S_\ell \right) \times S_{n-k\ell},$$

where the semidirect product emerges from the fact that the permutations that permute the cycles  $\sigma_1, \dots, \sigma_\ell$  induce a nontrivial automorphism on

$$\begin{aligned} N &= \prod_{i=1}^{\ell} \sigma_i \times S_{n-k\ell} \\ &\leq Z_{S_n}(\pi). \end{aligned}$$

Thus, we get the most general case.

**Theorem:** Let  $\pi$  be a product of  $\ell_1, \dots, \ell_t$  cycles with lengths  $k_1, \dots, k_t$  respectively. Then,

$$Z_{S_n}(\pi) \cong Z_{S_{k_1\ell_1}}(\pi_1) \times \cdots \times Z_{S_{k_t\ell_t}}(\pi_t) \times S_{\{\text{what's left}\}},$$

where each  $\pi_j \in S_{k_j\ell_j}$  is a product of  $\ell_j$  cycles with length  $k_j$ .

We now discuss the special case of conjugacy in the alternating group. Simply by definition, if  $\sigma \in A_n$ , then we know that  $K_{S_n}(\sigma) \leq A_n$ . The question is when  $K_{A_n}(\sigma) = K_{S_n}(\sigma)$ . There is in fact a very simple characterization.

**Lemma:** Let  $\sigma \in A_n$ . Then,

$$K_{A_n}(\sigma) = K_{S_n}(\sigma)$$

if and only if  $Z_{S_n}(\sigma)$  contains an odd permutation.

*Proof.* We have that  $K_{S_n}(\sigma) = K_{A_n}(\sigma)$  if and only if for all  $g \in S_n$ , there is  $h \in A_n$  such that  $g\sigma g^{-1} = h\sigma h^{-1}$ . Yet, this means that for all  $g \in S_n$ , there is  $h \in A_n$  such that  $h^{-1}g \in Z_{S_n}(\sigma)$ . By the definition of cosets, this holds if and only if  $gZ_{S_n}(\sigma) = hZ_{S_n}(\sigma)$ , whence it holds if and only if  $S_n = (A_n)(Z_{S_n}(\sigma))$ .

Since  $A_n$  consists exclusively of even permutations, and  $S_n$  contains odd permutations, it follows that this



holds if and only if  $Z_{S_n}(\sigma)$  contains an odd permutation. □

**Theorem:** The group  $A_5$  is simple.

*Proof.* We start by proving for  $n = 5$ . In order to do this, we count conjugacy classes in  $S_5$  and  $A_5$ . The partitions of 5 correspond to the following conjugacy classes.

- Cycles of type 5:

$$\begin{aligned} |K_{S_5}((1, 2, 3, 4, 5))| &= \frac{5!}{5} \\ &= 24. \end{aligned}$$

- Cycles of type 4 + 1:

$$\begin{aligned} |K_{S_5}(1, 2, 3, 4)| &= \frac{(5)(4)(3)(2)}{4} \\ &= 30. \end{aligned}$$

- Cycles of type 3 + 1 + 1:

$$\begin{aligned} |K_{S_5}((1, 2, 3))| &= \frac{(5)(4)(3)}{3} \\ &= 20. \end{aligned}$$

- Cycles of type 2 + 1 + 1 + 1:

$$\begin{aligned} |K_{S_5}((1, 2))| &= \binom{5}{2} \\ &= 10. \end{aligned}$$

- Cycles of type 3 + 2:

$$|K_{S_5}((1, 2, 3)(4, 5))| = 20$$

- Cycles of type 2 + 2 + 1:

$$\begin{aligned} |K_{S_5}((1, 2)(3, 4))| &= \frac{|S_5|}{|Z_{S_5}((1, 2)(3, 4))|} \\ &= \frac{5!}{2^3} \\ &= 15, \end{aligned}$$

where we use the fact that the stabilizer is isomorphic to  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \rtimes_f \mathbb{Z}/2\mathbb{Z}$ .

In  $A_5$ , we only care about the even permutations — that is, the 5-cycles, 3-cycles, and the cycles of type 2 + 2.

- The conjugacy class of the 1-cycle consists exclusively of  $e$ . Therefore, the 1-cycles endow  $A_5$  with 1 conjugacy class with 1 element.
- To find the conjugacy class(es) of the 3-cycles, observe that the centralizer  $Z_{S_n}((1, 2, 3))$  includes the transposition  $(4, 5)$ . Therefore, by the lemma,  $K_{A_5}((1, 2, 3)) = K_{S_5}((1, 2, 3))$ . Therefore, the 3-cycles endow  $A_5$  with 1 conjugacy class with 20 elements.

- To find the conjugacy class(es) of the cycles with type  $2 + 2$ , observe that  $(1, 2) \in Z_{S_5}((1, 2)(3, 4))$ . Therefore, by the lemma,  $K_{A_5}((1, 2)(3, 4)) = K_{S_5}((1, 2)(3, 4))$ . Therefore, the cycles of type  $2 + 2$  endow  $A_5$  with 1 conjugacy class with 15 elements.
- To find the conjugacy class(es) for the 5 cycles, observe that if  $\sigma$  is a 5-cycle, then  $Z_{S_5}(\sigma) = \langle \sigma \rangle$ , so  $K_{A_5}(\sigma) \subsetneq K_{S_5}(\sigma)$ . Yet, we have that  $Z_{S_5}(\sigma) = Z_{A_5}(\sigma)$ , so

$$\begin{aligned} |K_{A_5}(\sigma)| &= \left| \frac{A_5}{Z_{S_5}(\sigma)} \right| \\ &= \left| \frac{A_5}{Z_{A_5}(\sigma)} \right| \\ &= \frac{1}{2} |K_{S_5}(\sigma)| \\ &= 12. \end{aligned}$$

Therefore, the 5-cycles endow  $A_5$  with 2 disjoint conjugacy classes with 12 elements each.

The class equation for  $A_5$  thus yields

$$60 = 1 + 20 + 15 + 12 + 12.$$

Since no combination of 1 with any of 20, 15, 12, 12 divides 60 unless it is equal to 1 or 60, it follows that  $A_5$  has no proper normal subgroups, so  $A_5$  is simple.  $\square$

**Theorem:** For all  $n \geq 5$ ,  $A_n$  is simple.

*Proof.* We use induction on  $n \geq 5$ . Set  $G = A_n$ , and let  $G_i \leq A_n$  be the stabilizer of  $i$  in  $A_n$ , so that  $G_i \cong A_{n-1}$ . If  $N \trianglelefteq A_n$  with  $N \neq \{e\}$ , we will show that  $N = A_n$ .

It suffices to show that there exists  $1 \leq i \leq n$  such that  $N \cap G_i \neq \{e\}$ . It then follows that  $N \cap G_i \trianglelefteq G_i$ , so by the induction hypothesis,  $N \cap G_i = G_i$ . Since, for  $\sigma \in A_n$ , we have  $\sigma G_i \sigma^{-1} = G_{\sigma(i)}$ , and since  $N$  is normal,  $G_{\sigma(i)} \leq N$ , whence  $G_1, \dots, G_n \leq N$ . We claim that

$$A_n = \bigcup_{j=1}^n G_j \cdot G_1.$$

If  $g(1) = 1$ , then  $g \in A_n$ , and we are done. Else, if  $g(1) \neq 1$ , we pick  $1 \leq i \leq n$  with  $i \neq 1, g(1)$ , and  $1 \leq j \leq n$  with  $j \neq 1, g(1), i$  (this is why  $n \geq 5$ ). Taking  $h = (1, g(i), i)$ , we have that  $h \in G_j$ , with  $h^{-1}g(1) = 1$ , whence  $g \in G_j \cdot G_1$ .

To prove the theorem, we will construct  $\pi \in G_i \cap N$  with

$$\pi = \sigma^{-1}(\tau^{-1}\sigma\tau),$$

where  $\sigma$  is a fixed element of  $N$  and  $\sigma \neq e$ .

- If  $\sigma$  is a singular 3-cycle, then  $N = A_n$  since  $A_n$  is generated by the 3-cycles.
- Suppose the decomposition of  $\sigma$  contains a cycle of length at least 4. Then, via conjugacy, we may assume that  $\sigma$  is of the form

$$\sigma = (1, 2, \dots, m)\rho,$$

where  $\rho$  is some disjoint collection of cycles. If  $\tau = (1, 2, 3) \in A_5$ , we have that

$$\sigma\tau = (1, 2, \dots, m)\rho(1, 2, 3)$$

takes  $1 \mapsto 3$  and  $m \mapsto 1$ , where  $m \geq 4$ . Meanwhile, we have that

$$\tau\sigma = (1, 2, 3)(1, 2, \dots, m)\rho$$

takes  $1 \mapsto 3$  and  $m \mapsto 2$ . In particular,  $\sigma^{-1}\tau^{-1}\sigma\tau$  takes  $1 \mapsto 1$  and  $m \not\mapsto 1$ . Therefore,  $\pi = \sigma^{-1}\tau^{-1}\sigma\tau$  is contained in  $G_1 \cap N$ , and we are done in this case.

(iii) If  $\sigma$  contains cycle of length at most 3, and  $\sigma$  contains at least two such cycles, then we may write

$$\sigma = (1, 2, 3)(4, 5, 6)\rho$$

by conjugacy. Then, if  $\tau = (1, 2, 4)$ , we have

$$\begin{aligned}\sigma^{-1}\tau^{-1}\sigma\tau &= \rho^{-1}(4, 6, 5)(1, 3, 2)(1, 4, 2)(1, 2, 3)(4, 5, 6)\rho(1, 2, 4) \\ &= (4, 6, 5)(1, 3, 2)(1, 4, 2)(1, 2, 3)(4, 5, 6)(1, 2, 4) \\ &= (1, 4, 2, 6, 3).\end{aligned}$$

Thus,  $\pi \in G_5 \cap N$ .

(iv) If  $\sigma$  contains exactly one such 3-cycle, where

$$\sigma = (1, 2, 3)\rho,$$

with  $\rho$  exclusively consisting of transpositions, then

$$\sigma^2 = (1, 3, 2),$$

meaning that  $N$  contains a singular 3-cycle, so we reduce to case (i).

(v) If  $\sigma$  consists entirely of transpositions, then we may write

$$\sigma = (1, 2)(3, 4)\rho$$

with  $\rho$  consisting of disjoint transpositions. Observe that if  $\tau = (1, 2, 3)$ , then

$$\begin{aligned}\sigma^{-1}\tau^{-1}\sigma\tau &= \tau^{-1}(1, 2)(3, 4)(1, 3, 2)(1, 2)(3, 4)\tau(1, 2, 3) \\ &= (1, 3)(2, 4).\end{aligned}$$

Observe that  $\zeta = (1, 3, 5)$  is in  $A_n$  and satisfies

$$\begin{aligned}\pi(\zeta^{-1}\pi\zeta) &= (1, 3)(2, 4)(1, 5, 3)(1, 3)(2, 4)(1, 3, 5) \\ &= (1, 3, 5).\end{aligned}$$

Thus,  $N$  contains a singular 3-cycle, and we reduce to case (i).

□

## Solvable Groups

We like (finitely generated) abelian groups a lot because we know how to classify them. Specifically, it can be shown that every finitely generated abelian group is of the form

$$G \cong \mathbb{Z}^d \times \mathbb{Z}/p_1^{n_1}\mathbb{Z} \times \cdots \times \mathbb{Z}/p_r^{n_r}\mathbb{Z}$$

for some  $d \geq 0$ ,  $n_1, \dots, n_r > 0$ , and primes  $p_1, \dots, p_r$ . The question then becomes the following: what if our group is non-abelian? What is the next best decomposition of this group?

**Definition:** Let  $G$  be a group. The *commutator subgroup*  $[G, G] \leq G$  is the subgroup generated by the commutators  $xyx^{-1}y^{-1}$  for all  $x, y \in G$ .

We denote  $[G, G] = D^{(1)}(G)$  as the *first derived subgroup* of  $G$ . We iterate  $D^{(n)}(G) = [D^{(n-1)}(G), D^{(n-1)}(G)]$ .

Note that elements  $g \in [G, G]$  are not necessarily commutators themselves, but they are products of commutators.

**Definition:** A group  $G$  is called *solvable* if there is some  $n$  such that  $D^{(n)}(G) = \{e\}$ .

**Lemma:** Let  $G$  be a group. The following hold.

- (i) The commutator subgroup  $[G, G] \trianglelefteq G$  is normal.
- (ii) The quotient  $G/[G, G]$  is abelian.
- (iii) The commutator subgroup is the smallest normal subgroup such that (ii) holds.

**Remark:** Pretty much by definition, if  $\phi: G \rightarrow H$  is a group homomorphism, then  $\phi([x, y]) = [\phi(x), \phi(y)]$ . If  $\phi$  is surjective, then  $\phi([G, G]) = [H, H]$ .

*Proof.*

- (i) Apply the observation to the conjugation automorphism. In particular, for all  $g, x, y \in G$ ,  $g[x, y]g^{-1} = [gxg^{-1}, gyg^{-1}]$ , so that  $g[x, y]g^{-1} \in [G, G]$  for all generators of the commutator subgroup. In particular, this means that  $[G, G]$  is normal.
- (ii) Let  $g, h \in G$ . Observe then that  $gh[G, G] = hg[G, G]$  by definition, since  $ghg^{-1}h^{-1} \in [G, G]$ .
- (iii) Let  $N \trianglelefteq G$  be such that  $G/N$  is abelian. Then,  $ghN = hgN$  for all  $g, h \in G$ , so that  $ghg^{-1}h^{-1} \in N$ . Since  $g$  and  $h$  are arbitrary, it follows that  $[G, G] \subseteq N$ .

□

Note that from this result, we are able to consider a series of subgroups known as the *derived series* of the group, given by

$$G = D^{(0)}(G) \supseteq D^{(1)}(G) \supseteq \dots \supseteq D^{(n)}(G) \supseteq \dots,$$

where we say the group is solvable if the derived series stabilizes at  $\{e\}$ .

**Theorem:** For all  $n \geq 5$ ,  $S_n$  is not solvable.

*Proof.* We see that for all  $n \geq 3$ ,  $A_n$  is generated by the 3-cycles, since we may group pairs of transpositions that have one element in common into a 3-cycle; if we have a cycle of type  $2 + 2$ , conjugated into  $(1, 2)(3, 4)$ , we find that we can find a suitable product of 3-cycles given by  $(1, 3, 2)(1, 3, 4)$ .

Now, since  $S_n/A_n \cong \mathbb{Z}/2\mathbb{Z}$  for all  $n$ , we must have  $[S_n, S_n] \leq A_n$ . Yet, since  $[S_n, S_n] \trianglelefteq A_n$ , we have  $[S_n, S_n] \leq A_n$ . For all  $n \geq 5$ , we  $A_n$  is simple, so that  $[S_n, S_n] = A_n$ .

Finally, since  $A_n$  for all  $n \geq 5$  does not have any proper normal subgroups, and  $A_n$  is not commutative, it must be the case that  $[A_n, A_n] = A_n$ , meaning that the derived series does not stabilize at  $\{e\}$ . □

Solvability is useful because it has a similar inheritance property to Noetherian-ness in the case of modules.

**Proposition:** Let  $G$  be a solvable group.

- (i) Every subgroup of  $G$  is solvable.
- (ii) Every quotient of  $G$  is solvable.
- (iii) If  $N \trianglelefteq G$  and  $G/N$  are solvable, then  $G$  is solvable.

*Proof.*

- (i) By definition,  $D^{(n)}(H) \leq D^{(n)}(G)$ .
- (ii) Using the observation with the canonical projection  $\pi: G \rightarrow G/N$ , we find that  $D^{(n)}(G/N) = D^{(n)}(G)/N$ .

- (iii) Let  $\pi: G \rightarrow G/N$  be the canonical projection. Since  $G/N$  is solvable, there is some index  $n$  such that  $D^{(n)}(G)/N = D^{(n)}(G/N) = \{e\}$ . Therefore,  $D^{(n)}(G) \leq N$ . Since  $N$  is solvable, there is an index  $m$  such that  $D^{(m)}(N) = \{e\}$ . Thus,  $D^{(m+n)}(G) \leq D^{(m)}(N) = \{e\}$ .

□

**Corollary:** Every finite  $p$ -group is solvable.

*Proof.* Since  $|G| = p^r$  for some  $r \geq 1$ , we know a few cases. If  $r = 1$ , then  $G = \mathbb{Z}/p\mathbb{Z}$ , and  $G$  is then abelian.

If  $r > 1$ , then  $Z(G)$  is nontrivial, and if  $Z(G) = G$ ,  $G$  is abelian and we are done yet again. Else, both  $G/Z(G)$  and  $Z(G)$  are nontrivial  $p$ -groups with order less than  $p^r$ , so the induction hypothesis and the previous proposition give us our desired outcome. □

**Corollary:** If  $G$  is a group of order  $pq$  or order  $p^2q$  with  $p$  and  $q$  distinct primes, then  $G$  is solvable.

*Proof.* We have that  $G$  has some normal  $p$ -subgroup (or  $q$ -subgroup) from the Sylow theorem, so we may use the previous proposition. □

**Definition:** Let  $G$  be a group. A *subnormal* series for  $G$  is a sequence of subgroups

$$G = G_0 \geq G_1 \geq \cdots \geq G_n \geq \cdots$$

with  $G_{i+1} \trianglelefteq G_i$ . We call the quotients  $G_i/G_{i+1}$  the *factors* of the series. If each  $G_i \trianglelefteq G$  is normal, then we call the series a normal series.

**Proposition:** A group  $G$  is solvable if and only if  $G$  admits a subnormal series terminating at the identity,

$$G = G_0 \triangleright G_1 \triangleright \cdots \triangleright G_n = \{e\},$$

with all factors abelian.

*Proof.* In the forward direction, the derived subgroup series satisfies the condition.

Now, suppose  $G$  admits such a series. Since  $G_n = \{e\}$ , and  $G_{n-1} \cong G_{n-1}/G_n$  is abelian,  $G_{n-1}$  is solvable. Since  $G_{n-2}$  has a solvable normal subgroup and solvable quotient, it follows that  $G_{n-2}$  is solvable. Continuing up the chain gives the desired result. □

## Nilpotent Groups

**Proposition:** Let  $K \trianglelefteq G$  be a subgroup, and let  $K \leq H \leq G$  be a series of subgroups. Then,  $[H, G] \leq K$  if and only if  $H/K \leq Z(G/K)$ .

*Proof.* Let  $h \in H$  and  $g \in G$ . Then,  $[h, g] \in K$  if and only if  $[h, g]K = K$ , if and only if  $hKgK = gKhK$ , whence  $H/K \leq Z(G/K)$ . □

**Definition:** Let  $G$  be a group. Inductively, define

$$\begin{aligned} G_0 &= G \\ G_{i+1} &= [G_i, G]. \end{aligned}$$

The series of normal subgroups

$$G \triangleright G_1 \triangleright \cdots$$

is known as the *lower central series* of  $G$ .

**Definition:** Let  $G$  be a group. Inductively, define

$$\begin{aligned} Z_0(G) &= e \\ Z_{i+1}(G)/Z_i(G) &= Z(G/Z_i(G)). \end{aligned}$$

The series of normal subgroups

$$Z_0(G) \trianglelefteq Z_1(G) \trianglelefteq \cdots$$

is known as the *upper central series* of  $G$ .

We call a group  $G$  *nilpotent* if  $Z_n(G) = G$  for some  $n$ .

**Proposition:** The lower central series terminates if and only if the upper central series terminates. In particular, if there is  $n$  such that  $Z_n(G) = G$ , then  $G_{n+1} = e$ . Moreover,  $G_{i+1} \leq Z_{n-i}(G)$  for all  $i$ .

*Proof.* Suppose  $Z_n(G) = G$ . If  $i = 0$ , then  $G_1 = G = Z_n(G)$ , so the base case is proven.

If  $G_{i+1} \leq Z_{n-i}(G)$ , then

$$\begin{aligned} G_{i+2} &= [G_{i+1}, G] \\ &\leq [Z_{n-i}(G), G] \\ &\leq Z_{n-i-1}(G). \end{aligned}$$

Therefore, in particular, we have that  $G_{n+1} \leq Z_0(G) = e$ .

Now, suppose  $G_{n+1} = e$ . We will show that  $G_{n+1-j} \leq Z_j(G)$ . If  $j = 0$ , then  $Z_0(G) = G_{n+1} = 1$ .

If  $G_{n+1-j} \leq Z_j(G)$ , then by the third isomorphism theorem, we have the surjective homomorphism  $G/G_{n+1-j} \rightarrow G/Z_j(G)$ . Since  $[G_{n-j}, G] = G_{n+1-j}$ , we have from the proposition that  $G_{n-j}/G_{n+1-j} \leq Z(G/G_{n+1-j})$ .

From the surjection, we have

$$\begin{aligned} G_{n-j}Z_j(G)/Z_j(G) &\leq Z(G/Z_j(G)) \\ &= Z_{j+1}(G)/Z_j(G), \end{aligned}$$

whence  $G_{n-j} \leq G_{n-j}Z_j(G) \leq Z_{j+1}(G)$ . In particular, we have that  $G_1 = G = Z_n(G)$ . □

**Proposition:** A group  $G$  is nilpotent if and only if  $G/Z(G)$  is nilpotent.

*Proof.* Suppose  $G$  is nilpotent. Then, there is  $N$  such that

$$\begin{aligned} G_N &= [G_{N-1}, G] \\ &= \{e\}. \end{aligned}$$

Now, if  $\pi: G \rightarrow G/Z(G)$  is the canonical projection, then from the (implicit) induction hypothesis,

$$\begin{aligned} (G/Z(G))_n &= [(G/Z(G))_{n-1}, G/Z(G)] \\ &= [\pi(G_{n-1}), \pi(G)] \\ &= \pi([G_{n-1}, G]) \\ &= \pi(\{e\}) \\ &= \{e\}. \end{aligned}$$

Therefore,  $G/Z(G)$  is nilpotent.

Now if  $G/Z(G)$  is nilpotent, we have

$$(G/Z(G))_n = \{e\}$$

for some  $n$ . As above, if  $\pi: G \rightarrow Z(G)$  is the canonical projection, we have

$$\begin{aligned} \pi(G_n) &= (G/Z(G))_n \\ &= \{e\}, \end{aligned}$$

or that  $G_n \leq Z(G)$ . Thus, we have

$$\begin{aligned} G_{n+1} &= [G_n, G] \\ &\leq [Z(G), G] \\ &= \{e\}, \end{aligned}$$

whence  $G$  is nilpotent.  $\square$

**Theorem:** Every finite  $p$ -group is nilpotent.

*Proof.* Every finite  $p$ -group has nontrivial center. If, for some  $j$ , we have  $Z_j(G) < G$ , then  $Z(G/Z_j(G)) \neq e$ , so  $Z_j(G) < Z_{j+1}(G)$ . Since  $G$  is finite, there is some  $m$  such that  $Z_m(G) = G$ , whence  $G$  is nilpotent.  $\square$

**Theorem:** Every nilpotent group is solvable.

*Proof.* We see that  $D^{(1)}(G) \leq G$ , and inductively,

$$\begin{aligned} D^{(i+1)}(G) &= [D^{(i)}(G), D^{(i)}(G)] \\ &\leq [D^{(i)}(G), G_i] \\ &\leq [G_i, G_i] \\ &\leq [G_i, G] \\ &= G_{i+1}. \end{aligned}$$

Thus, since the lower central series terminates, so too does the derived series.  $\square$

## Free Groups

Given a set  $X$ , we will construct a group  $F(X)$  satisfying the following universal property: given any set map  $\phi: X \rightarrow G$  where  $G$  is a group, then there is a unique homomorphism  $\varphi: F(X) \rightarrow G$  such that the following diagram commutes.

$$\begin{array}{ccc} X & \xrightarrow{\phi} & G \\ \downarrow \iota & \nearrow \varphi & \\ F(X) & & \end{array}$$

To do this, we start by constructing a set  $X^{-1}$  that is a copy of  $X$ , which will be our “inverses,” and we introduce an extra element  $e$  with  $e^{-1} = e$ . From now on, we will call the identity element 1. Informally, we will treat  $X \cup X^{-1}$  as our “alphabet,” where the elements of  $X$  are letters. We will define  $F(X)$  to be the set of words on  $X$  subject to an equivalence relation.

A word on  $X$  is an expression  $w = x_1^{\pm 1} \cdots x_n^{\pm 1}$ , where the  $x_i \in X$  are not necessarily distinct. The set of words on the alphabet  $X$  is denoted  $W(X)$ . We will define two equivalence relations on  $W(X)$ .

(i) We allow for “insertion” of a pair  $xx^{-1}$ , so that

$$x_1x_2 \cdots x_n \sim x_1x_2 \cdots x_ixx^{-1}x_{i+1} \cdots x_n$$

for any  $x \in X \cup X^{-1}$ .

(ii) We allow for “deletion” of a pair  $xx^{-1}$ , so that

$$\begin{aligned} x_1x_2 \cdots x_ixx^{-1}x_{i+1} \cdots x_n &\sim x_1x_2 \cdots x_n \\ &\sim x_1x_2 \cdots x_i1x_{i+1} \cdots x_n. \end{aligned}$$

**Definition:** The free group  $F(X)$  is defined by

$$F(X) = W(X) / \sim,$$

where the operation is concatenation:

$$(x_1 \cdots x_n)(y_1 \cdots y_m) = x_1 \cdots x_n y_1 \cdots y_m.$$

We say a word  $w \in F(X)$  is *reduced* if there are no instances of 1 in the expression for  $w$ , and for any  $x_i \neq 1$ , we have  $x_{i+1} \neq x_i^{-1}$ .

From the definition of our equivalence relation, every word is equivalent to a reduced word.

**Proposition:** The free group  $F(X)$  satisfies the universal property.

*Proof.* Let  $\phi: X \rightarrow G$  be a set map. Extend  $\phi$  to  $\Phi: W(X) \rightarrow G$  by taking

$$\begin{aligned}\Phi(x^{-1}) &= \Phi(x)^{-1} \\ \Phi(w) &= \Phi(x_1)^{\varepsilon_1} \cdots \Phi(x_m)^{\varepsilon_m}.\end{aligned}$$

Then,  $\Phi$  factors through  $\sigma$ , giving  $\varphi: F(X) \rightarrow G$ .

By definition,  $\varphi$  is a homomorphism with  $\varphi|_X = \phi$ . As for uniqueness, if  $\psi: F(X) \rightarrow G$  is another homomorphism such that  $\psi|_X = \phi$ , then

$$\begin{aligned}\psi(w) &= \psi(x_1) \cdots \psi(x_m) \\ &= \phi(x_1) \cdots \phi(x_m) \\ &= \varphi(w).\end{aligned}$$

□

**Corollary:** Every group is a quotient of a free group.

**Definition:** If  $\pi: F(X) \rightarrow G$  is such a surjective homomorphism, then the subgroup  $R = \ker(\pi)$  is called a subgroup of *relations* of  $G$ .

Note that the relations subgroup is not unique, since we may select any generating set for  $G$ . We will write

$$G = \langle X | R \rangle,$$

where  $X$  is a generating set for  $G$  and  $R$  is the subgroup of relations. We call this a presentation for  $G$  in terms of the generators  $X$  and relations  $R$ .