

Observations on Excess Area Identities and Operator Symbols in Bergman Spaces

Avinash Iyer

Occidental College

Summary

- 1 Definitions and Notations
- 2 Motivation and Problem
- 3 Results and Observations
- 4 Remarks and Future Directions
- 5 REU Experience
- 6 Acknowledgements and References

Contents

- 1 Definitions and Notations
- 2 Motivation and Problem
- 3 Results and Observations
- 4 Remarks and Future Directions
- 5 REU Experience
- 6 Acknowledgements and References

Definition and Notations

- Ω : a region in \mathbb{C} e.g. $\mathbb{D}, D(0, r), \mathbb{A}(0, r, 1), \mathbb{C}$
- $\lambda(z) = \lambda(|z|) \in C^\infty(\Omega)$: weight function

Definition (λ -weighted Square-Integrable Functions)

$$L^2(\Omega, \lambda) = \left\{ f : \Omega \rightarrow \mathbb{C} \left| \int_{\Omega} |f(z)|^2 \lambda(z) \, dA(z) < \infty \right. \right\}$$

- $L^2(\Omega, \lambda)$ forms a Hilbert space with inner product

$$\langle f, g \rangle_{L^2(\Omega, \lambda)} = \int_{\Omega} f(z) \overline{g(z)} \lambda(z) \, dA(z)$$

inducing the norm

$$\|f\|_{L^2(\Omega, \lambda)}^2 = \int_{\Omega} |f(z)|^2 \lambda(z) \, dA(z)$$

Definitions and Notations

Definition (Holomorphic Function on Ω)

We say h is holomorphic on Ω , or $h \in \mathcal{O}(\Omega)$, if, for all $z \in \Omega$

$$\begin{aligned}\frac{\partial h(z)}{\partial \bar{z}} &= \frac{1}{2} \left(\frac{\partial h(z)}{\partial x} + i \frac{\partial h(z)}{\partial y} \right) \\ &= \frac{1}{2} \left(\left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + i \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right) \right) \\ &= 0.\end{aligned}$$

Definition (λ -weighted Bergman Space)

$$A^2(\Omega, \lambda) := \mathcal{O}(\Omega) \cap L^2(\Omega, \lambda).$$

Definitions and Notations

Definition ($A^{1,2}(\Omega, \lambda)$)

$$A^{1,2}(\Omega, \lambda) = \left\{ h \in A^2(\Omega, \lambda) \mid \frac{\partial h}{\partial z} \in A^2(\Omega, \lambda) \right\}$$

Definition (Weighted Image-Area)

Let $h \in A^{1,2}(\Omega, \lambda)$.

$$\begin{aligned} A_{\Omega, \lambda}(h) &= \int_{\Omega} \left| \frac{\partial h}{\partial z} \right|^2 \lambda(z) \, dA(z) \\ &= \left\| \frac{\partial h}{\partial z} \right\|_{L^2(\Omega, \lambda)}^2 \end{aligned}$$

Definitions and Notations

- $A^2(\Omega, \lambda)$ has a reproducing kernel i.e $\exists! K_{\Omega}^{\lambda}(\cdot, z) \in A^2(\Omega, \lambda) :$

$$h(z) = \langle h(\cdot), K_{\Omega}^{\lambda}(\cdot, z) \rangle_{L^2(\Omega, \lambda)}$$

- $A^2(\Omega, \lambda)$ is a closed subspace of $L^2(\Omega, \lambda)$.

Definition (Bergman Projection)

Let $P^{\Omega, \lambda} : L^2(\Omega, \lambda) \rightarrow A^2(\Omega, \lambda)$

$$\begin{aligned} (P^{\Omega, \lambda} h)(z) &:= \langle h(\cdot), K_{\Omega}^{\lambda}(\cdot, z) \rangle_{L^2(\Omega, \lambda)} \\ &= \int_{\Omega} h(w) \overline{K_{\Omega}^{\lambda}(w, z)} \lambda(w) \, dA(w) \end{aligned}$$

Definitions and Notations

Definition (Multiplication Operator)

Let $M_\varphi : L^2(\Omega, \lambda) \rightarrow L^2(\Omega, \lambda)$ where $\varphi \in L^\infty(\Omega)$

$$M_\varphi(h) := \varphi h$$

Definition (Toeplitz Operator)

$T_\varphi^{\Omega, \lambda} : A^2(\Omega, \lambda) \rightarrow A^2(\Omega, \lambda)$, where $\varphi \in L^\infty(\Omega)$

$$T_\varphi^{\Omega, \lambda} := P^{\Omega, \lambda} M_\varphi$$

Definitions and Notations

Definition (Commutator)

Let $[P^{\Omega,\lambda}, M_\varphi] : L^2(\Omega, \lambda) \rightarrow L^2(\Omega, \lambda)$

$$[P^{\Omega,\lambda}, M_\varphi] := P^{\Omega,\lambda} M_\varphi - M_\varphi P^{\Omega,\lambda}$$

Definition (Hankel Operator)

Let $H_\varphi^{\Omega,\lambda} : A^2(\Omega, \lambda) \rightarrow (A^2(\Omega, \lambda))^\perp$

$$\begin{aligned} H_\varphi^{\Omega,\lambda} &:= - [P^{\Omega,\lambda}, M_\varphi] \Big|_{A^2(\Omega,\lambda)} \\ &= (I - P^{\Omega,\lambda}) M_\varphi \\ &= M_\varphi - P^{\Omega,\lambda} M_\varphi \\ &= M_\varphi - T_\varphi^{\Omega,\lambda} \end{aligned}$$

Contents

- 1 Definitions and Notations
- 2 Motivation and Problem**
- 3 Results and Observations
- 4 Remarks and Future Directions
- 5 REU Experience
- 6 Acknowledgements and References

Motivations

- $\{z^n\}_{n=0}^\infty$ form a *complete orthogonal basis* for $A^2(\mathbb{D})$
- If h is holomorphic, then h is analytic:

$$h(z) = \sum_{n=0}^{\infty} h_n z^n$$

and

$$S_N := \sum_{n=0}^N h_n z^n$$

converges uniformly on compact subsets.

- Relationship between L^2 norm of h to the ℓ^2 norm of $\{h_k\}_{k=0}^\infty$:

$$\|h\|_{L^2(\mathbb{D})}^2 = \int_{\mathbb{D}} |h(z)|^2 dA(z) = \pi \sum_{k=0}^{\infty} \frac{|h_k|^2}{k+1}$$

- $[T_{\bar{z}}^{\mathbb{D}} M_z, D M_z](z^m) = 0$

- How can we expand **established identities concerning the area of the image of domains** under a holomorphic map in different Bergman spaces?
- Can we study the **structural properties of integral operators** (such as Toeplitz and Hankel operators) using the properties of Bergman spaces?

Literature Review on Previous Results I

- D'Angelo's Excess Area identity [D'A19]

Let $h \in A^{1,2}(\mathbb{D})$. Then,

$$\begin{aligned} A_{\mathbb{D}}(zh) - A_{\mathbb{D}}(h) &= \left\| \frac{\partial(zh)}{\partial z} \right\|_{L^2(\mathbb{D})}^2 - \left\| \frac{\partial h}{\partial z} \right\|_{L^2(\mathbb{D})}^2 \\ &= \frac{1}{2} \int_0^{2\pi} |f(e^{i\theta})|^2 d\theta \\ &= \pi \|Sh\|_{L^2(\partial\mathbb{D})}^2 \end{aligned}$$

where Sh is the restriction of h to the unit circle.

Literature Review on Previous Results II

- Excess Area identity with Blaschke product multiplier
- 'Excess Area' identity for harmonic functions [BÇGH22]
- Generating symbols for Toeplitz operators for a given initial p and target polynomial q on unit disc and polydisc, $T_{\varphi}^{\mathbb{D}}(p) = q$ and $T_{\varphi}^{\mathbb{D}^n}(p) = q$ [ÇDTR⁺24]
- Substituted derivatives for Toeplitz operators in Excess Area identity [ÇDTR⁺24]

Contents

- 1 Definitions and Notations
- 2 Motivation and Problem
- 3 Results and Observations**
- 4 Remarks and Future Directions
- 5 REU Experience
- 6 Acknowledgements and References

Summary of Results

1. Results and Observations influenced by the Area Difference of the image of \mathbb{D} between zh and h :
 - i. On $\mathcal{F}^2 = A^2(\mathbb{C}, e^{-|z|^2}), A^2(\mathbb{D}, \lambda), A^2(D(0, r))$
 - ii. On convergence of identities on certain weighted discs.
- 2 Results and Observations influenced by symbol-generating algorithm for Toeplitz Operators
 - i. On unweighted and weighted Toeplitz operators relation
 - ii. On creating symbols for Unweighted and weighted Hankel operators and commutator operators on $A^2(\mathbb{D})$

Methods Used

- Relation between L^2 norms of functions and ℓ^2 norms of Taylor series:

$$\|h\|_{L^2(\mathbb{D})}^2 = \sum_{k=0}^{\infty} \frac{|h_k|^2}{k+1}$$

- Integration by parts via Stokes's theorem on forms:

$$\begin{aligned}\oint_{\partial\Omega} f \, dz &= \int_{\Omega} \overline{\frac{\partial f}{\partial \bar{z}}} \, d\bar{z} \wedge dz \\ \oint_{\partial\Omega} f \, d\bar{z} &= \int_{\Omega} \frac{\partial f}{\partial z} \, dz \wedge d\bar{z}.\end{aligned}$$

- Inequalities e.g. Cauchy-Schwarz inequality, Hölder's inequality
- Beta, Gamma, and Hypergeometric functions

Using Integration by Parts to find Excess Area identity: Wedge Product I

The area is integrated with respect to $dA = dx \wedge dy$. The wedge product has the following properties:

$$(a + b) \wedge c = a \wedge c + b \wedge c$$

$$a \wedge b = -b \wedge a$$

$$a \wedge a = 0.$$

With $z = x + iy$, $\bar{z} = x - iy$, the substitution $x = \frac{z+\bar{z}}{2}$, $y = \frac{z-\bar{z}}{2i}$ yields

$$\begin{aligned} dx \wedge dy &= \frac{1}{2i} (d\bar{z} \wedge dz) \\ &= -\frac{1}{2i} (dz \wedge d\bar{z}). \end{aligned}$$

Using Integration by Parts to find Excess Area identity: Wedge Product II

The area integral is now rewritten as:

$$\begin{aligned}\left\langle \frac{\partial h}{\partial z}, \frac{\partial h}{\partial \bar{z}} \right\rangle_{L^2(\Omega, \lambda)} &= \int_{\Omega} \left(\overline{\frac{\partial h}{\partial z}} \right) \left(\frac{\partial h}{\partial \bar{z}} \right) \lambda(|z|) \, dx \wedge dy \\ &= \frac{1}{2i} \int_{\Omega} \lambda(|z|) \left(\left(\overline{\frac{\partial h}{\partial z}} \right) d\bar{z} \right) \wedge \left(\left(\frac{\partial h}{\partial \bar{z}} \right) dz \right)\end{aligned}$$

Using Integration by Parts to find Excess Area identity: Stokes's Theorem I

In particular,

$$\frac{\overline{\partial}}{\partial z} \left((\lambda(|z|)) \bar{h} \frac{\partial h}{\partial z} \right) d\bar{z} \wedge dz = \underbrace{(\lambda(|z|)) \frac{\overline{\partial} h}{\partial z} d\bar{z} \wedge \frac{\partial h}{\partial z} dz}_{\text{area integrand}} + \left(\frac{\overline{\partial}}{\partial z} \lambda(|z|) \right) \bar{h} \wedge \frac{\partial h}{\partial z} dz$$

meaning

$$\begin{aligned} \frac{1}{2i} \int_{\Omega} \frac{\partial h}{\partial z} \frac{\overline{\partial} h}{\partial z} \lambda(|z|) d\bar{z} \wedge dz &= \underbrace{\frac{1}{2i} \int_{\Omega} \frac{\overline{\partial}}{\partial z} \left(\lambda(|z|) \bar{h} \frac{\partial h}{\partial z} \right) d\bar{z} \wedge dz}_{\text{Integral A}} \\ &\quad - \frac{1}{2i} \int_{\Omega} \bar{h} \frac{\partial h}{\partial z} \left(\frac{\overline{\partial}}{\partial z} \lambda(|z|) \right) d\bar{z} \wedge dz. \end{aligned}$$

Using Integration by Parts to find Excess Area identity: Stokes's Theorem II

Turning our attention to Integral A,

$$\begin{aligned}\frac{1}{2i} &= \int_{\Omega} d \left(\lambda(|z|) \bar{h} \frac{\partial h}{\partial z} \right) d\bar{z} \wedge dz \\ &= \underbrace{\int_{\partial\Omega} \lambda(|z|) \bar{h} \frac{\partial h}{\partial z} dz}_{=0}.\end{aligned}$$

With this, the area integral is now

$$\frac{1}{2i} \int \frac{\partial h}{\partial z} \frac{\partial \bar{h}}{\partial \bar{z}} \lambda(|z|) d\bar{z} \wedge dz = -\frac{1}{2i} \int \bar{h} \frac{\partial h}{\partial z} \left(\frac{\partial}{\partial \bar{z}} \lambda(|z|) \right) d\bar{z} \wedge dz$$

Excess Area on Fock Spaces

D'Angelo's Excess Area identity:

$$A_{\mathbb{D}}(zh) - A_{\mathbb{D}}(h) = \frac{1}{2} \int_0^{2\pi} |f(e^{i\theta})|^2 d\theta = \pi \|Sh\|_{L^2(\mathbb{bD})}^2$$

Excess Area on Fock Space

Given $h \in \mathcal{F}^2$ with $\frac{\partial h}{\partial \bar{z}}$,

$$\begin{aligned} & A_{\mathcal{F}^2}(zh) - A_{\mathcal{F}^2}(h) \\ &= \pi \left\| z T_{\bar{z}}^{\mathcal{F}^2}(h) \right\|_{\mathcal{F}^2}^2 + \pi \left\| T_{\bar{z}}^{\mathcal{F}^2}(h) \right\|_{\mathcal{F}^2}^2 + \pi \left\| H_{\bar{z}}^{\mathcal{F}^2}(h) \right\|_{\mathcal{F}^2}^2 \end{aligned}$$

Here, the restriction of h to the unit circle in D'Angelo's Excess Area identity is replaced with the Bergman projection on \mathbb{C} .

Excess Area on $A^2(\mathbb{D}, \lambda)$

D'Angelo's Excess Area identity:

$$A_{\mathbb{D}}(zh) - A_{\mathbb{D}}(h) = \frac{1}{2} \int_0^{2\pi} |f(e^{i\theta})|^2 d\theta = \pi \|Sh\|_{L^2(\partial\mathbb{D})}^2$$

Excess Area on $A^2(\mathbb{D}, \lambda)$

Let $h \in A^{1,2}(\mathbb{D}, \lambda)$, $\lambda(z) = 1 - |z|^2$. Then,

$$A_{\mathbb{D}, \lambda}(z^{m+1}h) - A_{\mathbb{D}, \lambda}(z^m h) = \pi \|z^m h\|_{L^2(\mathbb{D}, \lambda)}^2.$$

Here, the restriction of h to the unit circle is replaced with the function itself.

Dilation and Contraction from $A^2(D(0, r))$ to $A^2(\mathbb{D})$

Contracting $h \in A^{1,2}(\mathbb{D})$ by taking $h_r = h(rz)$ for some $0 < r < 1$,

$$A_{\mathbb{D}}(zh_r) - A_{\mathbb{D}}(h_r) = \pi \|Sh_r\|_{L^2(b\mathbb{D})}^2 \quad (1)$$

$$A_{D(0,r)}(zh) - r^2 A_{D(0,r)}(h) = \pi r^2 \|Sh\|_{L^2(bD(0,r))}^2. \quad (2)$$

Dilating $h \in A^{1,2}(D(0, r))$ by taking $h_{\frac{1}{r}} = h(\frac{z}{r})$ for some $0 < r < 1$

$$A_{D(0,r)}(zh_{1/r}) - r^2 A_{D(0,r)}(h_{1/r}) = \pi r^2 \|Sh_{1/r}\|_{L^2(bD(0,r))}^2 \quad (3)$$

$$A_{\mathbb{D}}(zh) - A_{\mathbb{D}}(h) = \pi \|Sh\|_{L^2(b\mathbb{D})}^2 \quad (4)$$

Approximation for Sequences of Berezin Transform

Weighted Area on $D(0, r)$

Let $\lambda_r(z) = \chi_{D(0,r)} \left(1 - \frac{|z|^2}{r^2}\right)^{r^2}$ where $r > 0$. Then,

$$A_{D(0,r),\lambda_r}(h) = \int_{D(0,r)} |h'(z)|^2 \left(1 - \frac{|z|^2}{r^2}\right)^{r^2} dA(z)$$

We find that, as $r \rightarrow \infty$, $A_{D(0,r),\lambda_r}(h) \rightarrow A_{\mathcal{F}^2}(h)$.

Separately,

$$A_{\mathcal{F}^2}(h_\rho) = \left\| T_{\frac{\mathcal{F}^2}{z}} h_\rho \right\|_{\mathcal{F}^2}^2$$

Berezin Transform Convergence, Cont'd

Reproducing Kernel on $A^2(D(0, r), \lambda_r)$

$$\begin{aligned} K_{D(0, r)}^{\lambda_r}(w, z) &= \sum_{k=0}^{\infty} \frac{\bar{z}^k w^k}{\|w^k\|_{L^2(D(0, r), \lambda_r)}^2} \\ &= \frac{1}{\left(1 - \frac{\bar{z}w}{r^2}\right)^{r^2+2}} \end{aligned}$$

Reproducing Kernel on Fock Space

$$K_{\mathcal{F}^2}(w, z) = e^{\bar{z}w}$$

Berezin Transform Convergence, Cont'd

Definition (Berezin Transform ([Zhu07]))

Let

$$k_z^{\Omega, \lambda}(w) := \frac{K_{\Omega}^{\lambda}(w, z)}{\sqrt{K_{\Omega}^{\lambda}(z, z)}}$$

Then, for some bounded operator T on $L^2(\Omega, \lambda)$, define $\mathcal{B}^{\Omega, \lambda} : B(L^2(\Omega, \lambda)) \rightarrow L^2(\Omega, \lambda)$

$$(\mathcal{B}^{\Omega, \lambda} T)(z) := \left\langle T k_z^{\Omega, \lambda}, k_z^{\Omega, \lambda} \right\rangle_{L^2(\Omega, \lambda)}$$

Berezin Transform Convergence, Cont'd

Previous results:

- For $\varphi \in L^\infty(\Omega, \lambda)$, $\mathcal{B}^{\Omega, \lambda} T_\varphi = \mathcal{B}^{\Omega, \lambda} M_\varphi$. (see [AZ98a]).
- φ is harmonic if and only if $\mathcal{B}^{\Omega, \lambda} M_\varphi = \varphi$ (proof in [Eng94]).

New results:

- For $T_\varphi^{D(0,r), \lambda_r} = P^{D(0,r), \lambda_r} M_\varphi$, the Berezin transform $\mathcal{B}^{D(0,r), \lambda_r} T_\varphi^{D(0,r), \lambda_r}$ converges pointwise to $\mathcal{B}^{\mathcal{F}^2} T_\varphi^{\mathcal{F}^2}$ as $r \rightarrow \infty$.
- By Dini's Theorem, this convergence is uniform on compact subsets of \mathbb{C} (proof inspired by [GS20]).

Unweighted and Weighted Toeplitz Operators

Relation I

Using an extension of [ÇDTR⁺24, Lemma 2.1]

For weight $\lambda(z) = (1 - |z|^2)^\alpha$ ($\alpha \geq 0$) on the unit disc,
 $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$:

$$\frac{T_{\bar{z}^m}^{\mathbb{D}, \lambda_\alpha}(z^n)}{T_{\bar{z}^m}^{\mathbb{D}}(z^n)} = \begin{cases} \frac{\Gamma(m-n+\alpha-2)\Gamma(n+1)(m+1)}{\Gamma(m-n+2)\Gamma(n+\alpha+2)} & \text{if } m \leq n \\ \text{indeterminate} & \text{else} \end{cases}$$

$$T_{\bar{z}^m}^{\mathbb{D}, \lambda_\alpha}(z^n) = s_{n,m,\alpha} T_{\bar{z}^m}^{\mathbb{D}}(z^n), \text{ and } \lim_{n \rightarrow \infty} s_{n,m,\alpha} = 1$$

Unweighted and Weighted Commutator on $A^2(\mathbb{D})$

Existence of Commutator Symbols

Given p and q are harmonic polynomials and $\frac{\partial}{\partial \bar{z}}(p) \neq 0$, there does not exist a polynomial symbol ϕ , such that $[p^{\mathbb{D}}, M_{\phi}](p) = q$ or $[p^{\mathbb{D}, \lambda}, M_{\phi}](p) = q$.

Compare to [ÇDTR⁺24], who worked on constructing Toeplitz symbols mapping between holomorphic polynomials.

Unweighted and Weighted Hankel Operator on $A^2(\mathbb{D})$

Existence of Hankel Operator Symbols

Given some holomorphic polynomials p, q where p is not constant, there does not exist a polynomial symbol ϕ such that $H_{\phi}^{\mathbb{D}}(p) = \overline{q}$ or $H_{\phi}^{\mathbb{D}, \lambda}(p) = \overline{q}$

Contents

- 1 Definitions and Notations
- 2 Motivation and Problem
- 3 Results and Observations
- 4 Remarks and Future Directions**
- 5 REU Experience
- 6 Acknowledgements and References

Remarks on the Annulus

Toeplitz Operator on Monomials on $A^2(\mathbb{A}(0, r, 1))$

For all integers m and n ,

$$T_{\bar{z}^m}^{A(0,r,1)}(z^n) = \begin{cases} \frac{2m r^{2m} \ln(r)}{(r^{2m}-1)} z^{-m-1} & \text{if } n = -1 \\ \frac{r^{2m}-1}{2m \ln(r)} z^{-1} & \text{if } n = m-1 . \\ \frac{(n-m+1)(1-r^{2n+2})}{(n+1)(1-r^{2n-2m+2})} z^{n-m} & \text{else} \end{cases}$$

We attempted to find an algorithm to generate $\varphi \in L^\infty(\mathbb{A}(0, r, 1))$ such that $T_\varphi^{A(0,r,1)}(p) = q$ for given holomorphic Laurent polynomials p and q , but could not prove lack of existence if p has roots inside $\overline{\mathbb{A}(r, 0, 1)}$.

Future Directions

- Existence (or lack thereof) of bounded symbols for Toeplitz operators for a given initial polynomial p and target polynomial q on $\mathbb{A}(0, r, 1)$, $T_{\varphi}^{\mathbb{A}(0, r, 1)}(p) = q$
- Extension of 'Excess Area' identity to harmonic functions in $L^2(\mathbb{C}, e^{-|z|^2})$.
- Connection between non-weighted and weighted Toeplitz operators when the weight is exponential,
 $(1 - |z|^2)^A e^{\frac{-B}{(1-|z|^2)^\alpha}} (A \geq 0, B > 0, \alpha > 0)$.

Contents

- 1 Definitions and Notations
- 2 Motivation and Problem
- 3 Results and Observations
- 4 Remarks and Future Directions
- 5 REU Experience**
- 6 Acknowledgements and References

What is Research Like?

- The complex analysis group consisted of myself, Jennifer Yuan (NYU Abu Dhabi), and Sakia Akamah (Rose–Hulman Institute of Technology).
- Weeks were 9am to 5pm, mostly doing various calculations and updating our collected results document.
- We did not fully understand what we were doing a lot of the time.

Contents

- 1 Definitions and Notations
- 2 Motivation and Problem
- 3 Results and Observations
- 4 Remarks and Future Directions
- 5 REU Experience
- 6 Acknowledgements and References**

Acknowledgments

This work was only possible by the guidance of Dr. Çelik.

We would also like to express our gratitude to Texas A&M-Commerce and its Mathematics Department for hosting the REU where we conducted this research.

This research is based upon work supported by the National Science Foundation under Grant DMS-2243991.

This presentation was originally given August 2, 2024 at the Texas A&M University-Commerce. The presentation was prepared by Sakia Akamah, Avinash Iyer, and Jennifer Yuan.

References I



Lars Ahlfors.

Complex analysis—an introduction to the theory of analytic functions of one complex variable.

2021.



Stephane Attal.

Lectures in Quantum Noise Theory, Open Access Book.



Sheldon Axler and Željko Čučković.

Commuting Toeplitz operators with harmonic symbols.

Integral Equations Operator Theory, 14(1):1–12, 1991.



Sheldon Axler and Dechao Zheng.

The Berezin transform on the Toeplitz algebra.

Studia Math., 127(2):113–136, 1998.

References II



Sheldon Axler and Dechao Zheng.

Compact operators via the Berezin transform.

Indiana Univ. Math. J., 47(2):387–400, 1998.



Haley K. Bambico, Mehmet Çelik, Sarah T. Gross, and Francis Hall.

Generalization of the excess area and its geometric interpretation.

Zbl, 28, 2022.



Steven R. Bell.

The Cauchy transform, potential theory and conformal mapping.

Chapman & Hall/CRC, Boca Raton, FL, second edition, 2016.



F. A. Berezin.

Covariant and contravariant symbols of operators.

Izv. Akad. Nauk SSSR Ser. Mat., 36:1134–1167, 1972.

References III



Joaquin Bustoz and Mourad E. H. Ismail.

On gamma function inequalities.

Math. Comp., 47(176):659–667, 1986.



Mehmet Çelik, Luke Duane-Tessier, Ashley M. Rodriguez, Daniel Rodriguez, and Aden Shaw.

Area differences under analytic maps and operators.

Czechoslovak Math. J., 2024.



John P. D'Angelo.

Hermitian analysis.

Cornerstones. Birkhäuser/Springer, Cham, second edition, 2019.

From Fourier series to Cauchy-Riemann geometry.

References IV



Miroslav Engliš.

Functions invariant under the Berezin transform.

J. Funct. Anal., 121(1):233–254, 1994.



Robert E. Greene and Steven G. Krantz.

Function theory of one complex variable, volume 40 of *Graduate Studies in Mathematics*.

American Mathematical Society, Providence, RI, second edition, 2002.



Nihat Gökhan Göğüş and Sönmez Şahutoğlu.

On convergence of the Berezin transforms.

J. Math. Anal. Appl., 491(1):124295, 16, 2020.



Friedrich Haslinger.

The canonical solution operator to $\bar{\partial}$ restricted to spaces of entire functions.

Ann. Fac. Sci. Toulouse Math. (6), 11(1):57–70, 2002.



Friedrich Haslinger.

Complex analysis.

De Gruyter Graduate. De Gruyter, Berlin, 2018.

A functional analytic approach.



F. Holland and J. B. Twomey.

On hardy classes and the area function.

J. London Math. Soc. (2), 17(2):275–283, 1978.

References VI



Alexander Herbert and John Wermer.

Several complex variables and Banach algebras, volume 35 of *Graduate Texts in Mathematics*.

3 edition, 1998.



R. R. London and P. C. McCarthy.

On the area of the image of a bounded analytic function.

Bull. London Math. Soc., 18(3):277–283, 1986.



Paul-André Meyer.

Quantum probability for probabilists, volume 1538 of *Lecture Notes in Mathematics*.

Springer-Verlag, Berlin, 1993.

References VII



Nikolai Nikolski.

Toeplitz matrices and operators, volume 182 of *Cambridge Studies in Advanced Mathematics*.

Cambridge University Press, Cambridge, french edition, 2020.



Sivaguru Ravisankar and Yunus E. Zeytuncu.

A note on smoothing properties of the Bergman projection.

Internat. J. Math., 27(11):1650087, 10, 2016.



E.M. Stein and R. Shakarchi.

Complex Analysis.

Princeton lectures in analysis. Princeton University Press, 2010.



J. Michael Steele.

The Cauchy-Schwarz Master Class: An Introduction to the Art of Mathematical Inequalities.

Cambridge University Press, 2004.

References VIII



F. G. Tricomi and A. Erdélyi.

The asymptotic expansion of a ratio of gamma functions.

Pacific J. Math., 1:133–142, 1951.



K. J. Wirths and J. Xiao.

An image-area inequality for some planar holomorphic maps.

Results in Mathematics. Resultate der Mathematik, 38(1-2):172–179, 2000.



Yunus E. Zeytuncu.

Sobolev regularity of weighted Bergman projections on the unit disc.

Complex Var. Elliptic Equ., 58(3):309–315, 2013.



Kehe Zhu.

Operator theory in function spaces, volume 138 of *Mathematical Surveys and Monographs*.

2 edition, 2007.



Kehe Zhu.

Analysis on Fock spaces, volume 263 of *Graduate Texts in Mathematics*.
Springer, New York, 2012.