

I am using \bar{z} to denote the conjugate of a complex number and T^* to denote the adjoint of an operator.

Chapter 25 Problems

Problem 1

(a)

$$\begin{aligned}
 \| |1\rangle \|^2 &= \langle 1 | 1 \rangle \\
 &= (1) \overline{(1)} + (i) \overline{(i)} \\
 &= 2 \\
 \| |2\rangle \|^2 &= \langle 2 | 2 \rangle \\
 &= (-i) \overline{(-i)} + (2i) \overline{(2i)} \\
 &= 5 \\
 \| |3\rangle \|^2 &= \langle 3 | 3 \rangle \\
 &= \left(e^{i\phi} \right) \overline{(e^{i\phi})} + (-1) \overline{(-1)} \\
 &= 2 \\
 \| |4\rangle \|^2 &= \langle 4 | 4 \rangle \\
 &= (1) \overline{(1)} + (-2i) \overline{(-2i)} + (1) \overline{(1)} \\
 &= 6 \\
 \| |5\rangle \|^2 &= \langle 5 | 5 \rangle \\
 &= (i) \overline{(i)} + (1) \overline{(1)} + (i) \overline{(i)} \\
 &= 3.
 \end{aligned}$$

(b)

$$\begin{aligned}
 \langle 2 | 1 \rangle &= (1) \overline{(-i)} + (i) \overline{(2i)} \\
 &= \overline{-i(1) + 2i(i)} \\
 &= 2 + i \\
 &= \overline{\langle 1 | 2 \rangle} \\
 \langle 3 | 1 \rangle &= (1) \overline{(e^{i\phi})} + (i) \overline{(-1)} \\
 &= \overline{e^{i\phi}(1) + (-1)(i)} \\
 &= e^{-i\phi} - i \\
 &= \overline{\langle 1 | 3 \rangle} \\
 \langle 3 | 2 \rangle &= (-i) \overline{(e^{i\phi})} + (2i) \overline{(-1)} \\
 &= \overline{(e^{i\phi})(-i) + (-1)(2i)} \\
 &= -ie^{-i\phi} - 2i \\
 &= \overline{\langle 2 | 3 \rangle}. \\
 \langle 5 | 4 \rangle &= (1) \overline{(i)} + (-2i) \overline{(1)} + (1) \overline{(i)} \\
 &= \overline{i(1) + (1)(-2i) + (i)(1)} \\
 &= -4i
 \end{aligned}$$

$$= \overline{\langle 4 | 5 \rangle}$$

Problem 4

(a)

$$\begin{aligned} |u\rangle^* &= (M |v\rangle)^* \\ &= \left(\begin{pmatrix} 1 & i \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -i \end{pmatrix} \right)^* \\ &= \left(\begin{pmatrix} 2 \\ 2-i \end{pmatrix} \right)^* \\ &= \begin{pmatrix} 2 & 2+i \end{pmatrix} \\ &= \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ -i & 1 \end{pmatrix} \\ &= \langle u|. \end{aligned}$$

(b)

$$\begin{aligned} \langle w | v \rangle &= \langle w | Mv \rangle \\ &= \langle w | u \rangle \\ &= \begin{pmatrix} -1 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 2-i \end{pmatrix} \\ &= -i \\ &= \overline{\langle u | w \rangle} \\ &= \overline{\begin{pmatrix} 2 & 2+i \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix}} \\ &= \overline{(i)} \\ &= -i. \end{aligned}$$

Problem 5

$$\begin{aligned} \langle v | Lw \rangle &= \langle v | L | w \rangle \\ &= \langle L^* v | w \rangle \\ &= \overline{\langle w | L^* v \rangle} \\ &= \overline{\langle w | L^* | v \rangle}. \end{aligned}$$

Problem 6

(a)

$$\begin{aligned} \overline{\overline{\langle v | T | w \rangle}} &= \overline{\langle w | T^* | v \rangle} \\ &= \langle v | T^{**} | w \rangle. \end{aligned}$$

(b)

$$\begin{aligned} \langle v | (ST)^* | w \rangle &= \overline{\langle w | S (T | v \rangle)} \\ &= \overline{\langle w | S | u \rangle} \end{aligned}$$

$$|u\rangle = T |v\rangle$$

$$\begin{aligned}
&= \langle u | S^* | w \rangle \\
&= \langle Tv | S^* | w \rangle \\
&= \langle v | T^* S^* | w \rangle.
\end{aligned}$$

Alternatively,

$$\begin{aligned}
\langle v | (ST)^* | w \rangle &= \langle (ST) v | w \rangle \\
&= \langle Tv | S^* | w \rangle \\
&= \langle v | T^* S^* | w \rangle.
\end{aligned}$$

Problem 8

(a) It is clear that

$$\begin{aligned}
||1\rangle &= 1 \\
||2\rangle &= 1,
\end{aligned}$$

and

$$\begin{aligned}
\langle 1 | 2 \rangle &= \overline{(i \ 0)} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
&= 0.
\end{aligned}$$

(b) It is clear that

$$\begin{aligned}
||+\rangle &= 1 \\
||-\rangle &= 1,
\end{aligned}$$

and

$$\begin{aligned}
\langle + | - \rangle &= \frac{1}{2} \overline{(1 \ -i)} \begin{pmatrix} 1 \\ -i \end{pmatrix} \\
&= 0.
\end{aligned}$$

(c) Similarly, it is clear that

$$\begin{aligned}
||\uparrow\rangle &= 1 \\
||\downarrow\rangle &= 1.
\end{aligned}$$

Taking inner products, we have

$$\begin{aligned}
\langle \uparrow | \downarrow \rangle &= \frac{1}{2} \overline{(1 \ 1)} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\
&= 0.
\end{aligned}$$

(d)

$$\begin{aligned}
||I\rangle &= \frac{1}{9} \left(|i - \sqrt{3}|^2 + |1 + 2i|^2 \right) \\
&= \frac{1}{9} (9) \\
&= 1
\end{aligned}$$

$$\begin{aligned}
 \| |II\rangle \|^2 &= \frac{1}{9} \left(|1 - 2i|^2 + |i + \sqrt{3}|^2 \right) \\
 &= \frac{1}{9} (9) \\
 &= 1
 \end{aligned}$$

$$\begin{aligned}
 \langle I | II \rangle &= \frac{1}{9} \overline{(i - \sqrt{3} \quad 1 + 2i)} \begin{pmatrix} 1 - 2i \\ i + \sqrt{3} \end{pmatrix} \\
 &= 0.
 \end{aligned}$$

Problem 9

(a)

$$\begin{aligned}
 a_1 &= \langle 1 | A \rangle \\
 &= 1 - 2i \\
 a_2 &= \langle 2 | A \rangle \\
 &= 2 + 2i \\
 \| |A\rangle \|^2 &= |a_1|^2 + |a_2|^2 \\
 &= 10
 \end{aligned}$$

$$\begin{aligned}
 b_1 &= \langle 1 | B \rangle \\
 &= 1 + i \\
 b_2 &= \langle 2 | B \rangle \\
 &= 2i \\
 \| |B\rangle \|^2 &= |b_1|^2 + |b_2|^2 \\
 &= 6
 \end{aligned}$$

(b)

$$\begin{aligned}
 a_+ &= \langle + | A \rangle \\
 &= \frac{1}{\sqrt{2}} \overline{(1 \quad i)} \begin{pmatrix} 1 - i \\ 2 + 2i \end{pmatrix} \\
 &= \frac{1}{\sqrt{2}} (3 - 3i) \\
 a_- &= \langle - | A \rangle \\
 &= \frac{1}{\sqrt{2}} \overline{(1 \quad -i)} \begin{pmatrix} 1 - i \\ 2 + 2i \end{pmatrix} \\
 &= \frac{1}{\sqrt{2}} (-1 + i) \\
 \| |A\rangle \|^2 &= |a_+|^2 + |a_-|^2 \\
 &= 10
 \end{aligned}$$

$$\begin{aligned}
 b_+ &= \langle + | B \rangle \\
 &= \frac{1}{\sqrt{2}} \overline{(1 \quad i)} \begin{pmatrix} 1 + i \\ 2i \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\sqrt{2}} (3 + i) \\
b_- &= \langle - | B \rangle \\
&= \frac{1}{\sqrt{2}} (1 - i) \begin{pmatrix} 1 + i \\ 2i \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} (2i) \\
\| |B\rangle \|^2 &= |b_+|^2 + |b_-|^2 \\
&= 6
\end{aligned}$$

(c)

$$\begin{aligned}
a_\uparrow &= \langle \uparrow | A \rangle \\
&= \frac{1}{\sqrt{2}} (1 - 1) \begin{pmatrix} 1 - i \\ 2 + 2i \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} (3 + i) \\
a_\downarrow &= \langle \downarrow | A \rangle \\
&= \frac{1}{\sqrt{2}} (1 - 1) \begin{pmatrix} 1 - i \\ 2 + 2i \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} (-1 - 3i) \\
\| |A\rangle \|^2 &= |a_\uparrow|^2 + |a_\downarrow|^2 \\
&= 10
\end{aligned}$$

$$\begin{aligned}
b_\uparrow &= \langle \uparrow | B \rangle \\
&= \frac{1}{\sqrt{2}} (1 - 1) \begin{pmatrix} 1 + i \\ 2i \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} (1 + 3i) \\
b_\downarrow &= \langle \downarrow | B \rangle \\
&= \frac{1}{\sqrt{2}} (1 - 1) \begin{pmatrix} 1 + i \\ 2i \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} (1 - i) \\
\| |B\rangle \|^2 &= |b_\uparrow|^2 + |b_\downarrow|^2 \\
&= 6.
\end{aligned}$$

(d)

$$\begin{aligned}
a_I &= \langle I | A \rangle \\
&= \frac{1}{3} (\overline{i - \sqrt{3} - 1 + 2i}) \begin{pmatrix} 1 - i \\ 2 + 2i \end{pmatrix} \\
&= \frac{5\sqrt{3} - 1}{\sqrt{3}} - \frac{(1 + \sqrt{3})i}{\sqrt{3}}.
\end{aligned}$$

$$\begin{aligned}
a_{II} &= \langle II | A \rangle \\
&= \frac{1}{3} \overline{\begin{pmatrix} 1-2i & i+\sqrt{3} \end{pmatrix}} \begin{pmatrix} 1-i \\ 2+2i \end{pmatrix} \\
&= \frac{5\sqrt{3}+2}{\sqrt{3}} + \frac{(2-\sqrt{3})i}{\sqrt{3}} \\
\| |A\rangle \|^2 &= |a_I|^2 + |a_{II}|^2 \\
&= 10
\end{aligned}$$

$$\begin{aligned}
b_I &= \langle I | B \rangle \\
&= \frac{1}{3} \overline{\begin{pmatrix} i-\sqrt{3} & 1+2i \end{pmatrix}} \begin{pmatrix} 1+i \\ 2i \end{pmatrix} \\
&= \left(\frac{5-\sqrt{3}}{3} \right) + \left(\frac{1-\sqrt{3}}{3} \right) i \\
b_{II} &= \langle II | B \rangle \\
&= \frac{1}{3} \overline{\begin{pmatrix} 1-2i & i+\sqrt{3} \end{pmatrix}} \begin{pmatrix} 1+i \\ 2i \end{pmatrix} \\
&= \frac{1}{3} + \left(1 + \frac{2}{\sqrt{3}} \right) i \\
\| |B\rangle \|^2 &= |b_I|^2 + |b_{II}|^2 \\
&= 6.
\end{aligned}$$

Problem 13

$$\begin{aligned}
|\hat{A}\rangle &= \frac{|A\rangle}{\sqrt{\langle A | A \rangle}} \\
&= \frac{1}{\sqrt{3}} (|\hat{e}_1\rangle + |\hat{e}_2\rangle + |\hat{e}_3\rangle).
\end{aligned}$$

Problem 17

$$\begin{aligned}
|e_3\rangle &= |M_3\rangle - \langle \hat{e}_1 | M_3 \rangle |\hat{e}_1\rangle - \langle \hat{e}_2 | M_3 \rangle |\hat{e}_2\rangle \\
&= \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} - \frac{1}{2} \text{tr}(\hat{e}_1^* M_3) \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} - \frac{1}{2} \text{tr}(\hat{e}_2^* M_3) \left(\frac{i}{\sqrt{2}} \right) \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} - \frac{1+i}{4} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} + \frac{1+i}{4} \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} \\
&= \begin{pmatrix} \frac{1-i}{2} & 0 \\ 0 & -\frac{1-i}{2} \end{pmatrix} \\
|\hat{e}_3\rangle &= \frac{|e_3\rangle}{\| |e_3\rangle \|} \\
&= \frac{1-i}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}
\end{aligned}$$

To find $|e_4\rangle$, we can see that $\| |M_4\rangle \| = 1$ and $\langle \hat{e}_1 | M_4 \rangle = \langle \hat{e}_2 | M_4 \rangle = \langle \hat{e}_3 | M_4 \rangle = 0$. Thus, $|\hat{e}_4\rangle = |M_4\rangle$.

Problem 18

(a) Note that

$$\langle \phi_m | \phi_m \rangle = k_m$$

so

$$\begin{aligned} 1 &= \frac{1}{k_m} \langle \phi_m | \phi_m \rangle \\ &= \left\langle \frac{1}{\sqrt{k_m}} \phi_m \left| \frac{1}{\sqrt{k_m}} \phi_m \right. \right\rangle \\ &= \langle \hat{\phi}_m | \hat{\phi}_m \rangle. \end{aligned}$$

Thus,

$$\begin{aligned} |v\rangle &= \sum_m \langle \hat{\phi}_m | v \rangle |\hat{\phi}_m\rangle \\ &= \sum_m \left\langle \frac{1}{\sqrt{k_m}} \phi_m \left| v \right. \right\rangle \left| \frac{1}{\sqrt{k_m}} \phi_m \right\rangle \\ &= \sum_m \frac{1}{k_m} \langle \phi_m | v \rangle |\phi_m\rangle \\ &= \sum_m c_m |\phi_m\rangle. \end{aligned}$$

$$\text{Thus, } c_m = \frac{1}{k_m} \langle \phi_m | v \rangle.$$

(b)

$$\begin{aligned} \text{id}_V &= \sum_m |\hat{\phi}_m\rangle \langle \hat{\phi}_m| \\ &= \sum_m \left| \frac{1}{\sqrt{k_m}} \phi_m \right\rangle \left\langle \frac{1}{\sqrt{k_m}} \phi_m \right| \\ &= \sum_m \frac{1}{k_m} |\phi_m\rangle \langle \phi_m|. \end{aligned}$$

Problem 19

$$\begin{aligned} (M_{ij})^* &= (\langle \hat{e}_i | \mathcal{M} | \hat{e}_j \rangle)^* \\ &= \langle \hat{e}_j | \mathcal{M}^* | \hat{e}_i \rangle \\ &= \overline{M_{ji}}. \end{aligned}$$

Problem 26

We can see that $|\phi_1\rangle$ and $|\phi_2\rangle$ are linearly independent, and similarly are $|\phi_3\rangle$ and $|\phi_4\rangle$. Since $|\phi_2\rangle$ and $|\phi_3\rangle$ are necessarily linearly independent, the collection of $|\phi_i\rangle$ are linearly independent.

Therefore,

$$|\hat{e}_1\rangle = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

$$\begin{aligned}
 |e_2\rangle &= \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix} - \frac{1}{4} \left[\frac{1}{(1 \ 1 \ 1 \ 1)} \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix} \right] \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix} \\
 |\hat{e}_2\rangle &= \frac{1}{2} \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 |e_3\rangle &= |\phi_3\rangle - \langle \hat{e}_1 | \phi_3 \rangle - \langle \hat{e}_2 | \phi_3 \rangle \\
 &= \frac{i}{2} \begin{pmatrix} 0 \\ -1 \\ 0 \\ 1 \end{pmatrix} \\
 |\hat{e}_3\rangle &= \frac{i}{\sqrt{2}} \begin{pmatrix} 0 \\ -1 \\ 0 \\ 1 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 |e_4\rangle &= |\phi_4\rangle - \langle \hat{e}_1 | \phi_4 \rangle |\hat{e}_1\rangle - \langle \hat{e}_2 | \phi_4 \rangle |\hat{e}_2\rangle - \langle \hat{e}_3 | \phi_4 \rangle |\hat{e}_3\rangle \\
 &= \frac{1-i}{2} \begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \end{pmatrix} \\
 |\hat{e}_4\rangle &= \frac{1-i}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \end{pmatrix}
 \end{aligned}$$

Problem 29

(a)

$$\mathcal{T} = |2\rangle \langle 1| + |1\rangle \langle 2|$$

(b)

$$\begin{aligned}
 \mathcal{T} &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\
 &= \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \\
 &= \begin{pmatrix} 0 & \frac{1+i}{\sqrt{2}} \\ \frac{1+i}{\sqrt{2}} & 0 \end{pmatrix}
 \end{aligned}$$

Problem 30

$$\begin{aligned}\mathcal{S} &= |\hat{e}_1\rangle \langle \hat{e}_1| + |-\hat{e}_3\rangle \langle \hat{e}_3| \\ \mathcal{T} &= |\hat{e}_3\rangle \langle \hat{e}_1| + |\hat{e}_2\rangle \langle \hat{e}_2| + |\hat{e}_1\rangle \langle \hat{e}_3|.\end{aligned}$$

In the standard basis, we have

$$\begin{aligned}\mathcal{S} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \\ \mathcal{T} &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.\end{aligned}$$