

Problem 1

Find $\sup(A)$ and $\inf(A)$ where

(a) $A := \left\{ 1 - \frac{(-1)^n}{n} \mid n \in \mathbb{N} \right\}$

(b) $A := \left\{ \frac{1}{n} - \frac{1}{m} \mid m, n \in \mathbb{N} \right\}$

(c) $A := \left\{ \frac{m}{n} \mid m, n \in \mathbb{N}, m + n \leq 10 \right\}$

(a)

$\sup(A) = 2$: For any $t \in A$, $t < 2$, we can find a_t as follows:

$$a_t := \begin{cases} 1, & t < 1 \\ \frac{4}{3}, & 1 \leq t < \frac{4}{3} \\ 2, & t = \frac{4}{3} \end{cases}$$

$\inf(A) = \frac{1}{2}$: For any $t \in A$, $t > \frac{1}{2}$, we can find a_t as follows:

$$a_t := \begin{cases} 1, & t > 1 \\ \frac{3}{4}, & \frac{3}{4} < t \leq 1 \\ \frac{1}{2}, & t < \frac{3}{4} \end{cases}$$

(b)

$\sup(A) = 1$: For any $t \in A$, $t < 1$, we can find $a_t > t$ as follows:

(1) Take $|t| \geq t$.

(2) If $|t| < \frac{1}{2}$, find m such that $\frac{1}{m} < |t|$ (which exists by the Archimedean Property corollary). Set $a_t = 1 - \frac{1}{m}$.

(3) If $|t| > \frac{1}{2}$, then find m such that $\frac{1}{m} < 1 - |t|$, and set $a_t = 1 - \frac{1}{m}$.

In all three cases, $a_t > t$, meaning $\sup(A) = 1$

$\inf(A) = -1$

(c)

Since A is finite, $\sup(A) = \max(A) = 9$ and $\inf(A) = \min(A) = \frac{1}{9}$

Problem 2

Suppose $u = \sup(A)$ such that $u \notin A$. Show that there is a strictly increasing sequence

$$t_1 < t_2 < t_3 < \dots$$

With $t_n \in A$ and $t_n + \frac{1}{n} > u$ for all $n \geq 1$

Let $t_n = u - \frac{1}{2n}$. t_n must be a strictly increasing sequence because $t_{n+1} = u - \frac{1}{2n+2} > u - \frac{1}{2n} = t_n$.

Additionally, $t_n + \frac{1}{n} = u - \frac{1}{n} < u$, meaning $t_n \in A$.

Problem 3

If m is a lower bound for $A \subseteq \mathbb{R}$, show that the following are equivalent:

- (i) $m = \inf(A)$
- (ii) $\forall t > m, \exists a_t \in A \ni a_t < t$
- (iii) $\forall \varepsilon > 0, \exists a_\varepsilon \ni m + \varepsilon > a_\varepsilon$

Problem 4

Let $A, B \subseteq \mathbb{R}$ be bounded subsets.

- (a) Show that

$$\begin{aligned}\sup(A + B) &= \sup(A) + \sup(B) \\ \inf(A + B) &= \inf(A) + \inf(B)\end{aligned}$$

- (b) If $t > 0$, show that

$$\begin{aligned}\sup(tA) &= t \sup(A) \\ \inf(tA) &= t \inf(A)\end{aligned}$$

(a)

Let $a = \sup(A)$ and $b = \sup(B)$, and $x_a \in A$ and $x_b \in B$. Then

$$\begin{aligned}a &\geq x_a \\ a + x_b &\geq x_a + x_b && \text{by the ordering of } \mathbb{R} \\ a + b &\geq a + x_b && \text{by the definition of } \sup(B) \\ a + b &\geq x_a + x_b && \text{by the ordering of } \mathbb{R} \\ \sup(A) + \sup(B) &= \sup(A + B)\end{aligned}$$

Let $a' = \inf(A)$ and $b' = \inf(B)$, with x_a and x_b defined as above. Then

$$\begin{aligned}a' &\leq x_a \\ a' + x_b &\leq x_a + x_b && \text{by the ordering of } \mathbb{R} \\ a' + b' &\leq a' + x_b && \text{by the definition of } \inf(B) \\ a' + b' &\leq x_a + x_b && \text{by the ordering of } \mathbb{R} \\ \inf(A) + \inf(B) &= \inf(A + B)\end{aligned}$$

(b)

Let $a = \sup(A)$, $x_a \in A$, and $t > 0$. Then

$$\begin{aligned}a &\geq x_a \\ ta &\geq tx_a && \text{by the ordering of } \mathbb{R} \\ t \sup(A) &= \sup(tA)\end{aligned}$$

Let $a' = \inf(A)$, with x_a and t defined as above.

$$\begin{aligned}a' &\leq x_a \\ ta' &\leq tx_a && \text{by the ordering of } \mathbb{R} \\ t \inf(A) &= \inf(tA)\end{aligned}$$

Problem 5

Let $I = (0, 1)$ denote the open unit interval and consider $F : I \times I \rightarrow \mathbb{R}$, $F(x, y) = 2x + y$.

Compute

$$\sup_{y \in I} \left(\inf_{x \in I} F(x, y) \right)$$

and

$$\inf_{x \in I} \left(\sup_{y \in I} F(x, y) \right)$$

We start by finding $\inf_{x \in I} F(x, y)$, which is equal to $F(x, y) = y$ (as the infimum is the greatest lower bound on $2x$, which is $2(0) = 0$). So, $\sup_{y \in I} y = 1$.

We start by finding $\sup_{y \in I} F(x, y)$, which is $\sup_{y \in I} 2x + y$, which is $2x + 1$, as $\sup I = 1$. So, by similar reasoning, $\inf_{x \in I} 2x + 1 = 1$.

These values are the same.

Problem 6

Let D be a nonempty set and consider the vector space

$$\ell_\infty(D) := \{f \mid f : D \rightarrow \mathbb{R} \text{ is bounded}\}$$

with point-wise addition and scalar multiplication. Show that

$$\|f\|_u := \sup_{x \in D} |f(x)|$$

defines a norm on $\ell_\infty(D)$.

- (1) Because $\forall x \in \mathbb{R}, |x| \geq 0$, $\|\cdot\|_u \geq 0$.
- (2) $\|f + g\|_u = \sup_{x \in D} |f(x) + g(x)| \leq \sup_{x \in D} |f(x)| + \sup_{x \in D} |g(x)|$ (by the Triangle Inequality) $= \|f\|_u + \|g\|_u$.
- (3) $\|\mathbf{0}\| = \sup_{x \in D} |\mathbf{0}| = 0$.
- (4) Let $\|f\|_u = 0$. Then, $\sup_{x \in D} |f(x)| = 0$, meaning that $\nexists x' \in D$ such that $f(x') \neq 0$ (or else $\sup_{x \in D} |f(x)| = f(x')$), so $f(x) = \mathbf{0}$.
- (5) $\|tf\|_u = \sup_{x \in D} |tf(x)| = |t| \sup_{x \in D} |f(x)| = |t| \|f\|_u$.

Therefore, $\|\cdot\|_u$ is a norm on ℓ_∞ .

Problem 7

Let $f, g : D \rightarrow \mathbb{R}$ be bounded functions. Show that

- (a) $\sup_{x \in D} (f + g)(x) \leq \sup_{x \in D} f(x) + \sup_{x \in D} g(x)$
- (b) $\inf_{x \in D} (f + g)(x) \geq \inf_{x \in D} f(x) + \inf_{x \in D} g(x)$
- (c) $|\sup_{x \in D} f(x) - \sup_{x \in D} g(x)| \leq \sup_{x \in D} |f(x) - g(x)|$

Problem 8

Find $\bigcap_{n=1}^{\infty} I_n$ where

- (a) $I_n = [0, 1/n]$
- (b) $I_n = (0, 1/n)$
- (c) $I_n = [n, \infty)$

(a)

For all $k > 1$, $\bigcap_{n=1}^k = [0, 1/k]$, meaning that $\bigcap_{n=1}^{\infty} = \lim_{k \rightarrow \infty} [0, 1/k] = \{0\}$.