

# Contents

<b>Introduction</b>	<b>1</b>
<b>Algebras and <math>\ast</math>-Algebras</b>	<b>1</b>
Definitions and Examples . . . . .	1
Algebraic Constructions . . . . .	4
Subalgebras, Ideals, Products, Sums, and Tensor Products . . . . .	4
The Group $\ast$ -Algebra . . . . .	7
Distinguished Elements . . . . .	8
Algebra Homomorphisms . . . . .	9
Unitization . . . . .	10
Algebraic Spectrum . . . . .	13
<b>Banach and <math>C^\ast</math>-Algebras</b>	<b>15</b>
Examples . . . . .	15
Constructions . . . . .	17
Generating Sets . . . . .	17
Products, Sums, and Quotients . . . . .	19
Ideals in $C_0(\Omega)$ . . . . .	20
$C^\ast$ -norms . . . . .	22
Universal $C^\ast$ -Algebras . . . . .	25
Representations and the Group $C^\ast$ -algebra . . . . .	28
Unitizations of $C^\ast$ -Algebras . . . . .	30
Gelfand Theory . . . . .	34
Properties of the Spectrum . . . . .	34
The Character Space . . . . .	41
The Gelfand Transform . . . . .	46
Continuous Functional Calculus . . . . .	53
Ordering and Positive Elements . . . . .	61
Approximate Identities . . . . .	70
Representations, States, and the GNS Construction . . . . .	75

## Introduction

This is going to be part of my notes for my Honors Thesis independent study focused on Amenability and  $C^\ast$ -algebras. This set of notes will be focused on the theory of Banach algebras and  $C^\ast$ -algebras. The primary source for this section of notes will be Timothy Rainone's *Functional Analysis: En Route to Operator Algebras*.

I do not claim any of this work to be original.

## Algebras and $\ast$ -Algebras

A lot of the structures we encounter in functional analysis, like  $\mathbb{B}(X)$ , are not only vector spaces, but also come with an algebraic structure with them. We will learn these more in depth before venturing into the study of Banach and  $C^\ast$ -algebras.

## Definitions and Examples

We will let  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ .

**Definition.** An  $\mathbb{F}$ -algebra is a vector space  $A$  over the field  $\mathbb{F}$  with a multiplication operation  $(a, b) \mapsto a \cdot b$  satisfying the following for all  $a, b, c \in A$  and  $\alpha \in \mathbb{F}$ ;

- $a \cdot (b \cdot c) = (a \cdot b) \cdot c$ ;
- $a \cdot (b + c) = a \cdot b + a \cdot c$ ;
- $\alpha(a \cdot b) = (\alpha a) \cdot b = a \cdot (\alpha b)$ .

An algebra is called unital if there is a unique  $1_A \in A$  such that  $1_A \cdot a = a \cdot 1_A = a$ .

We say the algebra is commutative if multiplication is commutative, else it is called noncommutative.

**Remark:** Usually,  $\mathbb{F} = \mathbb{C}$  unless otherwise specified, and we drop the multiplication sign, writing  $ab$  for  $a \cdot b$ .

**Remark:** If  $A$  is an  $\mathbb{F}$ -vector space with basis  $B$ , we can always extend an associative map  $B \times B \rightarrow B$  to multiplication in  $A$  by defining multiplication by the associative map on the basis elements.

**Example (Functions).** Let  $\Omega$  be any nonempty set. The function space  $\mathcal{F}(\Omega, \mathbb{F})$ , equipped with pointwise addition, scalar multiplication, and pointwise multiplication is an algebra.

In general, if  $A$  is an  $\mathbb{F}$ -algebra, then  $\mathcal{F}(\Omega, A) = \{f \mid f: \Omega \rightarrow A\}$  is an  $\mathbb{F}$ -algebra. If  $A$  is unital, then the constant map  $u(x) = 1_A$  is the unit for  $\mathcal{F}(\Omega, A)$ .

**Example (Linear Maps).** If  $X$  is a vector space over  $\mathbb{F}$ , then  $\mathcal{L}(X)$ , the space of all linear maps from  $X$  to itself, is a unital  $\mathbb{F}$ -algebra with multiplication as composition.

**Example (Polynomials in One Variable).** If  $x$  is an abstract variable, then the linear space of all polynomials,

$$\mathbb{F}[x] = \left\{ \sum_{k=0}^n a_k x^k \mid a_k \in \mathbb{F}, n \in \mathbb{Z}_+ \right\}$$

is an  $\mathbb{F}$ -algebra. We define multiplication via ordinary multiplication of polynomials,

$$\left( \sum_{i=0}^n a_i x^i \right) \left( \sum_{j=0}^m b_j x^j \right) = \sum_{k=0}^{m+n} \left( \sum_{i=0}^k a_i b_{k-i} \right) x^k.$$

If we let  $x^0 = 1$ , this space is a commutative unital algebra.

**Example (General Polynomial).** If we have a set  $S = \{x_i\}_{i \in I}$  of abstract symbols, then  $\mathbb{F}\langle S \rangle$  is the space of all (not necessarily commuting) polynomials with symbols in  $S$ , where multiplication is defined by concatenation. This is a unital algebra.

If the symbols in  $S$  commute, then this is a commutative algebra, and we write  $\mathbb{F}[S]$ .

**Example.** If  $x$  is an abstract symbol, then

$$\mathbb{F}(x) = \left\{ \frac{p(x)}{q(x)} \mid p, q \in \mathbb{F}[x], q \neq 0 \right\}$$

is the unital commutative algebra of all rational functions.

**Definition.** Let  $A$  be an  $\mathbb{F}$ -algebra, and let  $a \in A$  be fixed. For  $p \in \mathbb{F}[x]$ , we define

$$p(a) = \sum_{k \geq 0} \alpha_k a^k.$$

It is assumed that  $\alpha_0 = 0$  when  $A$  does not have a unit.

Generally, if  $a_1, \dots, a_n \in A$  and  $p = \sum_I c_I x^I$  in  $\mathbb{F}\langle x_1, \dots, x_n \rangle$ , then

$$p(a_1, \dots, a_n) = \sum_I c_I a^I,$$

where  $a^I = a_1^{i_1} a_2^{i_2} \dots a_n^{i_n}$ , and  $(i_1, \dots, i_n) = I \in \mathbb{Z}_+^n$  is a multi-index.

**Remark:** The binomial theorem only holds for commutative algebras.

**Definition.** Let  $A$  be an algebra over  $\mathbb{C}$ . An involution on  $A$  is a self-map  $*$ :  $A \rightarrow A$  that satisfies the following, for all  $a, b \in A$  and  $\alpha \in \mathbb{C}$ :

- (1)  $(a + \alpha b)^* = a^* + \overline{\alpha} b^*$ ;
- (2)  $(ab)^* = b^* a^*$ ;
- (3)  $a^{**} = a$ .

If  $A$  admits an involution, then  $A$  is known as a  $*$ -algebra.

**Example.** The complex numbers,  $\mathbb{C}$ , is a unital commutative  $*$ -algebra with the usual operations, where  $z \mapsto \bar{z}$  is the involution.

**Example.** We can define an involution on  $\mathcal{F}(\Omega, \mathbb{C})$  by taking  $f^*(x) = \overline{f(x)}$ .

If  $A$  is a  $*$ -algebra, we may define the involution as  $f^*(x) = f(x)^*$ .

**Example (The Free  $*$ -Algebra).** Let  $E = \{x_i\}_{i \in I}$  be a set of abstract symbols. We may add a set of symbols disjoint from  $E$ , called  $E^* = \{x_i^*\}_{i \in I'}$ , and let  $S = E \cup E^*$ .

We consider  $\mathbb{C}\langle S \rangle$ , which is the set of general polynomials over  $S$ . The involution  $*$ :  $\mathbb{C}\langle S \rangle \rightarrow \mathbb{C}\langle S \rangle$  can be defined by

$$\left( \alpha x_{i_1}^{\epsilon_1} x_{i_2}^{\epsilon_2} \dots x_{i_n}^{\epsilon_n} \right)^* = \overline{\alpha} x_{i_n}^{\delta_n} \dots x_{i_2}^{\delta_2} x_{i_1}^{\delta_1},$$

where  $\epsilon_j \in \{1, *\}$  for each  $j = 1, \dots, n$ , and

$$\delta_j = \begin{cases} * & \epsilon_j = 1 \\ 1 & \epsilon_j = * \end{cases}.$$

The  $*$ -algebra,  $\mathbb{C}\langle E \cup E^* \rangle$  is referred to as the free  $*$ -algebra generated by  $E$ , denoted  $A^*(E)$ .

**Example (Matrix Algebra).** Let  $A$  be an algebra, and let

$$\text{Mat}_n(A) = \left\{ (a_{ij})_{ij} \mid 1 \leq i, j \leq n, a_{ij} \in A \right\}.$$

This is an algebra with element-wise addition and scalar multiplication, as well as traditional matrix multiplication. If  $A$  is unital, then  $\text{diag}(1_A, \dots, 1_A)$  is the unit for  $\text{Mat}_n(A)$ . When  $n \geq 2$ , this algebra is non-commutative. If  $A$  is a  $*$ -algebra, then  $\text{Mat}_n(A)$  is a  $*$ -algebra with the involution  $(a_{ij})_{ij}^* = (a_{ji}^*)_{ij}$ .

**Example.** Let  $(\Omega, \mathcal{M})$  be a measurable space, and let  $L_0(\Omega, \mathcal{M})$  be the space of measurable functions. This is a  $*$ -algebra when equipped with pointwise operations and involution.

If  $\mu$  is a measure, then  $L(\Omega, \mathcal{M})$  of  $\mu$ -equivalence classes is also a  $*$ -algebra when equipped with multiplication of equivalence classes and the involution

$$[f]_\mu^* = \left[ \bar{f} \right]_\mu.$$

## Algebraic Constructions

Algebras, like vector spaces and other algebraic objects, admit various sub-objects and super-objects.

### Subalgebras, Ideals, Products, Sums, and Tensor Products

**Definition.** Let  $A$  be a  $*$ -algebra over  $\mathbb{C}$ ,  $B \subseteq A$ .

- (1) If  $B \subseteq A$  is a linear subspace that is closed under multiplication, then  $B$  is known as a subalgebra. If  $1_A \in B$ , then  $B$  is unital.
- (2) If  $B \subseteq A$  is a subalgebra such that, for  $b \in B$  and  $a \in A$ , then  $ab, ba \in B$ , then we say  $B$  is an ideal.
- (3) If, for all  $x \in B$ ,  $x^* \in B$ , then  $B$  is called self-adjoint or  $*$ -closed.
- (4) If  $B$  is a subalgebra that is  $*$ -closed, then we say  $B$  is a  $*$ -subalgebra.
- (5) If  $B$  is an ideal that is  $*$ -closed, then we say  $B$  is a  $*$ -ideal.

**Example.** If  $\Omega$  is a nonempty set with  $\mathcal{F}(\Omega, \mathbb{C})$  its corresponding  $*$ -algebra, then  $\ell_\infty(\Omega) \subseteq \mathcal{F}(\Omega, \mathbb{C})$  is a unital  $*$ -subalgebra.

If  $(\Omega, \mathcal{M})$  is a measurable space, then  $B_\infty(\Omega, \mathcal{M})^I$  is a  $*$ -subalgebra of  $\ell_\infty(\Omega)$  and of  $L_0(\Omega, \mathcal{M})$ .

If  $\mu$  is a measure on  $(\Omega, \mathcal{M})$ , then  $L_\infty(\Omega, \mu)$  of  $\mu$ -essentially bounded functions is a unital  $*$ -subalgebra of  $L(\Omega, \mu)$ . Moreover,  $B_\infty(\Omega, \mu) \subseteq L_\infty(\Omega, \mu)$  is a unital  $*$ -subalgebra.

If  $\Omega$  is a LCH<sup>II</sup> space, then the string of  $*$ -subalgebras is

$$C_c(\Omega) \subseteq C_0(\Omega) \subseteq C_b(\Omega) \subseteq \ell_\infty(\Omega) \subseteq \mathcal{F}(\Omega, \mathbb{C}).$$

It is also the case that  $C_c(\Omega) \subseteq C_0(\Omega) \subseteq C_b(\Omega)$  is a string of  $*$ -ideals.

It is not the case that  $C_b(\Omega) \subseteq B_\infty(\Omega)$  is a  $*$ -ideal.

If  $\mu$  is a Radon measure, then the string of  $*$ -subalgebras is

$$C_c(\Omega, \mu) \subseteq C_0(\Omega, \mu) \subseteq C_b(\Omega, \mu) \subseteq B_\infty(\Omega, \mu) \subseteq L_\infty(\Omega, \mu).$$

**Example.** If  $\Omega \subseteq \mathbb{C}$  is a compact subset of the complex plane, then the set  $P(\Omega)$  of polynomials forms a unital subalgebra of  $C(\Omega)$ , but not a  $*$ -subalgebra. However,

$$\mathcal{Q}(\Omega) = \left\{ q: \Omega \rightarrow \mathbb{C} \mid q(z) = \sum_{k,l=0}^m c_{k,l} z^k \bar{z}^l, c_{k,l} \in \mathbb{C} \right\},$$

the space of Laurent polynomials, is a unital  $*$ -subalgebra of  $C(\Omega)$ .

If  $\Omega = \mathbb{T}$ , then  $\mathcal{Q}(\Omega)$  becomes the unital  $*$ -subalgebra of trigonometric polynomials,

$$\mathcal{T} = \left\{ \sum_{k=-n}^n c_k z^k \mid n \in \mathbb{N}, c_k \in \mathbb{C} \right\}.$$

<sup>I</sup>The space of bounded measurable functions.

<sup>II</sup>locally compact Hausdorff

**Definition.** Let  $A$  be an algebra, and let  $S \subseteq A$  be a subset. The subalgebra (or ideal) generated by  $S$ , denoted  $\text{alg}(S)$  or  $\text{ideal}(S)$ , is the smallest subalgebra or ideal that contains  $S$ :

$$\begin{aligned}\text{alg}(S) &= \bigcap \{B \mid B \supseteq S, B \subseteq A \text{ is a subalgebra}\} \\ \text{ideal}(S) &= \bigcap \{B \mid B \supseteq S, B \subseteq A \text{ is an ideal}\}.\end{aligned}$$

Similarly, we may define  $*\text{-alg}(S)$  and  $*\text{-ideal}(S)$ .

**Example.** Let  $X$  be a vector space, and let  $\mathcal{L}(X)$  be the unital algebra of linear operators on  $X$ . The collection  $\mathbb{F}(X) \subseteq \mathcal{L}(X)$  of finite-rank operators forms an ideal. If  $\dim(V) = \infty$ , then this ideal is proper.

If  $X$  is infinite-dimensional, then  $\mathbb{F}(X)$  is a non-commutative, non-unital subalgebra.

**Definition.** An algebra  $A$  is called simple if it has no nontrivial ideals.

**Example.** The algebra  $\text{Mat}_n(\mathbb{C})$  is simple.

To see this, if  $I \subseteq \text{Mat}_n(\mathbb{C})$  is a nontrivial ideal, and  $0 \neq a \in I$ , we select  $a_{ij} \neq 0$ . For every  $k \in \{1, \dots, n\}$ , we have

$$\begin{aligned}e_{kk} &= \frac{1}{a_{ij}} (a_{ij} e_{kk}) \\ &= \frac{1}{a_{ij}} (e_{ki} a e_{jk}) \\ &\in I,\end{aligned}$$

meaning  $I_n = \sum_k e_{kk}$  is in  $I$ , so  $I = \text{Mat}_n(\mathbb{C})$  is not proper.

**Definition.** If

$$\mathcal{I}_p(A) = \{I \mid I \subsetneq A \text{ is an ideal}\}$$

is the collection of proper ideals ordered by inclusion, we call a maximal element in  $\mathcal{I}_p(A)$  a maximal ideal.

**Theorem:** If  $A$  is a unital algebra, then every proper ideal  $J \subseteq A$  is contained in some maximal ideal  $M$ .

*Proof.* Order  $\mathcal{J} = \{I \mid J \subseteq I \subsetneq A, I \text{ is an ideal}\}$  by inclusion. If  $(I_\lambda)_{\lambda \in \Lambda}$  is a chain in  $\mathcal{J}$ , then  $I = \bigcup_{\lambda \in \Lambda} I_\lambda$  is an ideal in  $A$  containing  $J$ . If  $I = A$ , then  $1_\lambda \in I_\lambda$  for some  $\lambda$ , which contradicts the definition. Thus,  $I$  is proper and belongs to  $\mathcal{J}$ , so by Zorn's lemma, there is some maximal element  $M$  in  $\mathcal{J}$ .  $\square$

We can characterize the maximal ideals of the space  $C(\Omega)$ , where  $\Omega$  is compact. This will be very useful in the future.

**Proposition:** Let  $\Omega$  be a compact Hausdorff space. If  $I \subseteq C(\Omega)$  is a maximal ideal, then there is  $x_0 \in \Omega$  such that

$$\begin{aligned}I &= N_{x_0} \\ &= \{f \in C(\Omega) \mid f(x_0) = 0\}.\end{aligned}$$

Moreover, for every  $x \in \Omega$ ,  $N_x$  is a maximal ideal.

*Proof.* Suppose  $I \neq N_x$  for every  $x \in \Omega$ . Since  $N_x$  is a proper ideal, and  $I$  is maximal, this implies that there is some  $f_x \in I \setminus N_x$ , meaning  $f_x(x) \neq 0$ .

Let  $U_x = f_x^{-1}(\mathbb{C} \setminus \{0\})$ . We must have  $x \in U_x$  for all  $x \in \Omega$ , so

$$\Omega = \bigcup_{x \in \Omega} U_x.$$

Now,  $\Omega$  is compact, so we select  $\{x_1, \dots, x_j\} \subseteq \Omega$  such that

$$\Omega = \bigcup_{j=1}^n U_{x_j}.$$

Define

$$f = \sum_{j=1}^n |f_{x_j}|^2.$$

We have  $f \in I$ , and  $f > 0$  on  $\Omega$  by construction, so  $f$  is invertible in  $C(\Omega)$ . This implies that  $\frac{1}{f} \in C(\Omega)$ , so  $\mathbb{1}_\Omega = f \frac{1}{f} \in I$ , which means  $I = C(\Omega)$ , a contradiction.

Now, we fix  $x \in \Omega$ . If it is the case that  $N_x$  is not maximal, then there is some maximal ideal  $I$  such that  $N_x \subseteq I$ . We know that  $I = N_y$  for some  $y \in \Omega$ , so  $N_x \subseteq N_y$ . This means any continuous function that vanishes at  $x$  must vanish at  $y$ . However, by Urysohn's lemma, this is only possible if  $x = y$ , so  $N_x = I = N_y$ , so  $N_x$  is maximal.  $\square$

**Definition.** Let  $A$  be an algebra,  $J \subseteq A$  is an ideal. Then,  $A/J$  admits multiplication

$$(a + J) \cdot (b + J) = ab + J$$

that makes  $A/J$  into an algebra. If  $1_A \in A$ , then  $A/J$  has unit  $1_A + J$ , and if  $A$  is commutative, so too is  $A/J$ .

If  $A$  is a  $*$ -algebra, and  $J$  is a  $*$ -ideal, then  $A/J$  is a  $*$ -algebra with involution

$$(a + J)^* = a^* + J.$$

**Definition.** If  $\{A_i\}_{i \in I}$  is a family of  $*$ -algebras, the product and coproduct are respectively defined by

$$\prod_{i \in I} A_i = \left\{ f: I \rightarrow \bigcup_{i \in I} A_i \mid f(i) \in A_i \right\}$$

$$\bigoplus_{i \in I} A_i = \left\{ f \in \prod_{i \in I} A_i \mid \text{card}(\text{supp}(f)) < \infty \right\}.$$

Note that  $\bigoplus_{i \in I} A_i \subseteq \prod_{i \in I} A_i$  is a  $*$ -ideal.

**Example (The Universal  $*$ -Algebra).** Let  $E = \{x_i\}_{i \in I}$  be a collection of abstract symbols, and let  $A^*(E)$  be the free  $*$ -algebra generated by  $E$ . Given  $R \subseteq A^*(E)$ , let  $I(R)$  be the  $*$ -ideal generated by  $R$ . The quotient  $*$ -algebra

$$A^*(E|R) = A^*(E)/I(R)$$

is called the universal  $*$ -algebra generated by  $E$  with relations  $R$ . We write  $\pi_R(x_i) = z_i$ .

**Proposition:** Let  $A$  and  $B$  be  $*$ -algebras. The linear space  $A \otimes B$  admits a multiplication

$$(a \otimes b)(a' \otimes b') = aa' \otimes bb'$$

and an involution

$$(a \otimes b)^* = a^* \otimes b^*.$$

*Proof.* Fix  $a \in A$  and  $b \in B$ . Consider the linear maps  $L_a: A \rightarrow A$ , given by  $L_a(x) = ax$ , and  $L_b: B \rightarrow B$ , given by  $L_b(y) = by$ .

The maps  $a \mapsto L_a$  and  $b \mapsto L_b$  are both linear, meaning the map

$$A \times B \rightarrow \mathcal{L}(A) \otimes \mathcal{L}(B),$$

given by  $(a, b) \mapsto L_a \otimes L_b$ , is bilinear. Thus, there is a linear map

$$L: A \otimes B \rightarrow \mathcal{L}(A) \otimes \mathcal{L}(B)$$

given by  $a \otimes b \mapsto L_a \otimes L_b$ . There is a linear embedding  $\mathcal{L}(A) \otimes \mathcal{L}(B) \hookrightarrow \mathcal{L}(A \otimes B)$ , so we may identify the tensors in  $\mathcal{L}(A) \otimes \mathcal{L}(B)$  with the linear operators on  $A \otimes B$ .

We define

$$(A \otimes B) \times (A \otimes B) \rightarrow A \otimes B,$$

given by  $(t, s) \mapsto t \cdot s = L(t)(s)$ . This is a well-defined multiplication following from the fact that  $L$  is linear and  $L(t)$  is linear for all  $t \in A \otimes B$ .

For all  $a, a' \in A$  and  $b, b' \in B$ , we have

$$\begin{aligned} (a \otimes b)(a' \otimes b') &= L(a \otimes b)(a' \otimes b') \\ &= L_a \otimes L_b(a' \otimes b') \\ &= L_a(a') \otimes L_b(b') \\ &= aa' \otimes bb'. \end{aligned}$$

We write  $\overline{A \otimes B}$  for the conjugate vector space. The map

$$A \times B \rightarrow \overline{A \otimes B},$$

given by  $(a, b) \mapsto \overline{a' \otimes b'}$  is bilinear. Thus, there is a linear map  $\psi: A \otimes B \rightarrow \overline{A \otimes B}$  given by  $\psi(a \otimes b) = \overline{a' \otimes b'}$ .

The map  $\mu: \overline{A \otimes B} \rightarrow A \otimes B$ , given by  $\mu(\overline{t}) = t$  is conjugate linear. The composition,  $\nu = \mu \circ \psi$ , mapping  $A \otimes B \rightarrow A \otimes B$  is conjugate linear, and sends  $a \otimes b \mapsto a' \otimes b'$ . We define the involution  $t \mapsto t^* = \nu(t)$ . We have

$$\begin{aligned} ((a \otimes b)(c \otimes d))^* &= (ac \otimes bd)^* \\ &= (ac)^* \otimes (bd)^* \\ &= c^* a^* \otimes b^* d^* \\ &= (c^* \otimes d^*)(a^* \otimes b^*) \\ &= (c \otimes d)^*(a \otimes b)^*. \end{aligned}$$

□

## The Group \*-Algebra

Let  $\Gamma$  be a group, and let  $\mathbb{C}[\Gamma]$  be the free vector space on  $\Gamma$ . For each  $f, g \in \mathbb{C}[\Gamma]$ , we define  $f * g$  by convolution:

$$f * g(s) = \sum_{t \in \Gamma} f(t)g(t^{-1}s)$$

$$= \sum_{r \in \Gamma} f(sr^{-1})g(r).$$

This sum is finite since  $f$  and  $g$  have finite support.

This multiplication has the unit  $1_{\mathbb{C}[\Gamma]} = \delta_e$ .

The involution  $f \mapsto f^*$  in  $\mathbb{C}[\Gamma]$  is defined by

$$f^*(t) = \overline{f(t^{-1})}.$$

We can verify that this forms an involution.

$$\begin{aligned} (f \cdot g)^*(s) &= \overline{f \cdot g(s^{-1})} \\ &= \overline{\sum_{t \in \Gamma} f(t)g(t^{-1}s^{-1})} \\ &= \sum_{t \in \Gamma} \overline{f(t)g(t^{-1}s^{-1})} \\ &= \sum_{r \in \Gamma} \overline{f(r^{-1})g(rs^{-1})} \\ &= \sum_{r \in \Gamma} \overline{g((sr^{-1})^{-1})f(r^{-1})} \\ &= \sum_{r \in \Gamma} g^*(sr^{-1})f^*(r) \\ &= g^* \cdot f^*(s). \end{aligned}$$

The  $*$ -algebra  $\mathbb{C}[\Gamma]$  is known as the group  $*$ -algebra.

## Distinguished Elements

**Definition.** Let  $A$  be a  $*$ -algebra.

- (1) An element  $e \in A$  is said to be idempotent if  $e^2 = e$ . We write  $E(A)$  for the set of idempotents in  $A$ .
- (2) If  $A$  is unital, then  $x \in A$  is said to be invertible if there exists a unique  $y \in A$  with  $xy = yx = 1_A$ . We call  $y$  the inverse of  $x$ , and write  $x^{-1}$ . We write  $GL(A)$  to be the set of all invertible elements in  $A$ .
- (3) An element  $x \in A$  is said to be Hermitian or self-adjoint if  $x = x^*$ . We write  $A_{s.a.}$  for the set of self-adjoint elements in  $A$ .
- (4) An element  $a \in A$  is said to be positive if  $a = b^*b$  for some  $b \in A$ . We write  $A_+$  for the set of all positive elements in  $A$ .
- (5) A projection in  $A$  is a self-adjoint idempotent — that is,  $p^2 = p^* = p$ . We write  $\mathcal{P}(A)$  to be the set of projections in  $A$ .
- (6) If  $A$  is unital, an element  $u \in A$  is said to be unitary if  $u^*u = uu^* = 1_A$ . We write  $\mathcal{U}(A)$  to be the set of all unitaries in  $A$ .
- (7) An element  $z \in A$  is called normal if  $z^*z = zz^*$ . We write  $Nor(A)$  for the collection of normal elements in  $A$ .

**Fact.** Let  $A$  be a  $*$ -algebra.

- The following inclusions hold:



- $\mathcal{P}(A) \subseteq A_+ \subseteq A_{\text{s.a.}} \subseteq \text{Nor}(A)$ ;
- $\mathcal{U}(A) \subseteq \text{Nor}(A)$ .
- The linear span of  $A_{\text{s.a.}}$  is  $A$ . If  $x \in A$ , then

$$h = \frac{1}{2}(x + x^*)$$

$$k = \frac{i}{2}(x^* - x)$$

are self-adjoint with  $x = h + ik$ .

- The self-adjoint elements of  $A$  form a real vector space.
- If  $A$  is unital, then  $\text{GL}(A)$  is  $*$ -closed, with  $(x^*)^{-1} = (x^{-1})^*$ .
- If  $A$  is unital, then  $\mathcal{U}(A) \subseteq \text{GL}(A)$  is a subgroup with  $u^{-1} = u^*$  for all  $u \in \mathcal{U}(A)$ .

**Example.** The spectral theorem from linear algebra states that if a matrix  $a \in \text{Mat}_n(\mathbb{C})$  is normal, then there is a unitary matrix  $u$  with  $a = udu^*$ , where  $d = \text{diag}(\lambda_1, \dots, \lambda_n)$  is a diagonal matrix, and  $\lambda_1, \dots, \lambda_n$  is a complete list of eigenvalues.

Self-adjoint elements in  $\text{Mat}_n(\mathbb{C})$  are matrices that are conjugate symmetric.

A square matrix  $a$  is invertible if and only if  $\det(a) \neq 0$ .

**Example.** Let  $\mathcal{F}(\Omega)$  be the set of all  $\mathbb{C}$ -valued functions on  $\Omega$ . Every element in  $\mathcal{F}(\Omega)$  is normal. The following hold:

- $f \in \mathcal{F}(\Omega)_{\text{s.a.}}$  if and only if  $f(\Omega) \subseteq \mathbb{R}$ ;
- $f \in \mathcal{F}(\Omega)_+$  if and only if  $f(\Omega) \subseteq [0, \infty)$ ;
- $u \in \mathcal{U}(\mathcal{F}(\Omega))$  if and only if  $u(\Omega) \subseteq \mathbb{T}$ ;
- $\mathcal{P}(\mathcal{F}(\Omega)) = \{\mathbb{1}_E \mid E \subseteq \Omega\}$ .

## Algebra Homomorphisms

Now, we can learn about morphisms in the category of algebras and  $*$ -algebras.

**Definition.** Let  $A$  and  $B$  be  $\mathbb{F}$ -algebras.

- (1) An algebra homomorphism is a linear map  $\varphi: A \rightarrow B$  that is multiplicative.
- (2) A character on  $A$  is a nonzero homomorphism  $h: A \rightarrow \mathbb{F}$ . We write

$$\Omega(A) = \{h \mid h \text{ is a character on } A\}.$$

- (3) An algebra isomorphism is a bijective algebra homomorphism.
- (4) If  $A$  and  $B$  are  $*$ -algebras,  $\varphi: A \rightarrow B$  is said to be  $*$ -preserving if  $\varphi(a^*) = \varphi(a)^*$ .
- (5) If  $A$  and  $B$  are  $*$ -algebras, then a  $*$ -homomorphism (or  $*$ -isomorphism) is a homomorphism (or isomorphism) that is  $*$ -preserving.
- (6) An automorphism of a  $*$ -algebra is a  $*$ -isomorphism  $\alpha: A \rightarrow A$ . We write

$$\text{Aut}(A) = \{\alpha \mid \alpha: A \rightarrow A \text{ is a } *- \text{automorphism}\}.$$

(7) If  $A$  and  $B$  are  $*$ -algebras, then  $\phi: A \rightarrow B$  is said to be positive if  $\phi(A_+) \subseteq B_+$ .

(8) A positive map between  $*$ -algebras is called faithful if  $\ker(\phi) \cap A_+ = \{0\}$ .

**Theorem** (First Isomorphism Theorem): Let  $A, B$  be  $*$ -algebras, and let  $I \subseteq A$  be a  $*$ -ideal. If  $\varphi: A \rightarrow B$  is a  $*$ -homomorphism with  $I \subseteq \ker(\varphi)$ , then there exists a unique algebra  $*$ -homomorphism  $\phi: A/I \rightarrow B$  such that  $\phi \circ \pi = \varphi$ .

If  $I = \ker(\varphi)$ , then  $\phi$  is injective, and  $\phi: A/\ker(\varphi) \rightarrow \text{Ran}(\varphi)$  is a  $*$ -isomorphism.

If  $A, B$ , and  $\varphi$  are unital, then so is  $\phi$ .

**Example** (Universal Property of the Universal  $*$ -Algebra). Let  $\mathbb{A}^*(E|R)$  be the universal  $*$ -algebra generated by  $E = \{x_i\}_{i \in I}$  and  $R \subseteq \mathbb{A}^*(E)$ . Let  $B$  be a  $*$ -algebra admitting elements  $\{b_i\}_{i \in I}$  indexed by the same set  $I$  that satisfies the relations in  $R$ .

The evaluation  $*$ -homomorphism,  $\mathbb{A}^*(E) \rightarrow B$  defined by  $x_i \mapsto b_i$  sends  $I(R)$  to 0, so there is a unique  $*$ -homomorphism,  $x_i + I(R) \mapsto b_i$ .

**Corollary:** If  $A$  is an algebra, and  $h \in \Omega(A)$  is a character, then  $\ker(h) \subseteq A$  is a maximal ideal, and  $A/\ker(h) \cong \mathbb{C}$  are isomorphic as algebras.

*Proof.* Since  $h \neq 0$ , and  $h: A \rightarrow \mathbb{C}$  is a linear functional, it is the case that  $\text{Ran}(h) = \mathbb{C}$ .<sup>iii</sup> By the first isomorphism theorem, we have  $A/\ker(h) \cong \mathbb{C}$  as algebras.

Since  $\mathbb{C}$  is simple,  $A/\ker(h)$  is simple, so  $\ker(h) \subseteq A$  is a maximal ideal. □

There are some algebras that do not admit characters.

**Example.** Let  $A = \text{Mat}_n(\mathbb{C})$  for  $n \geq 2$ . If  $h: \text{Mat}_n(\mathbb{C}) \rightarrow \mathbb{C}$  is a character, then  $\ker(h) \subseteq \text{Mat}_n(\mathbb{C})$  is a proper ideal. However, since  $\text{Mat}_n(\mathbb{C})$  is simple,  $\ker(h) = 0$ . However, this means

$$\begin{aligned} n^2 &= \text{Mat}_n(\mathbb{C}) \\ &\leq \dim(\mathbb{C}) \\ &= 1, \end{aligned}$$

which is a contradiction. Thus,  $\Omega(\text{Mat}_n(\mathbb{C})) = \emptyset$ .

## Unitization

It is often the case that algebras lack a unit. However, we can create a “unitized” version of an algebra  $A$ ,  $\tilde{A}$ , such that  $A \subseteq \tilde{A}$  is an essential ideal.

**Definition.** Let  $A$  be an algebra,  $J \subseteq A$  an ideal. We say  $J$  is essential if for any other ideal  $I \subseteq A$ ,  $I \cap J \neq \{0\}$ .

**Proposition:** Let  $A$  be a complex algebra.

(1) The set  $A \times \mathbb{C}$ , equipped with

$$\begin{aligned} (a, \alpha) + (b, \beta) &= (a + b, \alpha + \beta) \\ z(a, \alpha) &= (za, z\alpha) \\ (a, \alpha)(b, \beta) &= (ab + \beta a + \alpha b, \alpha\beta) \end{aligned}$$

is a unital algebra, with unit  $1_{\tilde{A}} = (1, 0)$ . We denote this algebra  $\tilde{A}$ .

<sup>iii</sup>Just find  $f \in A$  such that  $h(f) = k$ , then take  $1 = h(f/k)$ .

(2) If  $A$  is a  $*$ -algebra, then  $\tilde{A}$  is a  $*$ -algebra, with

$$(a, \alpha)^* = (a^*, \alpha).$$

(3) The map  $\iota_A: A \rightarrow \tilde{A}$ , given by  $\iota_A(a) = (a, 0)$  is an injective  $*$ -homomorphism, and  $\pi_A: \tilde{A} \rightarrow \mathbb{C}$  is a surjective  $*$ -homomorphism.

The image,  $\iota_A(A) \subseteq \tilde{A}$  is a maximal  $*$ -ideal.

This yields an exact sequence of  $*$ -algebras:

$$0 \longrightarrow A \xrightarrow{\iota_A} \tilde{A} \xrightarrow{\pi_A} \mathbb{C} \longrightarrow 0$$

(4) If  $A$  is nonunital, then  $\iota_A(A) \subseteq \tilde{A}$  is an essential ideal.

*Proof.* We will prove (3) and (4).

(3) From the definition, we see that  $\iota_A$  is an injective  $*$ -homomorphism, and  $\pi_A$  is a surjective  $*$ -homomorphism, with  $\text{Ran}(\iota_A) = \ker(\pi_A)$ . Thus, by the first isomorphism theorem, we have  $\tilde{A}/\text{Ran}(\iota_A) \cong \mathbb{C}$ , so the  $*$ -ideal,  $\text{Ran}(\iota_A)$ , is maximal in  $\tilde{A}$ .

(4) Let  $I \subseteq \tilde{A}$  be a nonzero ideal, and let  $0 \neq (a, \alpha) \in I$ . If  $\alpha = 0$ , then  $0 \neq (a, 0) \in \iota(A) \cap I$ .

If  $a = 0$ , then  $\alpha \neq 0$ , so  $1_{\tilde{A}} = (0, 1) = \alpha^{-1}(0, \alpha) \in I$ , so  $I = \tilde{A}$ , so  $\iota(A) \cap I = \iota(A)$ .

We assume  $a, \alpha \neq 0$ . Multiplying by  $\alpha^{-1}$ , setting  $b = \alpha^{-1}a$ , we get  $(b, 1) \in I$ , and since  $I$  is an ideal, we have  $(xb + x, 0) \in I$  and  $(bx + x, 0) \in I$ . If  $xb + x = bx + x = 0$ , then  $(-b)$  is a multiplicative unit for  $A$ , which contradicts the fact that  $A$  is nonunital. Thus, there must be  $x \in A$  such that  $xb + x \neq 0$  or  $bx + x \neq 0$ . Thus,  $I \cap \iota(A) \neq \{0\}$ , so  $\iota(A)$  is an essential ideal.

□

When we talk about elements of  $\tilde{A}$ , we write  $(a, \alpha) = a + \alpha 1_{\tilde{A}}$ .

**Proposition:** Let  $A$  and  $B$  be  $*$ -algebras, and let  $\phi: A \rightarrow B$  be a  $*$ -homomorphism.

- (1) The map  $\tilde{\phi}(a, z) = (\phi(a), z)$  is a unital  $*$ -isomorphism that extends  $\phi$ . Moreover,  $\tilde{\phi}$  is injective (or surjective) if and only if  $\phi$  is injective (or surjective).
- (2) If  $B$  is unital, the map  $\bar{\phi}(a, z) = \phi(a) + z 1_B$  is a unital  $*$ -homomorphism that extends  $\phi$ . If  $A$  is nonunital, and  $\phi$  is injective, then so is  $\bar{\phi}$ .
- (3) If  $A$  is nonunital, and  $h: A \rightarrow \mathbb{C}$  is a character on  $A$ , then  $\bar{h}(a, \alpha) = h(a) + \alpha$  is a character on  $\tilde{A}$  extending  $h$ .

*Proof.* We will prove (3).

(3) If  $h$  is a character on  $A$ , then  $|\ker(h)| \neq A$ . If it were the case that  $\ker(\bar{h}) = \tilde{A}$ , then since  $A \subseteq \tilde{A}$  is an essential ideal,  $\ker(h) = \ker(\bar{h}) \cap A = \tilde{A} \cap A = A$ , which is a contradiction. Thus,  $\bar{h}$  is a character.

□

**Proposition:** Let  $X$  be a noncompact LCH space, and let  $X_\infty$  be the one-point compactification of  $X$ . There is a unital  $*$ -homomorphism  $\varphi: \widetilde{C_0(X)} \rightarrow C(X_\infty)$  that maps  $C_0(X)$  onto the ideal  $I = \{f \mid f(\infty) = 0\} \subseteq C(X_\infty)$ .

*Proof.* If  $f \in C_0(X)$ , we start by showing that  $\phi: X_\infty \rightarrow \mathbb{C}$ , given by

$$\phi(f)(x) = \begin{cases} f(x) & x \in X \\ 0 & x \in \infty \end{cases}$$

is continuous on  $X_\infty$ . It is the case that  $\phi(f)$  is continuous on  $X$ , since  $\phi(f)|_X = f$ , and  $X \subseteq X_\infty$  is open. Let  $(x_i)_i$  be a net in  $X_\infty$  converging to  $\infty$ , and let  $\varepsilon > 0$ . Since  $f$  vanishes at infinity, there is a compact subset  $K \subseteq X$  such that  $|f(x)| < \varepsilon$ , for  $x \notin K$ . The set  $X_\infty \setminus K$  is an open neighborhood of  $\infty$ , so  $x_i \in X_\infty \setminus K$  for large  $i$ . Thus,

$$(\phi(f)(x_i))_i \rightarrow 0 = \phi(f)(\infty).$$

We can see that  $\phi$  is a  $*$ -homomorphism by the way we have defined it, and that  $0 = \phi(f)(x)$  if and only if  $f = 0$  for all  $x$ , so  $\phi(f)$  is an injective  $*$ -homomorphism.

We will show that  $\text{Ran}(\phi) = I$ . Let  $g \in I$ . We have  $g|_X: X \rightarrow \mathbb{C}$  vanishes at infinity. Given  $\varepsilon > 0$ , since  $g(\infty) = 0$ , there is a neighborhood  $V$  of  $\infty$  with  $|g| < \varepsilon$  on  $V$ . This means we find compact  $K \subseteq X$  such that  $X \setminus K \subseteq V$ , so  $|g(x)| < \varepsilon$  for  $x \notin K$ . Thus,  $g|_X \in C_0(X)$ . Thus,  $g = \phi(g|_X)$ , so  $\text{Ran}(\phi) = I$ .

Since  $C_0(X)$  is nonunital, the extension  $\varphi: \widetilde{C_0(X)} \rightarrow C(X_\infty)$  is also injective. We will show that  $\varphi$  is onto. If  $k \in C(X_\infty)$ , then  $g = k - k(\infty)\mathbb{1}_{X_\infty} \in I$ , so there is  $f \in C_0(X)$  with  $\phi(f) = g$ . Thus,

$$\begin{aligned} \varphi(k, k(\infty)) &= \phi(k) + k(\infty)\mathbb{1}_{X_\infty} \\ &= g + k(\infty)\mathbb{1}_{X_\infty} \\ &= k. \end{aligned}$$

□

We have seen the character space on  $C(X)$  earlier when  $X$  is compact Hausdorff; now, we can see the character space on  $C_0(X)$ , where  $X$  is a LCH space.

**Corollary:** Let  $X$  be a LCH space. If  $x \in X$ , then  $\delta_x: C_0(X) \rightarrow \mathbb{C}$ , given by  $\delta_x(f) = f(x)$  is a character on  $C_0(X)$ . Moreover, the map  $\delta: X \rightarrow \Omega(C_0(X))$ , given by  $x \mapsto \delta_x$  is a bijection.

*Proof.* Each  $\delta_x: C_0(X) \rightarrow \mathbb{C}$  is a character, and  $\delta_x \neq 0$  by Urysohn's lemma.

Let  $h: C_0(X) \rightarrow \mathbb{C}$  be a character. The unitization,  $\bar{h}: \widetilde{C_0(X)} \rightarrow \mathbb{C}$  is a character. Let  $\varphi: \widetilde{C_0(X)} \rightarrow C(X_\infty)$  be the  $*$ -isomorphism to the one-point compactification of  $X$ . Thus, there is a  $\xi \in X_\infty$  with  $\delta_\xi = \bar{h} \circ \varphi^{-1}$ .

Thus, we see that  $\delta_\xi \circ \phi = \delta_\xi \circ \varphi \circ \iota = \bar{h} \circ \iota = h$  on  $C_0(X)$ , where  $\iota: C_0(X) \rightarrow \widetilde{C_0(X)}$  is the natural inclusion. Since  $h \neq 0$  and  $\phi(f)(\infty) = 0$  for every  $f \in C_0(X)$ , we must have  $\xi = x \in X$ , so

$$\begin{aligned} h(f) &= \delta_x \circ \phi(f) \\ &= f(x) \\ &= \delta_x(f) \end{aligned}$$

for every  $f \in C_0(X)$ , so  $\delta$  is onto. Since  $C_0(X)$  separates points,  $\delta$  is injective. □

We are interested in identifying the character space of a nonunital algebra with a subset of the character space of its unitization. Note that the projection  $\pi: \widetilde{A} \rightarrow \mathbb{C}$  given by  $(a, \alpha) \mapsto \alpha$  is a character, but  $\pi|_A$  is not a character on  $A$ . This is indeed the desired character that will extend the character space on  $A$ .

**Proposition:** Let  $A$  be a nonunital algebra. The map  $\Omega(A) \rightarrow \Omega(\widetilde{A})$ , given by  $h \mapsto \bar{h}$ , is injective.

Moreover,  $\Omega(\widetilde{A}) = \Omega(A) \cup \{\pi\}$ , where  $\pi: \widetilde{A} \rightarrow \mathbb{C}$  is given by  $(a, \alpha) \mapsto \alpha$ .

*Proof.* It is clear from the definition of  $\bar{h}$  that  $h \mapsto \bar{h}$  is well-defined and injective. Moreover, if  $\phi \in \Omega(\tilde{A})$  is such that  $h = \phi|_A \neq 0$ , then  $h \in \Omega(A)$  is such that  $\bar{h} = \phi$ . Since  $\phi$  is unital, we have

$$\begin{aligned}\phi(a + \alpha 1_{\tilde{A}}) &= \phi(a) + \alpha \\ &= h(a) + \alpha \\ &= \bar{h}(a + \alpha 1_{\tilde{A}}).\end{aligned}$$

Thus, if  $\phi \in \Omega(\tilde{A})$  is such that  $\phi|_A = 0$ , then  $\phi = \pi$ . □

## Algebraic Spectrum

**Definition.** Let  $A$  be a unital  $\mathbb{F}$ -algebra. Let  $a \in A$ .

(1) The resolvent of  $a$  is the set

$$\rho(a) = \{\lambda \in \mathbb{F} \mid a - \lambda 1_A \in GL(A)\}.$$

(2) The spectrum of  $a$  is the complement of the resolvent,  $\sigma(a) = \mathbb{F} \setminus \rho(a)$ .

We may refine the definition for the case of a nonunital algebra.

**Definition.** Let  $A$  be a nonunital algebra, and let  $a \in A$ . We define

$$\rho(a) = \rho(\iota_A(a)),$$

where  $\iota_A : A \hookrightarrow \tilde{A}$  is the canonical embedding of  $A$  into its unitization. We define  $\sigma(a) = \mathbb{F} \setminus \rho(a)$ .

**Exercise:** Prove that similar elements in an algebra have the same resolvent.

**Solution:** Let  $a = zbz^{-1}$  for some  $z \in GL(\tilde{A})$ . We will show that  $\rho(a) = \rho(b)$ .

Let  $\lambda \in \rho(a)$ . Then,  $a - \lambda 1_{\tilde{A}} \in GL(\tilde{A})$ . Note that since  $z \in GL(\tilde{A})$ , and  $GL(\tilde{A})$  is a group, we have  $z(a - \lambda 1_{\tilde{A}})z^{-1} \in GL(\tilde{A})$ , so  $b - \lambda 1_{\tilde{A}} \in GL(\tilde{A})$ .

**Example.** If  $A = \mathbb{C}$ , then for every  $z \in \mathbb{C}$ , the spectrum  $\sigma(z) = \{z\}$ . This is because every non-zero number is invertible.

**Example.** Let  $A = \text{Mat}_n(\mathbb{C})$ . Then,  $\lambda$  is an eigenvalue for a matrix  $a \in A$  if and only if  $a - \lambda I_n$  is not invertible. In this case, we say  $\sigma(a) = \sigma_p(a)$ , the point spectrum of  $a$ .

**Example.** Let  $A = \ell_\infty(\Omega)$ . We find

$$\begin{aligned}\lambda \in \rho(f) &\Leftrightarrow f - \lambda 1_\Omega \in GL(A) \\ &\Leftrightarrow \inf_{x \in \Omega} |f(x) - \lambda| > 0 \\ &\Leftrightarrow \lambda \notin \overline{\text{Ran}(f)}.\end{aligned}$$

Thus,  $\sigma(f) = \overline{\text{Ran}(f)}$ .

**Example.** Let  $\Omega$  be compact Hausdorff, and let  $A = C(\Omega)$ . If  $f \in A$ , we have

$$\begin{aligned}\lambda \in \rho(f) &\Leftrightarrow f - \lambda 1_\Omega \in GL(A) \\ &\Leftrightarrow f(x) - \lambda \neq 0 \\ &\Leftrightarrow \lambda \notin \text{Ran}(f).\end{aligned}$$

Thus,  $\sigma(f) = \text{Ran}(f)$ .

If  $\Omega$  is a noncompact LCH space with one-point compactification  $\Omega_\infty$ , then if  $A = C_0(\Omega)$ , we find

$$\sigma(f) = \text{Ran}(f) \cup \{0\}.$$

**Example.** Let  $A = L_\infty(\Omega, \mu)$ , where  $(\Omega, \mu)$  is a measure space. Fix  $f \in A$ . To understand  $\sigma(f)$ , we must understand the essential range of  $f$ .

Write  $U(z, \varepsilon)$  for the open disk centered at  $z \in \mathbb{C}$  with radius  $\varepsilon > 0$ . The essential range of  $f$  is

$$\text{ess ran}(f) = \left\{ \lambda \in \mathbb{F} \mid \mu\left(f^{-1}(U(\lambda, \varepsilon))\right) > 0, \forall \varepsilon > 0 \right\}.$$

The essential range can also be viewed as the support of the pushforward measure  $f_*\mu$  on  $\mathbb{C}$ , where  $f_*\mu(E) = \mu(f^{-1}(E))$  for measurable  $E \subseteq \mathbb{C}$ .

A number  $\lambda \notin \text{ess ran}(f)$  if and only if  $f - \lambda 1_A$  is essentially bounded below. Thus, we see that  $\lambda \in \rho(f)$  if and only if  $\lambda \notin \text{ess ran}(f)$ . Thus,  $\sigma(f) = \text{ess ran}(f)$ .

**Proposition:** Let  $\varphi: A \rightarrow B$  be a unital algebra homomorphism. Then,  $\sigma(\varphi(a)) \subseteq \sigma(a)$ , with equality if  $\varphi$  is bijective.

*Proof.* If  $\lambda \in \rho(a)$ , then  $a - \lambda 1_A \in GL(A)$ . Thus,  $\varphi(a) - \lambda 1_B \in GL(B)$ , so  $\lambda \in \rho(\varphi(a))$ . Thus,  $\rho(a) \subseteq \rho(\varphi(a))$ , so  $\sigma(\varphi(a)) \subseteq \sigma(a)$ .

If  $\varphi$  is bijective, then  $\varphi^{-1}: B \rightarrow A$  is also a unital homomorphism, meaning

$$\begin{aligned} \sigma(a) &= \sigma\left(\varphi^{-1}(\varphi(a))\right) \\ &\subseteq \sigma(\varphi(a)). \end{aligned}$$

□

**Corollary:** If  $A$  is an algebra, and  $h \in \Omega(A)$ , then  $h(a) \in \sigma(a)$ . We have

$$\{h(a) \mid h \in \Omega(A)\} \subseteq \sigma(a).$$

*Proof.* Assume  $A$  is unital. Taking  $B = \mathbb{C}$  in the above proposition, and noting  $\sigma(z) = \{z\}$ , we prove the statement.

If  $A$  is nonunital, then  $h(a) = \overline{h}(\iota_A(a))$ .

□

**Fact.** Let  $A$  be a unital algebra with  $a, b \in A$ . Then,  $\sigma(ab) \cup \{0\} = \sigma(ba) \cup \{0\}$ .

*Proof.* If  $\lambda \neq 0$  belongs to  $\rho(ab)$ , then  $ab - \lambda 1_A \in GL(A)$ , meaning  $\lambda^{-1}ab - 1_A \in GL(A)$ . Thus,  $\lambda^{-1}ba - 1_A \in GL(A)$ , so  $ba - \lambda 1_A \in GL(A)$ , so  $\lambda \in \rho(ba)$ . Thus,  $\rho(ba) \setminus \{0\} \subseteq \rho(ab) \setminus \{0\}$ , so, repeating this argument, we get  $\rho(ba) \setminus \{0\} = \rho(ab) \setminus \{0\}$ .

□

**Proposition (Spectral Mapping):** Let  $A$  be a unital  $\mathbb{C}$ -algebra, and let  $a \in A$  with  $\sigma(a) \neq \emptyset$ . If  $p \in \mathbb{C}[x]$ , then  $\sigma(p(a)) = p(\sigma(a))$ .

*Proof.* Splitting  $p$ , we take

$$p(z) = \alpha \prod_{j=1}^n (z - \alpha_j)$$

for  $\alpha, \alpha_1, \dots, \alpha_n \in \mathbb{C}$ . Then,

$$p(a) = \alpha \prod_{j=1}^n (a - \alpha_j 1_A).$$

The factors  $a - \alpha_j 1_A$  commute, and since  $GL(A)$  is a group, we get

$$p(a) \in GL(A) \Leftrightarrow a - \alpha_j 1_A \in GL(A)$$

$$\begin{aligned}
&\Leftrightarrow \alpha_j \in \rho(a) \\
&\Leftrightarrow \alpha_j \notin \sigma(a) \\
&\Leftrightarrow p(z) \neq 0 \forall z \in \sigma(A) \\
&\Leftrightarrow 0 \notin p(\sigma(a)).
\end{aligned}$$

Thus,  $p(a) \notin GL(A)$  if and only if  $0 \in p(\sigma(a))$ . □

## Banach and $C^*$ -Algebras

In the notes on Hilbert space operators, we established the spectral theorem for compact normal operators. In order to establish the spectral theorem for all normal operators, we will study the unital  $C^*$ -algebra generated by the normal operator. This will hinge on understanding the abstract theory of Banach algebras and  $C^*$ -algebras.

We start with some examples of Banach and  $C^*$ -algebras, as well as discussing some constructions of and with  $C^*$ -algebras.

### Examples

**Definition.** A Banach  $*$ -algebra is a Banach algebra  $A$  with an involution map  $A \rightarrow A, a \mapsto a^*$ , satisfying

$$\|a^*\| = \|a\|.$$

If  $A$  is a Banach  $*$ -algebra that satisfies the  $C^*$  property,  $\|a^*a\| = \|a\|^2$ , for every  $a \in A$ , then  $A$  is called a  $C^*$ -algebra.

We know that  $*$ -algebras admit a variety of distinguished elements. We can add two more to that list.

**Definition.** Let  $A$  be a  $C^*$ -algebra, and  $w \in A$ .

- We say  $w$  is an isometry if  $w^*w = 1_A$ .
- If  $w$  is an isometry, and  $ww^* \neq 1_A$ , then we say  $w$  is a proper isometry.

We may also speak of partial isometries.

**Lemma:** If  $A$  be a  $C^*$ -algebra with  $v \in A$ . The following are equivalent:

- (i)  $v^*v$  is a projection;
- (ii)  $vv^*v = v$ ;
- (iii)  $vv^*$  is a projection;
- (iv)  $v^*vv^* = v^*$ .

Such an element is called a partial isometry.

*Proof.* We obtain the implication (i) implying (ii) through verifying

$$(vv^*v - v)^*(vv^*v - v) = 0,$$

meaning  $vv^*v - v = 0$ . Similarly, the implication (iii) implying (iv) is similar. □

**Example.** The complex numbers  $\mathbb{C}$  with involution  $z \mapsto \bar{z}$  and norm  $z \mapsto |z|$  is a  $C^*$ -algebra.

**Example.** We know that  $\mathcal{B}(\mathcal{H})$ , the space of bounded linear operators on a Hilbert space, is a  $C^*$ -algebra.

**Example.** If  $n \geq 2$ , then  $\text{Mat}_n(\mathbb{C})$  is a unital noncommutative  $*$ -algebra. We know that  $(\text{Mat}_n(\mathbb{C}), \|\cdot\|_{\text{op}})$  is a Banach space.

We want to show that  $(\text{Mat}_n(\mathbb{C}), \|\cdot\|_{\text{op}})$  is a  $C^*$ -algebra isomorphic to  $\mathcal{B}(\ell_2^n)$ .

We can establish a unital isomorphism  $\text{Mat}_n(\mathbb{C}) \rightarrow \mathcal{L}(\mathbb{C}^n)$  by sending the matrix  $a$  to its corresponding transformation  $T_a$ .

Since  $\mathbb{C}^n$  is a finite-dimensional Hilbert space,  $\mathcal{B}(\ell_2^n) = \mathcal{L}(\mathbb{C}^n)$ . We have a unital isomorphism of algebras  $T(a) = T_a$  between  $\text{Mat}_n(\mathbb{C})$  and  $\mathcal{B}(\ell_2^n)$ .

By the definition of the operator norm,  $\|a\|_{\text{op}} = \|T_a\|_{\text{op}}$ , so  $T: \text{Mat}_n(\mathbb{C}) \rightarrow \mathcal{B}(\ell_2^n)$  is an isometry.

If  $a, b \in \text{Mat}_n(\mathbb{C})$ , then

$$\begin{aligned} \|ab\|_{\text{op}} &= \|T_{ab}\|_{\text{op}} \\ &= \|T_a T_b\|_{\text{op}} \\ &\leq \|T_a\|_{\text{op}} \|T_b\|_{\text{op}} \\ &= \|a\|_{\text{op}} \|b\|_{\text{op}}. \end{aligned}$$

Next,  $\|I_n\|_{\text{op}} = \|T_{I_n}\|_{\text{op}} = \|\text{id}_{\ell_2^n}\|_{\text{op}} = 1$ , and

$$\begin{aligned} \|a^*\|_{\text{op}} &= \|T_{a^*}\|_{\text{op}} \\ &= \|T_a^*\|_{\text{op}} \\ &= \|T_a\|_{\text{op}} \\ &= \|a\|_{\text{op}}. \end{aligned}$$

Similarly,

$$\|a^* a\|_{\text{op}} = \|a\|_{\text{op}}^2.$$

Thus,  $(\text{Mat}_n(\mathbb{C}), \|\cdot\|_{\text{op}})$  is a  $C^*$ -algebra.

**Example.** The space  $\ell_\infty(\Omega)$  of bounded functions on  $\Omega$  is a unital and commutative  $*$ -algebra under point-wise operations, which is also a Banach space under  $\|\cdot\|_{\text{u}}$ .

We can also see that  $\|fg\|_{\text{u}} \leq \|f\|_{\text{u}} \|g\|_{\text{u}}$ , and  $\|f^* f\|_{\text{u}} = \|f\|_{\text{u}}^2$  for all  $f, g \in \ell_\infty(\Omega)$ , meaning  $\ell_\infty(\Omega)$  is a unital and commutative Banach algebra.

Finally,

$$\begin{aligned} \|f^* f\|_{\text{u}} &= \sup_{x \in \Omega} |(f^* f)(x)| \\ &= \sup_{x \in \Omega} |f^*(x) f(x)| \\ &= \sup_{x \in \Omega} |\overline{f(x)} f(x)| \\ &= \sup_{x \in \Omega} |f(x)|^2 \\ &= \|f\|_{\text{u}}^2. \end{aligned}$$



**Lemma:** Let  $B$  be a Banach algebra/Banach  $*$ -algebra/ $C^*$ -algebra. If  $A \subseteq B$  is a norm closed subalgebra/ $*$ -subalgebra, then  $A$  is a Banach algebra/Banach  $*$ -algebra/ $C^*$ -algebra.

**Definition.** If  $B$  is a  $C^*$ -algebra, and  $A \subseteq B$  is a norm-closed  $*$ -subalgebra, then  $A$  is a  $C^*$ -subalgebra of  $B$ .

If  $B$  is unital, then  $A \subseteq B$  is a unital  $C^*$ -subalgebra if  $1_B \in A$ .

If  $\mathcal{H}$  is a Hilbert space, then a  $C^*$ -subalgebra  $A \subseteq \mathcal{B}(\mathcal{H})$  is sometimes called a concrete  $C^*$ -algebra.

**Example.** The compact operators,  $\mathcal{K}(\mathcal{H})$  is an operator norm-closed  $*$ -subalgebra of  $\mathcal{B}(\mathcal{H})$ . It is unital if and only if  $\dim(\mathcal{H}) < \infty$ .

**Example.** Let  $(\Omega, \mathcal{M})$  be a measurable space, The bounded measurable functions,  $B_\infty(\Omega)$ , is a unital  $*$ -subalgebra.

Equipped with the  $\infty$  norm,  $B_\infty(\Omega)$  is a Banach space, meaning  $B_\infty(\Omega) \subseteq \ell_\infty(\Omega)$  is norm-closed, and is thus a unital commutative  $C^*$ -algebra.

**Example.** Let  $(\Omega, \mathcal{M}, \mu)$  be a measure space. The essentially bounded functions,  $L_\infty(\Omega, \mu)$ , is a Banach space with the ess sup norm. It is also the case that  $L_\infty(\Omega, \mu)$  is a unital commutative  $*$ -algebra. We can show that  $\|f^*f\|_\infty = \|f\|_\infty^2$ , so  $L_\infty(\Omega, \mu)$  is a unital  $C^*$ -algebra.

**Example.** Let  $\Omega$  be a LCH space. We know that  $C_b(\Omega)$  and  $C_0(\Omega)$  are  $*$ -subalgebras of  $\ell_\infty(\Omega)$ . We also know these are uniform norm-closed, meaning  $C_b(\Omega)$  and  $C_0(\Omega)$  are  $C^*$ -algebras. The  $C^*$ -algebra  $C_b(\Omega)$  is always unital, but  $C_0(\Omega)$  is unital if and only if  $\Omega$  is compact.

Note that if  $\Omega$  is given the discrete topology, then  $\ell_\infty(\Omega) = C_b(\Omega)$ , as any function on a discrete space is continuous.

The map  $C_b(\Omega) \rightarrow C(\beta\Omega)$ , given by  $f \mapsto f^\beta$  is an isometric isomorphism of Banach spaces. We can also verify that this is a  $*$ -isomorphism, as  $(fg)^\beta = f^\beta g^\beta$ , as these agree on the dense subset  $\Delta(\Omega) \subseteq \beta\Omega$ . Similarly,  $(f^*)^\beta = (f^\beta)^*$ . Thus,  $C_b(\Omega)$  and  $C(\beta\Omega)$  are isomorphic as  $C^*$ -algebras.

We get the isometric  $*$ -isomorphism  $\ell_\infty = C_b(\mathbb{N}) = C(\beta\mathbb{N})$ .

## Constructions

We are interested in constructing new  $C^*$ -algebras from old.

### Generating Sets

We may start with closures.

**Lemma:** Let  $B$  be a Banach algebra/Banach  $*$ -algebra, and let  $A \subseteq B$  be a subalgebra/ $*$ -subalgebra. The closure,  $\overline{A} \subseteq B$ , is a Banach subalgebra/ $*$ -subalgebra.

If  $B$  is a  $C^*$ -algebra with  $A \subseteq B$  a  $*$ -subalgebra, then  $\overline{A}$  is a  $C^*$ -subalgebra of  $B$ .

Given a collection of operators  $S \subseteq \mathcal{B}(\mathcal{H})$ , we are interested in constructing the picture of the smallest  $C^*$ -subalgebras of  $\mathcal{B}(\mathcal{H})$  containing  $S$ . In a more general case, we may consider any  $C^*$ -algebra  $B$  and the subset  $S \subseteq \mathcal{B}(\mathcal{H})$ .

**Definition.** Let  $B$  be a Banach algebra/ $*$ -algebra, and let  $S \subseteq B$  be any subset. The Banach algebra/ $*$ -algebra generated by  $S$  is the smallest Banach subalgebra/ $*$ -subalgebra containing  $S$ .

If  $B$  is a  $C^*$ -algebra, then the  $C^*$ -subalgebra generated by  $S$  is the smallest  $C^*$ -subalgebra of  $B$  containing  $S$ , denoted

$$C^*(S) = \bigcap \{A \mid S \subseteq A, A \subseteq B \text{ is a } C^*\text{-subalgebra}\}.$$

Notationally, we write  $C^*(a_1, \dots, a_n)$  if  $\{a_1, \dots, a_n\}$  is a finite subset of  $B$ .

Obviously, we need a more workable picture of the  $C^*$ -subalgebra generated by a set, at the very least we need something we can imagine.

**Lemma:** Let  $B$  be a Banach algebra and suppose  $S \subseteq B$  is any subset.

- (1) The Banach algebra generated by  $S$  is the closed span of the set of finite words in  $S$ . In other words, it is equal to  $\overline{\text{span}}(W)$ , where

$$W = \{x_1 x_2 \cdots x_n \mid n \in \mathbb{N}, x_j \in S\}.$$

- (2) If  $B$  is a Banach  $*$ -algebra or  $C^*$ -algebra, then the Banach  $*$ -algebra or  $C^*$ -algebra generated by  $S$  is the closed span of the set of finite words in  $S$  and  $S^*$ . In other words, it is equal to  $\overline{\text{span}}(W)$ , where

$$W = \{x_1 x_2 \cdots x_n \mid n \in \mathbb{N}, x_j \in S \cup S^*\}.$$

*Proof.* We will prove (2).

Note that  $S$  is closed under multiplication and involution, so  $\text{span}(W)$  is a  $*$ -algebra containing  $S$ , so  $\overline{\text{span}}(W)$  is a  $C^*$ -subalgebra of  $B$  containing  $S$ , so  $C^*(S) \subseteq \overline{\text{span}}(W)$ .

In the reverse inclusion, any  $C^*$ -subalgebra of  $B$  containing  $S$  must contain  $\text{span}(W)$ , so  $\overline{\text{span}}(W) \subseteq C^*(S)$ .  $\square$

**Proposition:** Let  $B$  be a  $C^*$ -algebra, and suppose  $a \in B$  is a normal element. The  $C^*$ -algebra generated by  $a$ ,  $C^*(a)$ , is a commutative  $C^*$ -subalgebra. If  $B$  is unital, then  $C^*(a, 1_B)$  is a unital commutative  $C^*$ -algebra.

*Proof.* We see that, in the notation of the lemma, if  $S = \{a\}$  or  $S = \{1_B, a\}$ , if  $w_1, w_2 \in W$ , we have  $w_1 w_2 = w_2 w_1$  (since  $aa^* = a^*a$ ), so  $\text{span}(W)$  is a commutative  $*$ -subalgebra, hence  $\overline{\text{span}}(W)$  is commutative.  $\square$

**Example.** Let  $\iota: [0, 1] \rightarrow \mathbb{C}$  be the inclusion map,  $\iota(t) = t$ . By the Stone–Weierstrass theorem, we have

$$\begin{aligned} C^*(\iota, \mathbb{1}_{[0,1]}) &= C([0, 1]) \\ C^*(\iota) &= \{f \in C([0, 1]) \mid f(0) = 0\} \\ &\cong C_0((0, 1]). \end{aligned}$$

Note that if  $\iota: \mathbb{T} \rightarrow \mathbb{C}$  is the inclusion  $\iota(z) = z$ , then  $C^*(\iota) = C^*(\mathbb{T})$ .

**Exercise:** Let  $\Delta$  be the Cantor set. Let

$$\mathcal{C} = \{\mathbb{1}_C \mid C \subseteq \Delta \text{ is clopen}\}.$$

Show that  $C^*(\mathcal{C}) = C(\Delta)$ .

**Solution:** Since  $\mathcal{C}$  separates points and contains the constant function  $\mathbb{1}_\Delta$ , the Stone–Weierstrass theorem provides that  $C^*(\mathcal{C}) = C(\Delta)$ .

**Definition.** Recall that the definition of the right shift is such that  $R^* = L$ , where  $L$  is the left shift. We know that  $R^*R = I$ , but  $RR^* \neq I$ , since it has a nontrivial kernel.

The Toeplitz algebra is the  $C^*$ -algebra generated by the right shift. In other words,

$$\mathcal{T} = C^*(R).$$

**Exercise:** Prove that the Toeplitz algebra contains the compact operators.

**Solution:** We start by showing that the rank-one projection of  $e_j$  onto  $e_i$ , where  $(e_n)_n$  are the canonical orthonormal basis of  $\ell_2$ , is generated by the right shift as follows.

$$\theta_{e_i, e_j} = R^{i-1}(I - RR^*)(R^*)^{j-1}.$$

Note that we only need to show this equivalence when applied to  $e_n$ :

$$\begin{aligned}\theta_{e_i, e_j}(e_n) &= \langle e_n, e_j \rangle e_i \\ &= \delta_{nj} e_i.\end{aligned}$$

Applying in steps, we start with

$$\begin{aligned}R^{i-1}(I - RR^*)(R^*)^{j-1}(e_n) &= R^{i-1}(I - RR^*)(e_{n-j+1}) \\ &= \begin{cases} R^{i-1}(e_{n-j+1}) & n = j \\ R^{i-1}(0) & n \neq j \end{cases} \\ &= \delta_{nj} e_i.\end{aligned}$$

Thus, since the rank-one projections are contained in the Toeplitz algebra, the finite-rank operators are contained in the Toeplitz algebra, hence the compact operators are contained in the Toeplitz algebra.

**Example.** Consider the following isometries on  $\ell_2$ :

$$\begin{aligned}V(\alpha_1, \alpha_2, \alpha_3, \dots) &= (\alpha_1, 0, \alpha_2, 0, \alpha_3, 0, \dots) \\ W(\alpha_1, \alpha_2, \alpha_3, \dots) &= (0, \alpha_1, 0, \alpha_2, 0, \alpha_3, \dots).\end{aligned}$$

The operators  $V$  and  $W$  satisfy

$$\begin{aligned}V^*V &= I \\ W^*W &= I \\ VV^* + WW^* &= I.\end{aligned}$$

The Cuntz algebra,  $\mathcal{O}_2$ , is the  $C^*(V, W)$ .

## Products, Sums, and Quotients

In the category of  $C^*$ -algebras, we can also look at products and coproducts.

**Definition.** Let  $\{A_i\}_{i \in I}$  be a family of Banach algebras/Banach  $*$ -algebras/ $C^*$ -algebras. Then, we define the following two constructions with pointwise operations and the  $\infty$  norm.

- (1) The  $\ell_\infty$  product is defined as

$$\prod_{i \in I} A_i = \left\{ (a_i)_i \mid a_i \in A_i, \|(a_i)_i\| = \sup_{i \in I} \|a_i\| < \infty \right\}.$$

- (2) For the case of  $I = \mathbb{N}$ , we may consider the  $c_0$  sum

$$\bigoplus_{n \in \mathbb{N}} A_n = \left\{ a = (a_n)_n \mid a_n \in A_n, \lim_{n \rightarrow \infty} \|a_n\| = 0 \right\}$$

as a subset of the  $\ell_\infty$  product of  $\{A_n\}_{n \in \mathbb{N}}$ . This is a closed  $*$ -ideal.

- (3) In the case where  $I = \mathbb{N}$  and  $A_n = A$  is fixed, we write  $\ell_\infty(A) = \prod_{n \in \mathbb{N}} A_n$  and  $c_0(A) = \bigoplus_{n \in \mathbb{N}} A_n$ .

- (4) For a finite family  $\{A_n\}_{n=1}^N$ , the  $c_0$  sum equals the  $\ell_\infty$  product. We decorate the notation to write  $A_1 \oplus_\infty \dots \oplus_\infty A_N$ .

**Example.** For  $n_1, \dots, n_r \in \mathbb{N}$ , the  $C^*$ -algebra

$$M = \text{Mat}_{n_1}(\mathbb{C}) \oplus_{\infty} \text{Mat}_{n_2} \oplus_{\infty} \dots \oplus_{\infty} \text{Mat}_{n_r}(\mathbb{C})$$

is finite-dimensional. It is actually the case that every finite-dimensional  $C^*$ -algebra is of this form.

We can also take quotients.

**Proposition:** Let  $A$  be a normed  $*$ -algebra. Let  $I \subseteq A$  be a closed  $*$ -ideal. The quotient space  $A/I$  equipped with the quotient norm is a normed  $*$ -algebra.

If  $A$  is complete, then so is  $A/I$ . If  $A$  is commutative or unital, then so is  $A/I$ .

*Proof.* We know that  $A/I$  with its quotient norm is a normed vector space, and that  $A/I$  is a  $*$ -algebra. We need to show that the quotient norm is submultiplicative and that the involution is isometric.

Let  $a, b \in A$  and  $\varepsilon > 0$ . Then, there are  $x, y$  such that  $\|a + I\| + \varepsilon \geq \|a - x\|$ , and  $\|b + I\| + \varepsilon \geq \|b - y\|$ . Note that  $ay + xb - xy \in I$ , so

$$\begin{aligned} \|(a + I)(b + I)\| &= \|ab + I\| \\ &= \text{dist}_I(ab) \\ &\leq \|ab - (ay + xb - xy)\| \\ &= \|(a - x)(b - y)\| \\ &\leq \|a - x\| \|b - y\| \\ &\leq (\|a + I\| + \varepsilon)(\|b + I\| + \varepsilon). \end{aligned}$$

Sending  $\varepsilon \rightarrow 0$ , we get submultiplicativity. Regarding the involution, we get

$$\begin{aligned} \|(a + I)^*\| &= \|a^* + I\| \\ &= \inf_{x \in I} \|a^* - x\| \\ &= \inf_{y \in I} \|a^* - y^*\| \\ &= \inf_{y \in I} \|(a - y)^*\| \\ &= \inf_{y \in I} \|a - y\| \\ &= \|a + I\|. \end{aligned}$$

Completeness follows from the case of the quotient space in Banach spaces. □

## Ideals in $C_0(\Omega)$

Earlier, we characterized the maximal ideal space of  $C(\Omega)$ , where  $\Omega$  was compact Hausdorff. We are interested in applying this to characterizing the closed ideals of  $C_0(\Omega)$ , where  $\Omega$  is a LCH space.

**Definition.** Let  $\Omega$  be a LCH space.

- (a) For a subset  $K \subseteq \Omega$ , we write  $N_K$  to be continuous hull of  $K$ , i.e.

$$N_K = \{f \in C_0(\Omega) \mid f(x) = 0, \forall x \in K\}.$$

If  $K = \{x\}$ , we write  $N_x$ .

- (b) For any map  $f: \Omega \rightarrow \mathbb{C}$ , we denote the zero set of  $f$  by

$$Z(f) = f^{-1}(\{0\}).$$

(c) If  $I \subseteq C_0(\Omega)$  is any subset, the kernel of  $I$  is

$$\ker(I) = \bigcap_{f \in I} Z(f).$$

**Fact.**

- (1) If  $K \subseteq \Omega$  is nonempty, then  $N_K$  is a closed proper  $*$ -ideal in  $C_0(\Omega)$ .
- (2) If  $I \subseteq C_0$  is any subset, then  $\ker(I) \subseteq \Omega$  is closed.
- (3) If  $K \subseteq L \subseteq \Omega$ , then  $N_K \supseteq N_L$ .
- (4) If  $I \subseteq J \subseteq C_0(\Omega)$ , then  $\ker(I) \supseteq \ker(J)$ .

To show that every closed ideal in  $C_0(\Omega)$  is of the form  $N_K$  for some closed  $K \subseteq \Omega$ , we start with the case of  $C_c(\Omega)$ . We will finish the proof by taking closures.

**Lemma:** Let  $\Omega$  be a LCH space, and let  $I \subseteq C_0(\Omega)$  be an ideal. If  $g \in C_c(\Omega)$  with  $\text{supp}(g) \cap \ker(I) = \emptyset$ , then  $g \in I$ .

*Proof.* Let  $g \in C_c(\Omega)$  and  $C = \text{supp}(g)$ . For each  $x \in C$ , define  $h_x \in I$  such that  $h_x(x) \neq 0$  on  $C$ , and let  $U_x$  be the open neighborhood on which  $h_x \neq 0$ . The open cover  $\{U_x\}_{x \in C}$  admits a finite subcover,

$$C \subseteq \bigcup_{j=1}^n U_{x_j}.$$

We define the function

$$h = \sum_{j=1}^n |h_{x_j}|^2,$$

which belongs to  $I$  and is strictly positive on  $C$  by construction. Since  $C$  is compact,  $\inf_C(h) > 0$ . Let

$$f(x) = \begin{cases} \frac{g(x)}{h(x)} & x \in C \\ 0 & x \notin C \end{cases}.$$

Then,  $f$  is supported on  $C$ , and  $g = fh$ , so  $g \in I$ . □

**Proposition:** Let  $\Omega$  be a LCH space. If  $I \subseteq C_0(\Omega)$  is a closed proper ideal, then  $K = \ker(I)$ , and  $I = N_K$ .

*Proof.* Set  $J = \{g \in C_c(\Omega) \mid \text{supp}(g) \cap K = \emptyset\}$ . By the above lemma, we know that  $J \subseteq I$ . If  $K$  were empty, we would have  $J = C_c(\Omega)$ , implying

$$\begin{aligned} C_0(\Omega) &= \overline{C_c(\Omega)} \\ &= \overline{J} \\ &\subseteq \overline{I} \\ &= I, \end{aligned}$$

which would contradict the assumption that  $I$  is a proper ideal.

We can see that  $I \subseteq N_K$  by the definition of  $N_K$ . We will now show that every function in  $N_K$  can be approximated arbitrarily by a member in  $J$ . We will establish the reverse inclusion,  $J \subseteq N_K$ .

Let  $f \in N_K$ ,  $\varepsilon > 0$ , and set

$$C_\varepsilon = \{x \in \Omega \mid |f(x)| > \varepsilon\}.$$

Since  $f$  vanishes at infinity,  $C_\varepsilon$  is compact, and  $C_\varepsilon \cap K = \emptyset$ . By Urysohn's lemma, there is  $g \in C_c(\Omega, [0, 1])$  with  $g|_{C_\varepsilon} = 1$  and  $\text{supp}(g) \subseteq K^c$ . Thus,  $h = fg \in J$ , and  $\|f - h\|_\infty \leq \varepsilon$ . □

**Proposition:** Let  $\Omega$  be a LCH space. If  $K \subseteq \Omega$  is closed, then  $K = \ker(N_K)$ .

*Proof.* We can see that  $K \subseteq \ker(N_K)$ . If the inclusion is strict, then there is a point  $x \in \ker(N_K) \setminus K$ , and, by Urysohn's lemma, there is an  $f \in C_c(\Omega, [0, 1])$  with  $f|_K = 0$  and  $f(x) = 1$ . Thus,  $f \in N_K$ .

Since  $x \in \ker(N_K)$ , we must also have  $f(x) = 0$ , which is a contradiction. Thus,  $K = \ker(N_K)$ .  $\square$

We arrive at the following characterization of the closed ideals of  $C_0(\Omega)$ .

**Corollary:** Let  $\Omega$  be a LCH space. There is an order-reversing one-to-one correspondence between closed subsets of  $\Omega$  and closed ideals of  $C_0(\Omega)$ , given by

$$\Omega \supseteq K \leftrightarrow N_K \subseteq C_0(\Omega).$$

**Exercise:** Show that every maximal ideal of  $C_0(\Omega)$  is of the form  $N_x$ .

**Solution:** Via the containment ordering, we see that every maximal element of  $\Omega$  with this ordering is of the form  $\{x\}$ , meaning that every ideal of the form  $N_x$  is maximal.

Indeed, we may go further. Letting  $\Omega$  be a LCH space, and  $\Lambda \subseteq \Omega$  be open, we know that both  $\Lambda$  and  $\Lambda^c$  are locally compact. We can identify  $C_0(\Lambda)$  with the closed ideal  $N_K$ , where  $K = \Lambda^c$ .

Given  $f \in C_0(\Lambda)$ , define

$$f'(x) = \begin{cases} f(x) & x \in \Lambda \\ 0 & x \in \Lambda^c \end{cases}.$$

Clearly,  $f' \in C_0(\Omega)$ , and by definition,  $f \in N_K$ . Additionally, the inclusion map  $\iota: C_0(\Omega) \rightarrow N_K$ , defined by  $f \mapsto f'$ , is an isometric  $*$ -homomorphism.

**Exercise:** If  $g \in N_K$ , then  $g|_\Lambda \in C_0(\Lambda)$ , and  $(g|_\Lambda)' = g$ .

**Solution:** If  $g \in N_K$ , then  $g = 0$  on  $\Lambda^c$ , so for all  $\varepsilon > 0$ , there is some compact  $S \subseteq \Lambda$  such that  $|g|_{S^c} < \varepsilon$ . Thus,  $g \in C_0(\Lambda)$ .

By the definition of  $\iota$ , we must have  $g \mapsto g'$  is an isometric  $*$ -homomorphism, and since  $g$  is 0 on  $\Lambda^c$ , we have that  $(g|_\Lambda)' = g$ .

Thus, we come to the conclusion that every closed ideal in  $C_0(\Omega)$  is of the form  $C_0(\Lambda)$ , where  $\Lambda \subseteq \Omega$  is open.

## C\*-norms

We are interested in turning  $*$ -algebras into Banach  $*$ -algebras or  $C^*$ -algebras. To do this, we can actually use the Banach space completion,  $\overline{\iota(A)}^{\|\cdot\|_{\text{op}}} \subseteq A^{**}$ , where  $\iota$  is the canonical injection.

**Lemma:** If  $A_0$  is a normed  $*$ -algebra, then its Banach space completion is a Banach  $*$ -algebra, and the inclusion  $A_0 \hookrightarrow A$  is an injective  $*$ -homomorphism.

*Proof.* We know that  $A$  is a Banach space, and the inclusion  $A_0 \hookrightarrow A$  is an isometry. We show that  $A$  has an algebra structure that extends  $A_0$ , and the norm on  $A$  is submultiplicative.

Let  $x, y \in A$ , and let  $(x_n)_n, (y_n)_n$  be sequences in  $A_0$  converging to  $x$  and  $y$  respectively. Then,  $\sup_n \|x_n\| = C_1 < \infty$  and  $\sup_n \|y_n\| = C_2 < \infty$ , since convergent sequences are bounded. For  $m, n \in \mathbb{N}$ , we have

$$\begin{aligned} \|x_n y_n - x_m y_m\| &= \|x_n y_n - x_n y_m + x_n y_m - x_m y_m\| \\ &\leq \|x_n(y_n - y_m)\| + \|(x_n - x_m)y_m\| \\ &\leq C_1 \|y_n - y_m\| + C_2 \|x_n - x_m\|, \end{aligned}$$

meaning  $(x_n y_n)_n$  is Cauchy in  $A$ , and converges to  $x \cdot y = \lim_{n \rightarrow \infty} x_n y_n$ .

The map  $(x, y) \mapsto x \cdot y$  extends the multiplication on  $A_0$ , and endows  $A$  with the structure of an algebra.

For  $x, y \in A$ , and  $(x_n)_n, (y_n)_n$  sequences in  $A_0$  converging to  $x$  and  $y$  respectively, we get

$$\begin{aligned} \|xy\| &= \left\| \lim_{n \rightarrow \infty} x_n y_n \right\| \\ &= \lim_{n \rightarrow \infty} \|x_n y_n\| \\ &\leq \lim_{n \rightarrow \infty} \|x_n\| \|y_n\| \\ &= \|x\| \|y\|. \end{aligned}$$

Thus,  $A$  is a Banach algebra.

To see that  $A$  is a Banach  $*$ -algebra, we show that  $A$  admits the involution defined by, for  $x \in A$  and  $(x_n)_n \subseteq A_0$  with  $(x_n)_n \rightarrow x$ ,

$$x^* = \lim_{n \rightarrow \infty} x_n^*.$$

Similarly, we find that

$$\begin{aligned} \|x^*\| &= \lim_{n \rightarrow \infty} \|x_n^*\| \\ &= \lim_{n \rightarrow \infty} \|x_n\| \\ &= \|x\|, \end{aligned}$$

so  $A$  is a Banach  $*$ -algebra. □

**Definition.** Let  $A_0$  be a  $*$ -algebra. A  $C^*$ -norm on  $A_0$  is a norm satisfying

- (i)  $\|ab\| \leq \|a\| \|b\|$ ;
- (ii)  $\|a^*\| = \|a\|$ ;
- (iii)  $\|a^* a\| = \|a\|^2$

for all  $a, b \in A_0$ . We can define  $C^*$ -seminorms analogously.

On any given  $*$ -algebra, there can be many  $C^*$ -norms.

**Example.** Let  $\mathcal{T}$  be the unital  $*$ -algebra of trigonometric polynomials in  $C(\mathbb{T})$ . For every closed infinite set  $F \subseteq \mathbb{T}$ , we have a  $C^*$ -norm, given by

$$\|p\|_F = \sup_{z \in F} |p(z)|.$$

This is pretty clearly a  $C^*$ -seminorm, but it isn't clear at first sight that this is a norm. We can show this as follows.

Suppose  $\|p\|_F = 0$ , meaning  $p(z) = 0$  for all  $z \in F$ . Write

$$\begin{aligned} p(z) &= \sum_{k=-n}^n c_k z^k \\ q(z) &= z^n p(z). \end{aligned}$$

Then,  $q(z)$  is a polynomial, that vanishes on  $F$ . However, since  $q$  is a polynomial with degree  $2n$ ,  $q$  can have at most  $2n$  distinct roots by the fundamental theorem of algebra. Thus,  $q = 0$ , so  $p = 0$ .

We can generate  $C^*$ -norms and seminorms via morphisms into  $C^*$ -algebras.

**Lemma:** Let  $A_0$  be a  $*$ -algebra, and let  $\phi: A_0 \rightarrow B$  be a  $*$ -homomorphism into a  $C^*$ -algebra  $B$ . Then,

$$\|a\|_\phi = \|\phi(a)\|$$

defines a  $C^*$ -seminorm on  $A_0$ . If  $\phi$  is injective, then  $\|\cdot\|_\phi$  is a norm.

*Proof.* We will prove that this is a  $C^*$ -(semi)norm.

$$\begin{aligned} \|ab\|_\phi &= \|\phi(ab)\| \\ &= \|\phi(a)\phi(b)\| \\ &\leq \|\phi(a)\| \|\phi(b)\| \\ &= \|a\|_\phi \|b\|_\phi \end{aligned}$$

$$\begin{aligned} \|a^*\|_\phi &= \|\phi(a^*)\| \\ &= \|\phi(a)^*\| \\ &= \|\phi(a)\| \\ &= \|a\|_\phi \end{aligned}$$

$$\begin{aligned} \|a^*a\|_\phi &= \|\phi(a^*a)\| \\ &= \|\phi(a)^*\phi(a)\| \\ &= \|\phi(a)\|^2 \\ &= \|a\|_\phi^2. \end{aligned}$$

□

We can pass from seminorms to norms by modding out by the null set.

**Lemma:** Let  $p$  be a  $C^*$ -seminorm on the  $*$ -algebra  $A_0$ . The set

$$N_p = \{x \in A \mid p(x) = 0\}$$

is a  $*$ -ideal, and the map

$$\|a + N_p\|_{A/N_p} = p(a)$$

is a well-defined  $C^*$ -norm on  $A_0/N_p$ .

Now that we have defined a  $C^*$ -norm, we can extend this norm to the norm completion of the  $*$ -algebra  $A_0$ .

**Lemma:** Let  $\|\cdot\|$  be a  $C^*$ -norm on a  $*$ -algebra  $A_0$ . The norm completion  $A$  is a  $C^*$ -algebra, and the inclusion  $A_0 \hookrightarrow A$  is an isometric  $*$ -homomorphism.

*Proof.* We know that  $A$  is a Banach  $*$ -algebra, and the inclusion is an isometric  $*$ -homomorphism. We only need to check that the  $C^*$  property holds in  $A$ . Let  $x \in A$ ,  $(x_n)_n \rightarrow x$  in  $A_0$ . Then,

$$\begin{aligned} \|x^*x\| &= \lim_{n \rightarrow \infty} \|x_n^*x_n\| \\ &= \lim_{n \rightarrow \infty} \|x_n\|^2 \\ &= \|x\|^2. \end{aligned}$$

□

**Definition.** Let  $A_0$  be a  $*$ -algebra equipped with  $C^*$ -seminorm  $p$ . The norm completion of the  $*$ -algebra  $A_0/N_p$  with respect to  $\|\cdot\|_{A_0/N_p}$  is called the Hausdorff completion, or enveloping  $C^*$ -algebra, of the pair  $(A_0, p)$ .



## Universal C\*-Algebras

We are now interested in a sort of maximal Hausdorff completion of  $A_0$ .

**Definition.** Let  $A_0$  be a  $*$ -algebra, and let  $\mathcal{P}$  be the collection of all C\*-seminorms on  $A_0$ . For each  $a \in A_0$ , we set

$$\|a\|_u = \sup_{p \in \mathcal{P}} p(a).$$

If  $\|a\|_u < \infty$  for all  $a \in A_0$ , then  $\|\cdot\|_u$  defines a C\*-seminorm on  $A_0$ , called the universal C\*-seminorm. In this case, the universal enveloping C\*-algebra of  $A_0$  is the enveloping algebra of  $(A_0, \|\cdot\|_u)$ .

Recall that given a set of generators  $E = \{x_i\}_{i \in I}$  and relations  $R \subseteq \mathbb{A}^*(E)$ , we can construct the quotient  $*$ -algebra  $\mathbb{A}^*(E|R) = \mathbb{A}(E)/I(R)$ , where  $I(R)$  is the  $*$ -ideal generated by  $R$  contained in the free  $*$ -algebra on  $E$ . We write  $z_i = x_i + I(R)$ .

We also saw that  $\mathbb{A}^*(E|R)$  admits a universal property, wherein if  $B$  is any  $*$ -algebra admitting elements  $\{b_i\}_{i \in I}$  that satisfy  $R$ , then there is a  $*$ -homomorphism  $\phi_B: \mathbb{A}^*(E|R) \rightarrow B$ , defined by  $\phi_B(z_i) = b_i$ .

We can define a universal C\*-algebra by looking at the universal enveloping algebra of  $\mathbb{A}^*(E|R)$ , provided it exists.

**Definition.** Let  $E$  be a set of abstract symbols, and  $R \subseteq \mathbb{A}^*(E)$  is a set of relations. If the universal C\*-algebra of  $\mathbb{A}(E|R)$  exists — i.e., if  $\|a\|_u < \infty$  for all  $a \in \mathbb{A}^*(E|R)$  — then we write  $C^*(E|R)$  to denote this C\*-algebra, and call it the universal C\*-algebra generated by  $E$  with relations  $R$ .

Just as in the case of the universal  $*$ -algebra, we see that the universal C\*-algebra admits an analogous universal property.

**Proposition:** Let  $E = \{x_i\}_{i \in I}$  be a set of abstract symbols, and let  $R \subseteq \mathbb{A}^*(E)$  be a collection of relations. Let  $C^*(E|R)$  exist. If  $B$  is a C\*-algebra admitting elements  $\{b_i\}_{i \in I}$  that satisfy the relations, then there is a unique contractive  $*$ -homomorphism  $\phi_B: C^*(E|R) \rightarrow B$ , defined by  $\phi_B(v_i) = b_i$ , where  $v_i = (x_i + I(R)) + N_u$ .

*Proof.* By the universal property of  $\mathbb{A}^*(E|R)$ , we have  $\phi_B: \mathbb{A}^*(E|R) \rightarrow B$ , defined by  $\phi_B(z_i) = b_i$ , where  $z_i = x_i + I(R)$ .

We have the C\*-seminorm given by  $a \mapsto \|\phi_B(a)\|$ , where  $\|\phi_B(a)\| \leq \|a\|_u$  for all  $a \in \mathbb{A}^*(E|R)$ . Additionally, we must have that  $\phi_B$  kills the  $*$ -ideal

$$N_u = \{a \in \mathbb{A}^*(E|R) \mid \|a\|_u = 0\}.$$

By the first isomorphism theorem, we get the  $*$ -homomorphism  $\widetilde{\phi_B}: \mathbb{A}^*(E|R)/N_u \rightarrow B$ , given by  $z_i + N_u \mapsto b_i$ . This map is still contractive, so we can continuously extend  $\widetilde{\phi_B}$  to the desired contractive  $*$ -homomorphism,  $\phi_B: C^*(E|R) \rightarrow B$ , mapping  $z_i + N_u \mapsto b_i$ .

Uniqueness follows from the fact that  $\mathbb{A}^*(E|R)/N_u$  is dense in its completion.  $\square$

**Example.** It is sometimes the case that  $C^*(E|R)$  doesn't exist. Consider  $E = \{x\}$  and  $R = \{x - x^*\}$ . We write  $z = x + I(R)$ . For a  $t > 0$ , we find a C\*-algebra  $B_t$  and a self-adjoint  $b_t \in B_t$  with  $\|b_t\| = t$ .

For each  $t > 0$ , the universal property for  $\mathbb{A}^*(E|R)$  gives a  $*$ -homomorphism  $\phi_t: \mathbb{A}^*(E|R) \rightarrow B_t$ , with  $\phi_t(z) = b_t$ . We get a C\*-seminorm  $p_t$  on  $\mathbb{A}^*(E|R)$  given by  $p_t(a) = \|\phi_t(a)\| = t$ , meaning that the universal C\*-seminorm is

$$\|z\|_u \geq \sup_{t>0} p_t(z)$$

$$\begin{aligned}
&= \sup_{t>0} \|\phi_t(z)\| \\
&= \sup_{t>0} \|b_t\| \\
&= \sup_{t>0} t \\
&= \infty.
\end{aligned}$$

To verify that the universal  $C^*$ -seminorm is finite for every element in  $\mathbb{A}^*(E|R)$ , we can use a simpler characterization.

**Lemma:** Let  $E = \{x_i\}_{i \in I}$  be a set of symbols and suppose  $R \subseteq \mathbb{A}^*(E|R)$  is a collection of relations. Write  $z_i = x_i + I(R)$ . If there is a  $C \geq 0$  for which  $p(z_i) \leq C$  for every  $i \in I$  and every  $C^*$ -seminorm  $p$  on  $\mathbb{A}^*(E|R)$ , then  $C^*(E|R)$  exists.

We can consider the  $C^*$ -algebra of  $n \times n$  matrices over  $\mathbb{C}$ , and construct this  $C^*$ -algebra using the universal  $C^*$ -algebra.

**Example.** Let  $n \geq 1$ , and let  $E_n = \{x_{ij} \mid 1 \leq i, j \leq n\}$ . Let

$$R = \left\{ x_{ij}^* - x_{ji}, x_{ij}x_{kl} - \delta_{jk}x_{il} \mid i, j \in \{1, \dots, n\} \right\}$$

be our set of relations.<sup>iv</sup> Let  $z_{ij} = x_{ij} + I(R)$ . Then, if  $p$  is any  $C^*$ -seminorm on  $\mathbb{A}^*(E_n|R)$ , we have

$$\begin{aligned}
p(z_{jj})^2 &= p(z_{jj}^* z_{jj}) \\
&= p(z_{jj} z_{jj}) \\
&= p(z_{jj}),
\end{aligned}$$

so  $p(z_{jj}) \subseteq \{0, 1\}$ , and we also have

$$\begin{aligned}
p(z_{ij})^2 &= p(z_{ij}^* z_{ij}) \\
&= p(z_{ji} z_{ij}) \\
&= p(z_{jj}) \\
&\in \{0, 1\}.
\end{aligned}$$

Thus,  $C^*(E_n|R)$  exists. Write  $v_{ij} = z_{ij} + N_u$ . We will show that  $C^*(E_n|R)$  is not trivial.

The matrix units  $\{e_{ij} \mid 1 \leq i, j \leq n\}$  satisfy the relations, so by the universal property of  $C^*(E_n|R)$ , we have a contractive  $*$ -homomorphism  $\varphi: C^*(E_n|R) \rightarrow \text{Mat}_n(\mathbb{C})$  given by  $\varphi(v_{ij}) = e_{ij}$ . Since  $\text{span}(\{e_{ij}\}_{i,j}) = \text{Mat}_n(\mathbb{C})$ , we must have  $C^*(E_n|R) \cong \text{Mat}_n(\mathbb{C})$ .

Consequently,  $C^*(E_n|R)$  is simple. Additionally, if  $B$  is any other  $C^*$ -algebra admitting elements  $\{b_{ij} \mid 1 \leq i, j \leq n\}$  with  $b_{ij}^* = b_{ji}$  and  $b_{ij}b_{kl} = \delta_{jk}b_{il}$ , then there is a unique injective  $*$ -homomorphism between  $\text{Mat}_n(\mathbb{C})$  and  $B$  such that  $\varphi(e_{ij}) \cong b_{ij}$ .

We can also obtain  $\text{Mat}_n(\mathbb{C})$  another way. Consider  $F_n = \{x_1, \dots, x_n\}$  and the relations

$$R' = \{x_i^* x_j - \delta_{ij} x_1 \mid i, j = 1, \dots, n\}.$$

We write  $z_i = x_i + I(R')$ . If  $p$  is any  $C^*$ -seminorm on  $\mathbb{A}^*(F_n|R')$ , then

$$p(z_i)^2 = p(z_i^* z_i)$$

<sup>iv</sup>The first set of relations denotes the (conjugate) transpose,  $x_{ij}^* = x_{ji}$  and the second set of relations denotes  $x_{ij}x_{kl} = \delta_{jk}x_{il}$ , which is the index notation definition of matrix multiplication.

$$= p(z_1),$$

so  $p(z_i) \in \{0, 1\}$  for all  $i$ . Thus,  $C^*(F_n|R')$  exists.

Write  $v_i = z_i + N_u$ , and set

$$b_{ij} = v_i v_j^*.$$

We have  $b_{ij}^* = b_{ji}$ , and since  $v_i^* v_i = v_1$  is a projection for every  $i$ , each  $v_i$  is a partial isometry, meaning

$$\begin{aligned} b_{ij} b_{kl} &= v_i (v_j^* v_k) v_l^* \\ &= v_i (\delta_{jk} v_1) v_l^* \\ &= v_i (\delta_{jk} v_i^* v_i) v_l^* \\ &= \delta_{jk} (v_i v_i^* v_i) v_l^* \\ &= \delta_{jk} v_i v_l^*. \end{aligned}$$

Thus, there is a  $*$ -homomorphism between  $\text{Mat}_n(\mathbb{C})$  and  $C^*(F_n|R')$ , given by  $\psi(e_{ij}) = b_{ij}$ . Since  $\text{Mat}_n(\mathbb{C})$  is simple,  $\psi$  is injective.

We also have

$$\begin{aligned} \psi(e_{i1}) &= b_{i1} \\ &= v_i v_1^* \\ &= v_i v_1 \\ &= v_i v_i^* v_i \\ &= v_i, \end{aligned}$$

so  $\psi$  is onto. Thus  $C^*(F_n|R') \cong \text{Mat}_n(\mathbb{C})$ .

**Example.** Let  $E = \{1, x\}$  and

$$R = \{x^*x - 1, xx^* - 1, 1x - x, x1 - x, 1^2 - 1, 1^* - 1\}.$$

We see that  $\mathbb{A}^*(E|R)$  is unital with unit  $1 + I(R)$ , and that  $x + I(R)$  is invertible with inverse  $x^* + I(R)$ . Writing  $z = x + I(R)$ , we see that

$$\mathbb{A}^*(E|R) = \left\{ \sum_{k \in \mathbb{Z}} \alpha_k z^k \mid \alpha_k \in \mathbb{C}, \text{ finitely many nonzero} \right\},$$

where  $z^{-1} = z^*$  and  $z^0 = 1$ .

If  $p$  is any seminorm on  $\mathbb{A}^*(E|R)$ , we have

$$\begin{aligned} p(1)^2 &= p(1^*1) \\ &= p(1^2) \\ &= p(1), \end{aligned}$$

so  $p(1) \in \{0, 1\}$ , and

$$\begin{aligned} p(z)^2 &= p(z^*z) \\ &= p(1) \\ &\in \{0, 1\}. \end{aligned}$$

Thus,  $C^*(E|R)$  exists. We write  $u = z + N_u$ . The universal property states that if  $w$  is a unitary in any unital  $C^*$ -algebra  $B$ , then there is a surjective  $*$ -homomorphism between  $C^*(E|R)$  and  $C^*(w) \subseteq B$ , given by  $u \mapsto w$ .

Eventually, we will show that  $C^*(E|R) \cong C(\mathbb{T})$ .

## Representations and the Group C\*-algebra

We can realize  $*$ -algebras as  $*$ -subalgebras of bounded operators on a Hilbert space. This allows us to get a  $C^*$ -norm for free, and get a  $C^*$ -algebra by completion.

**Definition.** Let  $A_0$  be a  $*$ -algebra. A representation of  $A_0$  is a pair  $(\pi_0, \mathcal{H})$ , where  $\mathcal{H}$  is a Hilbert space and  $\pi_0: A_0 \rightarrow \mathcal{B}(\mathcal{H})$  is a  $*$ -homomorphism. We will refer to the representation by  $\pi_0$  if the Hilbert space is understood.

If  $A_0$  is unital, and  $\pi(1_A) = I_{\mathcal{H}}$ , then we say  $\pi$  is a unital representation.

**Lemma:** Let  $A_0$  be a  $*$ -algebra, and suppose  $(\pi_0, \mathcal{H})$  is a representation of  $A_0$ . Then,

$$\|a\|_{\pi_0} = \|\pi_0(a)\|_{\text{op}}$$

is a  $C^*$ -seminorm on  $A_0$ . If  $\pi_0$  is injective, then  $\|\cdot\|_{\pi_0}$  is a  $C^*$ -norm.

**Lemma:** Let  $A_0$  and  $B_0$  be normed  $*$ -algebras with respective completions  $A$  and  $B$ . If  $\varphi_0: A_0 \rightarrow B_0$  is a bounded  $*$ -homomorphism, then the continuous extension  $\varphi: A \rightarrow B$  is a  $*$ -homomorphism.

*Proof.* Let  $x, y \in A$  with  $(x_n)_n \rightarrow x$  and  $(y_n)_n \rightarrow y$  sequences in  $A_0$ . Then,

$$\begin{aligned} \varphi(xy) &= \varphi\left(\lim_{n \rightarrow \infty} x_n y_n\right) \\ &= \lim_{n \rightarrow \infty} \varphi(x_n y_n) \\ &= \lim_{n \rightarrow \infty} \varphi_0(x_n y_n) \\ &= \lim_{n \rightarrow \infty} \varphi_0(x_n) \varphi_0(y_n) \\ &= \lim_{n \rightarrow \infty} \varphi(x_n) \varphi(y_n) \\ &= \varphi(x) \varphi(y). \end{aligned}$$

A similar process, using the continuity of the involution, gives  $\varphi(x^*) = \varphi(x)^*$ . □

**Corollary:** Let  $A_0$  be a  $*$ -algebra, and suppose  $\pi: A_0 \rightarrow \mathcal{B}(\mathcal{H})$  is an injective representation. The completion  $A$  of  $A_0$  with respect to the  $C^*$ -norm  $\|\cdot\|_{\pi_0}$  is a  $C^*$ -algebra, and the continuous extension  $\pi: A \rightarrow \mathcal{B}(\mathcal{H})$  is an isometric  $*$ -homomorphism.

The  $C^*$ -algebra that arises from a group is an important example of a  $C^*$ -algebra.<sup>v</sup>

Given a group  $\Gamma$ , we can construct the group  $*$ -algebra,  $\mathbb{C}[\Gamma]$ . An element  $a \in \mathbb{C}[\Gamma]$  is a finitely supported complex-valued function on  $\Gamma$ , written as a finite sum

$$a = \sum_{s \in \Gamma} a(s) \delta_s,$$

where  $\delta_s: \Gamma \rightarrow \mathbb{C}$  is the indicator function for  $s$ ,  $\delta_s(t) = \delta_{st}$ .

Unitary representations of  $\Gamma$  are related to representations of the group  $*$ -algebra  $\mathbb{C}[\Gamma]$ .

**Proposition:** Let  $\Gamma$  be a group, and let  $\mathcal{H}$  be a Hilbert space.

(1) If  $u: \Gamma \rightarrow \mathcal{U}(\mathcal{H})$  is a unitary representation of  $\Gamma$ , then the map  $\pi_u: \mathbb{C}[\Gamma] \rightarrow \mathcal{B}(\mathcal{H})$  given by

$$\pi_u(a) = \sum_{s \in \Gamma} a(s) u_s$$

is a representation of  $\mathbb{C}[\Gamma]$ .

---

<sup>v</sup>It's partially the subject of my Honors thesis.

(2) If  $\pi: \mathbb{C}[\Gamma] \rightarrow \mathbb{B}(\mathcal{H})$  is a unital representation, then the map  $u: \Gamma \rightarrow \mathcal{U}(\mathcal{H})$ , given by

$$u(s) = \pi(\delta_s)$$

is a unitary representation of  $\Gamma$ .

*Proof.*

(1) The map  $s \mapsto u_s \in \mathbb{B}(\mathcal{H})$  extends to a linear map  $\pi_u: \mathbb{C}[\Gamma] \rightarrow \mathbb{B}(\mathcal{H})$ , satisfying  $\pi_u(\delta_s) = u_s$  by the universal property of the free vector space.

For  $s, t \in \Gamma$ , we have

$$\begin{aligned} \pi_u(\delta_s \delta_t) &= \pi_u(\delta_{st}) \\ &= u_{st} \\ &= u_s u_t \\ &= \pi_u(\delta_s) \pi_u(\delta_t) \end{aligned}$$

$$\begin{aligned} \pi_u(\delta_s^*) &= \pi_u(\delta_s^{-1}) \\ &= u_{s^{-1}} \\ &= u_s^* \\ &= \pi_u(\delta_s)^*. \end{aligned}$$

Using the linearity of  $\pi_u$ , we see that  $\pi_u$  is multiplicative and  $*$ -preserving.

(2) Every  $\delta_s \in \mathbb{C}[\Gamma]$  is unitary, and since unital  $*$ -homomorphisms map unitaries to unitaries, we know that each  $u(s)$  is unitary. Moreover, for  $s, t \in \Gamma$ , we have

$$\begin{aligned} u(st) &= \pi(\delta_{st}) \\ &= \pi(\delta_s \delta_t) \\ &= \pi(\delta_s) \pi(\delta_t) \\ &= u(s) u(t), \end{aligned}$$

meaning  $u$  is a unitary representation. □

For the group  $\Gamma$  is a group with neutral element  $e$ , we have defined the group  $*$ -algebra and the left-regular representation  $\lambda: \Gamma \rightarrow \mathcal{U}(\ell_2(\Gamma))$ . We thus get a representation of the group  $*$ -algebra

$$\pi_\lambda(a) = \sum_{s \in \Gamma} a(s) \lambda_s.$$

We claim that  $\pi_\lambda$  is injective. Suppose  $\pi_\lambda(a) = 0$  for some  $a = \sum_{s \in \Gamma} a(s) \delta_s \in \mathbb{C}[\Gamma]$ . Evaluating  $\delta_e$ , we have

$$\begin{aligned} 0 &= \pi_\lambda(a)(\delta_e) \\ &= \left( \sum_{s \in \Gamma} a(s) \lambda_s \right) (\delta_e) \\ &= \sum_{s \in \Gamma} a(s) \lambda_s(\delta_e) \\ &= \sum_{s \in \Gamma} (\delta_s). \end{aligned}$$

Since the vectors  $\{\delta_t\}_{t \in \Gamma}$  are linearly independent, we must have  $a(s) = 0$  for all  $s \in \Gamma$ , so  $a = 0$ .

Thus, we have a  $C^*$ -norm on  $\mathbb{C}[\Gamma]$  given by  $\|a\|_\lambda = \|\pi_\lambda(a)\|_{\text{op}}$ . The  $\|\cdot\|_\lambda$ -completion of  $\mathbb{C}[\Gamma]$  is a  $C^*$ -algebra denoted by  $C_\lambda^*(\Gamma)$ . This is known as the left-regular group  $C^*$ -algebra.

Similarly, we may begin with the right-regular representation  $\rho: \mathbb{C}[\Gamma] \rightarrow \mathcal{U}(\ell_2(\Gamma))$ , and construct the representation

$$\pi_\rho(a) = \sum_{s \in \Gamma} a(s) \rho_s,$$

which induces the  $C^*$ -norm  $\|\cdot\|_\rho$  on  $\mathbb{C}[\Gamma]$ , which gives rise to the right-regular group  $C^*$ -algebra,  $C_\rho^*(\Gamma)$ .

We often refer to  $C_\lambda^*(\Gamma)$  as the reduced group  $C^*$ -algebra of  $\Gamma$ , often denoted  $C_r^*(\Gamma)$ .

There is also a full group  $C^*$ -algebra, with the full norm defined by

$$\|a\|_u = \sup \{ \|\pi(a)\| \mid \pi: \mathbb{C}[\Gamma] \rightarrow \mathbb{B}(\mathcal{H}_\pi) \text{ is a representation} \}.$$

To see that this quantity is finite, note that for every representation  $\pi: \mathbb{C}[\Gamma] \rightarrow \mathbb{B}(\mathcal{H}_\pi)$ , the elements  $\pi(\delta_s)$  are unitaries in  $\mathbb{B}(\mathcal{H})$ , hence having norm 1. So, we have

$$\begin{aligned} \|\pi(a)\| &= \left\| \pi \left( \sum_{s \in \Gamma} a(s) \delta_s \right) \right\| \\ &= \left\| \sum_{s \in \Gamma} a(s) \pi(\delta_s) \right\| \\ &\leq \sum_{s \in \Gamma} \|a(s) \delta_s\| \\ &= \sum_{s \in \Gamma} |a(s)|, \end{aligned}$$

so  $\|a\|_u \leq \sum_{s \in \Gamma} |a(s)| < \infty$ . This is a  $C^*$ -norm, as if  $\|a\|_u = 0$ , then  $\|a\|_\lambda = 0$ , as  $\pi_\lambda$  is one of the representations, and since  $\|\cdot\|_\lambda$  is a norm, we must have  $a = 0$ . Thus, completing  $\mathbb{C}[\Gamma]$  with respect to  $\|\cdot\|_u$  yields the full (or universal) group  $C^*$ -algebra, denoted  $C^*(\Gamma)$ .

The full group  $C^*$ -algebra admits a universal property.

**Proposition:** Let  $\Gamma$  be a discrete group. Given any unitary representation  $u: \Gamma \rightarrow \mathcal{U}(\mathcal{H})$ , there is a contractive  $*$ -homomorphism  $\pi_u: C^*(\Gamma) \rightarrow \mathbb{B}(\mathcal{H})$  satisfying  $\pi_u(\delta_s) = u(s)$  for every  $s \in \Gamma$ .

*Proof.* We have a representation  $\pi_u: \mathbb{C}[\Gamma] \rightarrow \mathbb{B}(\mathcal{H})$  that extends  $u: \Gamma \rightarrow \mathcal{U}(\mathcal{H})$ . By definition, the universal norm provides  $\|\pi(a)\|_u \leq \|a\|_u$ .

The continuous extension  $\pi_u: C^*(\Gamma) \rightarrow \mathbb{B}(\mathcal{H})$  is contractive and a  $*$ -homomorphism.  $\square$

### Unitizations of $C^*$ -Algebras

Given a non-unital algebra  $A$ , there is a unital algebra,  $\tilde{A}$ , that contains  $A$  as a maximal and essential ideal. We will now examine the analytical component of unitization — given a Banach algebra or  $C^*$ -algebra, we want the resulting unitization to also be a Banach algebra or  $C^*$ -algebra.

**Proposition:** Let  $A$  be a Banach  $*$ -algebra. The unitization  $\tilde{A}$  is a unital Banach  $*$ -algebra with the norm

$$\|(a, \alpha)\| = \|a\| + |\alpha|.$$

The inclusion  $\iota_A: A \rightarrow \tilde{A}$ , given by  $\iota(a) = (a, 0)$ , is an isometric  $*$ -isomorphism.

*Proof.* Let  $A$  be a Banach  $*$ -algebra. We know that the unitization,  $\tilde{A}$ , is a unital  $*$ -algebra, and  $\iota_A$  is a  $*$ -homomorphism.

We can see that  $\|\cdot\|$  is a norm on the vector space  $\tilde{A}$  from its definition. To verify that it is a norm on the algebra, we have

$$\begin{aligned} \|(a, \alpha)(b, \beta)\| &= \|(ab + \alpha b + \beta a, \alpha\beta)\| \\ &= \|ab + \alpha b + \beta a\| + |\alpha\beta| \\ &\leq \|a\|\|b\| + |\alpha|\|b\| + |\beta|\|a\| + |\alpha||\beta| \\ &= (\|a\| + |\alpha|)(\|b\| + |\beta|) \\ &= \|(a, \alpha)\| \|(b, \beta)\|. \end{aligned}$$

We also have

$$\begin{aligned} \|(a, \alpha)^*\| &= \|(a^*, \bar{\alpha})\| \\ &= \|a^*\| + |\bar{\alpha}| \\ &= \|a\| + |\alpha| \\ &= \|(a, \alpha)\|. \end{aligned}$$

To see that the norm on  $\tilde{A}$  is complete, recall that the projection  $\pi: \tilde{A} \rightarrow \mathbb{C}$ , given by  $(a, \alpha) \mapsto \alpha$ , is a 1-quotient mapping, so  $\tilde{A}/A$  is isometrically isomorphic to  $\mathbb{C}$ , hence complete. Since  $A$  is also complete, we must have  $\tilde{A}$  is complete, as it is a two of three spaces property.  $\square$

Turning our attention to  $C^*$ -algebras, we know that the traditional unitization converts  $A$  into a Banach  $*$ -algebra. However, this norm is not a  $C^*$ -norm. Instead, we embed  $A$  isometrically into an algebra of bounded operators in order to obtain the unitization.

If  $A$  is an algebra, we let  $L_a(x) = ax$  be left-multiplication by  $a$ . If  $A$  is normed, we can see that  $L_a(x)$  is continuous:

$$\begin{aligned} \|L_a(x)\| &= \|ax\| \\ &\leq \|a\|\|x\|. \end{aligned}$$

Thus, we have a map  $L: A \rightarrow \mathcal{B}(A)$  given by  $a \mapsto L_a$ . We can also see that  $L_{a+\alpha b} = L_a + \alpha L_b$ , and  $L_{ab} = L_a \circ L_b$ , so  $L$  is an algebra homomorphism. We may extend to the unitization, so we obtain the unital algebra homomorphism

$$\bar{L}(a, \alpha) = L_a + \alpha \text{id}_A.$$

We know that if  $A$  is nonunital and  $L$  is injective, then  $\bar{L}$  is injective. This will allow us to unitize a nonunital  $C^*$ -algebra.

**Lemma:** Let  $A$  be a normed algebra, and let  $L: A \rightarrow \mathcal{B}(A)$  and  $\bar{L}: \tilde{A} \rightarrow \mathcal{B}(A)$  be as above.

- (1)  $L$  is a contractive algebra homomorphism, and  $\text{Ran}(L) \subseteq \mathcal{B}(A)$  is a subalgebra.
- (2) If  $A$  is a  $C^*$ -algebra, then  $L$  is isometric, and  $\text{Ran}(L) \subseteq \mathcal{B}(A)$  is closed in operator norm.
- (3) If  $A$  is a nonunital  $C^*$ -algebra, then  $\bar{L}$  is an injective algebra homomorphism, restricting to an isometry on  $A$ , and  $\text{Ran}(\bar{L}) \subseteq \mathcal{B}(A)$  is closed in operator norm.

*Proof.* We have proven (1) already, so we prove (2) and (3).

(2) If  $A$  is a  $C^*$ -algebra, then we see that

$$\begin{aligned}\|L_a\|_{\text{op}} &\geq \left\| L_a \left( \frac{a^*}{\|a\|} \right) \right\| \\ &= \frac{\|aa^*\|}{\|a\|} \\ &= \frac{\|a\|^2}{\|a\|} \\ &= \|a\|.\end{aligned}$$

Thus,  $\|L_a\|_{\text{op}} = \|a\|$ , so  $L$  is isometric. Since  $L$  is complete, and  $L$  is an isometry,  $\text{Ran}(L)$  is complete, so it is closed in  $\mathbb{B}(A)$ .

We have seen that  $L$  is isometric, hence injective. Since  $A$  is nonunital,  $\bar{L}$  is injective too. Since  $\text{Ran}(L)$  is closed, the sum  $\text{Ran}(L) + \mathbb{C} \text{id}_A$  is closed as well.

□

**Proposition:** Let  $A$  be a  $C^*$ -algebra, and let  $L: A \rightarrow \mathbb{B}(A)$ ,  $\bar{L}: A \rightarrow \mathbb{B}(A)$  be as above.

(1) The quantity

$$\|(a, \alpha)\|_L = \|L_a + \alpha \text{id}_A\|_{\text{op}}$$

is a  $C^*$ -seminorm on  $\tilde{A}$ .

(2) If  $A$  is nonunital, then  $\|\cdot\|_L$  is a  $C^*$ -norm on  $\tilde{A}$ , and  $(\tilde{A}, \|\cdot\|_L)$  is a unital  $C^*$ -algebra. The inclusion  $\iota_A: A \hookrightarrow (\tilde{A}, \|\cdot\|_L)$  is an isometric  $*$ -homomorphism.

(3) The quantity

$$\|(a, \alpha)\|_1 = \max(\|(a, \alpha)\|_L, |\alpha|)$$

is a  $C^*$ -norm on  $\tilde{A}$ .

(4)  $(\tilde{A}, \|\cdot\|_1)$  is a unital  $C^*$ -algebra, and the inclusion,  $\iota_A: A \hookrightarrow (\tilde{A}, \|\cdot\|_1)$  is an isometric  $*$ -homomorphism.

*Proof.*

(1) Since  $\bar{L}$  is an algebra homomorphism, we know that  $\|\cdot\|_L$  is a seminorm. We will show the rest of the definitions simultaneously:

$$\begin{aligned}\|(a, \alpha)\|_L^2 &= \sup_{x \in B_A} \|ax + \alpha x\|^2 \\ &= \sup_{x \in B_A} \|(ax + \alpha x)^*(ax + \alpha x)\| \\ &= \sup_{x \in B_A} \|x^* a^* ax + \alpha x^* a^* x + \bar{\alpha} x^* ax + |\alpha|^2 x^* x\|\end{aligned}$$

Thus,  $\|(a, \alpha)\|_L \leq \|(a, \alpha)^*\|_L$ , and  $\|(a, \alpha)^*\|_L \leq \|(a, \alpha)\|_L$ , so  $\|(a, \alpha)^*\|_L = \|(a, \alpha)\|_L$ . This means all the inequalities above are indeed equalities, so we also recover the  $C^*$  identity.

(2) Since  $\bar{L}: A \rightarrow \mathbb{B}(A)$  is injective,  $\|\cdot\|_L$  is a norm.



Additionally, we know that  $\bar{L}: (\tilde{A}, \|\cdot\|_L) \rightarrow (\text{Ran}(\bar{L}), \|\cdot\|_{\text{op}})$  is an isometric isomorphism. Since  $(\text{Ran}(\bar{L}), \|\cdot\|_{\text{op}})$  is a Banach algebra, so too is  $(\tilde{A}, \|\cdot\|_L)$ , so  $(\tilde{A}, \|\cdot\|_L)$  is a  $C^*$ -algebra.

We can also see that  $\iota_A$  is isometric, since

$$\begin{aligned} \|\iota(a)\|_L &= \|(a, 0)\|_L \\ &= \|\bar{L}(a, 0)\|_{\text{op}} \\ &= \|L_a\|_{\text{op}} \\ &= \|a\|. \end{aligned}$$

- (3) That  $\|\cdot\|_1$  is a  $C^*$ -seminorm follows from (1), and  $a \mapsto |a|$  is a  $C^*$ -norm on  $\mathbb{C}$ . If  $\|(a, \alpha)\|_1 = 0$ , then  $\alpha = 0$ , so  $\|(a, 0)\|_L = 0$ , meaning  $\|L_a\|_{\text{op}} = 0$ , so  $a = 0$ .
- (4) Let  $((a_n, \alpha_n))_n$  be a  $\|\cdot\|_1$ -Cauchy sequence in  $\tilde{A}$ .

It follows that  $(\alpha_n)_n$  is Cauchy in  $\mathbb{C}$ , and  $(L_{a_n} + \alpha_n \text{id}_A)_n$  is Cauchy in  $\mathcal{B}(A)$ . Thus, there are  $\alpha \in \mathbb{C}$  and  $T \in \mathcal{B}(A)$  that these sequences respectively converge to.

We see that  $(\alpha_n \text{id}_A)_n \rightarrow \alpha \text{id}_A$ , so  $L_{a_n} \rightarrow T - \alpha \text{id}_A$ . Since  $\text{Ran}(L)$  is closed,  $T - \alpha \text{id}_A = L_a$  for some  $a \in A$ , meaning  $T = L_a + \alpha \text{id}_A$ . Thus,  $((a_n, \alpha_n))_n \xrightarrow{\|\cdot\|_1} (a, \alpha)$ , so  $\|\cdot\|_1$  is complete.

It is clear that  $\iota_A$  is isometric, as

$$\begin{aligned} \|\iota(a)\|_1 &= \|(a, 0)\|_1 \\ &= \|(a, 0)\|_L \\ &= \|a\|. \end{aligned}$$

□

**Definition.** Let  $A$  be a  $C^*$ -algebra.

- (1) If  $A$  is non-unital, then  $(\tilde{A}, \|\cdot\|_L)$  is known as the minimal  $C^*$ -unitization of  $A$ .
- (2) The  $C^*$ -algebra  $(\tilde{A}, \|\cdot\|_1)$  is known as the forced unitization of  $A$ , referred to as  $A^1$  or  $A^+$ .

**Proposition:** Let  $A$  be a  $C^*$ -algebra.

- (1) If  $A$  is nonunital, then  $\|\cdot\|_1$  and  $\|\cdot\|_L$  are equal.
- (2) If  $A$  is unital, then there is an isometric  $*$ -isomorphism between  $A \oplus \mathbb{C} \rightarrow A^1$ .

*Proof.* We will prove (2).

- (2) Consider the  $*$ -isomorphism  $A \oplus \mathbb{C} \rightarrow \tilde{A}$ , given by  $(a, z) = a + zp$ , where  $p = 1_{\tilde{A}} - 1_A$ . We only need to show that this is an isometry when  $A \oplus \mathbb{C}$  is equipped with the infinity norm and  $\tilde{A}$  is equipped with  $\|\cdot\|_1$ .

Note that

$$\bar{L}(a + zp) = \bar{L}((a - z1_A, z))$$

$$\begin{aligned}
&= L_{a-z1_A} + z \operatorname{id}_A \\
&= L_a - z \operatorname{id}_A + z \operatorname{id}_A \\
&= L_a.
\end{aligned}$$

Thus,

$$\begin{aligned}
\|a + zp\|_1 &= \max(\|L_a\|_{\operatorname{op}}, |z|) \\
&= \max(\|a\|, |z|) \\
&= \|(a, z)\|_{\infty}.
\end{aligned}$$

□

## Gelfand Theory

Diving deeper into the theory of  $C^*$ -algebras, we are interested in characterizing  $C^*$ -algebras, ultimately realizing commutative  $C^*$ -algebras as continuous function algebras, and non-commutative  $C^*$ -algebras as bounded operators on Hilbert spaces.

### Properties of the Spectrum

We can create an analytic characterization of invertibility in a Banach algebra.

**Proposition** (Carl Neumann Series): Let  $A$  be a unital Banach algebra. If  $x \in A$  with  $\|x\| < 1$ , then  $1_A - x \in \operatorname{GL}(A)$ , and

$$(1_A - x)^{-1} = \sum_{k=0}^{\infty} x^k.$$

Moreover,

$$\|(1_A - x)^{-1}\| \leq \frac{1}{1 - \|x\|}.$$

*Proof.* Since  $A$  is a normed algebra, we have  $\|x^k\| \leq \|x\|^k$ , so

$$\begin{aligned}
\sum_{k=0}^{\infty} \|x^k\| &\leq \sum_{k=0}^{\infty} \|x\|^k \\
&= \frac{1}{1 - \|x\|},
\end{aligned}$$

so the series  $\sum_{k=0}^{\infty} x^k$  converges absolutely, hence converges as  $A$  is complete. Note that this also means  $\lim_{n \rightarrow \infty} x^n = 0$ . We compute

$$\begin{aligned}
(1_A - x) \left( \sum_{k=0}^{\infty} x^k \right) &= \lim_{n \rightarrow \infty} (1_A - x) \left( \sum_{k=0}^n x^k \right) \\
&= \lim_{n \rightarrow \infty} \left( \sum_{k=0}^n x^k - \sum_{k=0}^n x^{k+1} \right) \\
&= \lim_{n \rightarrow \infty} (1_A - x^{n+1}) \\
&= 1_A.
\end{aligned}$$

Similarly,  $(\sum_{k=0}^{\infty} x^k)(1_A - x) = 1_A$ .

□

**Corollary:** Let  $A$  be a unital Banach algebra, and let  $a \in A$  with  $\|1_A - a\| < 1$ . Then,  $a \in GL(A)$ , and

$$a^{-1} = \sum_{k=0}^{\infty} (1_A - a)^k$$

$$\|a^{-1}\| \leq \frac{1}{1 - \|1_A - a\|}.$$

**Proposition:** Let  $A$  be a unital Banach algebra.

- (1) The group of invertible elements,  $GL(A) \subseteq A$ , is open.
- (2) The inverse map,  $i: GL(A) \rightarrow GL(A)$ , given by  $i(a) = a^{-1}$ , is a homeomorphism.

*Proof.*

- (1) Let  $a \in GL(A)$ , and set  $\delta = \|a^{-1}\|^{-1}$ . We will show that  $U(a, \delta) \subseteq GL(A)$ . Let  $b \in A$  be such that  $\|a - b\| < \delta$ . Then,

$$\begin{aligned} \|1_A - a^{-1}b\| &= \|a^{-1}(a - b)\| \\ &\leq \|a^{-1}\| \|a - b\| \\ &< 1. \end{aligned}$$

Thus,  $a^{-1}b \in GL(A)$  with

$$\|b^{-1}a\| \leq \frac{1}{1 - \|1 - a^{-1}b\|}.$$

Thus, we get  $b = a(a^{-1}b) \in GL(A)$ .

- (2) It suffices to show that  $i$  is continuous. Let  $a, b \in GL(A)$ . Notice that

$$\begin{aligned} \|b^{-1} - a^{-1}\| &= \|b^{-1}(1_A - ba^{-1})\| \\ &= \|b^{-1}(a - b)a^{-1}\| \\ &\leq \|b^{-1}\| \|a - b\| \|a^{-1}\|. \end{aligned}$$

We wish to control  $\|b^{-1}\|$ , so we find

$$\begin{aligned} \|b^{-1}\| &= \|b^{-1}aa^{-1}\| \\ &\leq \|b^{-1}a\| \|a^{-1}\| \\ &\leq \frac{\|a^{-1}\|}{1 - \|1_A - a^{-1}b\|}. \end{aligned}$$

Additionally, we have

$$\begin{aligned} \|1_A - a^{-1}b\| &= \|a^{-1}(a - b)\| \\ &\leq \|a^{-1}\| \|a - b\|, \end{aligned}$$

so

$$1 - \|1_A - a^{-1}b\| \geq 1 - \|a^{-1}\| \|a - b\|.$$

Combining, we get

$$\|b^{-1}\| \leq \frac{\|a^{-1}\|}{1 - \|a^{-1}\| \|a - b\|}.$$

Thus, we get

$$\|b^{-1} - a^{-1}\| \leq \frac{\|a^{-1}\|^2 \|a - b\|}{1 - \|a^{-1}\| \|a - b\|}.$$

For any sequence  $(b_n)_n$  in  $GL(A)$  that converges to  $a \in GL(A)$ , we see that  $\|i(b_n) - i(a)\| \rightarrow 0$ , so  $i$  is continuous.

□

We can now examine the analytic properties of the resolvent and spectrum.

**Theorem:** Let  $A$  be a Banach algebra with  $a \in A$ .

- (1) The resolvent  $\rho(a) \subseteq \mathbb{C}$  is open, and the spectrum  $\sigma(a) \subseteq \mathbb{C}$  is closed.
- (2) If  $\lambda \in \mathbb{C}$  with  $|\lambda| > \|a\|$ , then  $\lambda \in \rho(a)$ . Thus,  $\sigma(a) \subseteq \overline{D(0, \|a\|)}$ , meaning  $\sigma(a)$  is compact.
- (3) If  $A$  is unital, then  $R_a: \rho(a) \rightarrow A$ , given by  $R_a(a - \lambda 1_A)^{-1}$  is holomorphic on  $\rho(a)$ .
- (4) The spectrum  $\sigma(a)$  is nonempty. If  $A$  is nonunital,  $0 \in \sigma(a)$ .

*Proof.*

- (1) We assume  $A$  is unital. Let  $t: \mathbb{C} \rightarrow A$  be given by  $t(\lambda) = a - \lambda 1_A$ , which is continuous. Since  $t^{-1}(GL(A)) = \rho(a)$ , and  $GL(A)$  is open,  $\rho(a)$  is open. The complement,  $\sigma(a)$ , is closed.

If  $A$  is not unital, then  $\rho(a) = \rho(\iota_A(a))$  by definition, which is yet again open, so  $\sigma(a)$  is yet again closed.

- (2) We assume  $A$  is unital. If  $\lambda \in \mathbb{C}$  with  $|\lambda| \geq \|a\|$ , then  $\|\lambda^{-1}a\| = |\lambda|^{-1}\|a\| < 1$ , so  $1 - \lambda^{-1}a \in GL(A)$ , meaning  $a - \lambda 1_A = -\lambda(1_A - \lambda^{-1}a) \in GL(A)$ , so  $\lambda \in \rho(a)$ .

If  $A$  is nonunital, then the canonical inclusion  $\iota_A$  is an isometry, so

$$\begin{aligned} \sigma(a) &= \sigma(\iota_A(a)) \\ &\subseteq \overline{D(0, \|\iota_A(a)\|)} \\ &= \overline{D(0, \|a\|)}. \end{aligned}$$

- (3) The common denominator expansion

$$\begin{aligned} R_a(\mu) - R_a(\lambda) &= (a - \mu 1_A)^{-1} - (a - \lambda 1_A)^{-1} \\ &= (a - \mu 1_A)((a - \lambda 1_A) - (a - \mu 1_A))(a - \lambda 1_A)^{-1} \\ &= R_a(\mu)(\mu - \lambda)1_A R_a(\lambda). \end{aligned}$$

When  $\mu \neq \lambda$ , we have

$$\frac{R_a(\mu) - R_a(\lambda)}{\mu - \lambda} = R_a(\mu)R_a(\lambda).$$

We note that  $R_a = i \circ t: \rho(a) \rightarrow A$  is continuous, so for all  $\lambda \in \rho(a)$ , we have

$$\begin{aligned} R_a(\lambda) &= \lim_{\mu \rightarrow \lambda} \frac{R_a(\mu) - R_a(\lambda)}{\mu - \lambda} \\ &= \lim_{\mu \rightarrow \lambda} R_a(\mu)R_a(\lambda) \\ &= R_a(\lambda)^2. \end{aligned}$$

- (4) Let  $A$  be unital, and assume  $a \neq 0$ . Suppose toward contradiction  $\sigma(a) = \emptyset$ , meaning  $\rho(a) = \mathbb{C}$ . Then,  $R_a: \mathbb{C} \rightarrow A$  is entire, so for  $\lambda \neq 0$ , we have

$$\begin{aligned} \|R_a(\lambda)\| &= \|(a - \lambda 1_A)^{-1}\| \\ &= \left\| \left( \lambda \left( \lambda^{-1} a - 1_A \right) \right)^{-1} \right\| \\ &= \left\| \lambda^{-1} \left( \lambda^{-1} a - 1_A \right)^{-1} \right\| \\ &= |\lambda|^{-1} \left\| \left( \lambda^{-1} a - 1_A \right)^{-1} \right\| \\ &\rightarrow 0. \end{aligned}$$

Thus,  $R_a$  is also bounded, and specifically  $R_a \in C_0(\mathbb{C})$ , so by Liouville's theorem,  $R_a = 0$ . However, since  $R_a(0) = a^{-1} \neq 0$ , we have a contradiction. Thus,  $\sigma(a) \neq \emptyset$ .

If  $A$  is nonunital, then  $\sigma(a) = \sigma(\iota_A(a)) \neq \emptyset$ , and  $0 \in \sigma(a)$  since  $A$  does not contain any invertible elements. □

**Definition.** Let  $A$  be a Banach algebra. For each  $a \in A$ , we define the spectral radius to be

$$r(a) = \sup_{\lambda \in \sigma(a)} |\lambda|.$$

**Example.** Let  $\Omega$  be a compact Hausdorff space, and let  $f \in C(\Omega)$ . By compactness, there is  $x \in \Omega$  with  $|f(x)| = \|f\|_u$ . Thus, for some  $\omega \in \mathbb{T}$ , we have  $f(x) = \omega \|f\|_u$ .

It follows that  $f - \omega \|f\|_u 1_\Omega$  is noninvertible, so  $\omega \|f\|_u \in \sigma(f)$ . Thus,  $\|f\|_u = r(f)$ .

**Corollary:** If  $A$  is a Banach algebra with  $a \in A$ , then there is  $\lambda \in \sigma(a)$  with  $|\lambda| = r(a)$ .

*Proof.* The spectrum is nonempty and compact, and the map  $\lambda \mapsto |\lambda|$  is continuous, so its supremum is attained. □

What makes Banach algebras special is that we are able to relate the algebraic aspects of an element (like the spectrum) and the analytic aspects of that element (like the norm), through the following result.

**Proposition:** Let  $A$  be a Banach algebra, and let  $a \in A$ . Then,

$$r(a) = \lim_{n \rightarrow \infty} \|a^n\|^{1/n}.$$

*Proof.* We assume  $A$  admits a unit, and assume  $a \neq 0$ .

Fix  $\lambda \in \sigma(a)$ . From spectral mapping of polynomials, we know that  $\lambda^n \in \sigma(a^n)$ . Thus, we get

$$\begin{aligned} |\lambda|^n &= |\lambda^n| \\ &\leq \|a^n\|, \end{aligned}$$

meaning

$$|\lambda| \leq \|a^n\|^{1/n}.$$

We see that the sequence  $\left( \|a^n\|^{1/n} \right)_n$  is bounded, as

$$\|a^n\|^{1/n} \leq (\|a\|^n)^{1/n}$$

$$= \|a\|,$$

so

$$|\lambda| \leq \liminf_{n \rightarrow \infty} \|a^n\|^{1/n} < \infty.$$

Since this holds for all  $\lambda \in \sigma(a)$ , we have  $r(a) \leq \liminf_{n \rightarrow \infty} \|a^n\|^{1/n}$ .

To establish the opposite direction, let  $\Omega$  be the set of all complex numbers with modulus strictly less than  $r(a)^{-1}$ , where we set  $r(a)^{-1} = \infty$  if  $r(a) = 0$ . If  $0 \neq z \in \Omega$ , then  $|z|^{-1} > r(a)$ , so  $z^{-1} \in \rho(a)$ , and  $a - z^{-1}1_A \in GL(A)$ . Thus, we get that  $1_A - za = -z(a - z^{-1}1_A) \in GL(A)$ . If  $z = 0$ , then it is clear that  $1_A - za = 1_A \in GL(A)$ .

We consider the map  $F: \Omega \rightarrow A$ , given by  $F(z) = (1_A - za)^{-1}$ . We can see that  $F$  is holomorphic on  $\Omega$ . Fix  $\varphi \in A^*$ , and set  $f = \varphi \circ F: \Omega \rightarrow \mathbb{C}$ . Since  $f$  is the composition of holomorphic maps,  $f$  is holomorphic. From complex function theory, we have that  $f$  is analytic, so there is a unique sequence  $(\alpha_n)_n \subseteq \mathbb{C}$  such that

$$f(z) = \sum_{n=0}^{\infty} \alpha_n z^n,$$

which converges absolutely and uniformly on compact subsets of  $\Omega$ . Seeing as  $r(a) \leq \|a\|$ , we have  $\frac{1}{\|a\|} \leq \frac{1}{r(a)}$ , so the open disk  $D(0, \|a\|^{-1})$  is a subset of  $\Omega$ . For any  $w \in D(0, \|a\|^{-1})$ , we see that  $\|wa\| = |w|\|a\| < 1$ , so we get that

$$\begin{aligned} F(w) &= (1_A - wa)^{-1} \\ &= \sum_{n=0}^{\infty} w^n a^n. \end{aligned}$$

Applying  $\varphi$ , we get

$$\begin{aligned} \sum_{n=0}^{\infty} \alpha_n w^n &= f(w) \\ &= \varphi(F(w)) \\ &= \varphi\left(\sum_{n=0}^{\infty} w^n a^n\right) \\ &= \sum_{n=0}^{\infty} w^n \varphi(a^n), \end{aligned}$$

so, since Taylor expansions of holomorphic functions are unique, we have

$$f(z) = \sum_{n=0}^{\infty} \varphi(a^n) z^n.$$

Let  $t > r(a)$ , and set  $\zeta = \frac{1}{t} \in \Omega$ . Since the series for  $f(z)$  converges for all  $z$ , we have that  $|\varphi(a^n)\zeta^n| \rightarrow 0$  as  $n \rightarrow \infty$ , meaning  $\sup_{n \geq 0} |\varphi(\zeta^n a^n)| < \infty$ . Since this holds for all  $\varphi \in A^*$ , the uniform boundedness principle gives  $\sup_{n \geq 0} \|\zeta^n a^n\| = C < \infty$ . Thus, we have  $\|a^n\| \leq \frac{C}{|\zeta|^n}$ , so  $\|a^n\|^{1/n} \leq \frac{C^{1/n}}{|\zeta|}$ . Thus,

$$\limsup_{n \rightarrow \infty} \|a^n\|^{1/n} \leq \limsup_{n \rightarrow \infty} \frac{C^{1/n}}{|\zeta|}$$

$$\begin{aligned}
&= \frac{1}{|\zeta|} \\
&= t.
\end{aligned}$$

Since this holds for all  $t > r(a)$ , we get

$$\limsup_{n \rightarrow \infty} \|a^n\|^{1/n} \leq r(a).$$

Thus, we have

$$\begin{aligned}
r(a) &\leq \liminf_{n \rightarrow \infty} \|a^n\|^{1/n} \\
&\leq \limsup_{n \rightarrow \infty} \|a^n\|^{1/n} \\
&\leq r(a).
\end{aligned}$$

If  $A$  is nonunital, we consider the unitization,  $\tilde{A}$ , and find

$$\begin{aligned}
r(a) &= r(\iota_A(a)) \\
&= \lim_{n \rightarrow \infty} \|\iota_A(a)^n\|^{1/n} \\
&= \lim_{n \rightarrow \infty} \|\iota_A(a^n)\|^{1/n} \\
&= \lim_{n \rightarrow \infty} \|a^n\|^{1/n}.
\end{aligned}$$

□

We can use the established fact that, for a normal element  $b \in A$ ,

$$\|b\|^{2^k} = \|b^{2^k}\|$$

for all  $k \geq 0$  to establish an important fact about the spectral radius of normal elements.

**Proposition:** Let  $A$  be a  $C^*$ -algebra. For any normal  $b \in A$ , we have

$$r(b) = \|b\|$$

*Proof.* We compute

$$\begin{aligned}
r(b) &= \lim_{n \rightarrow \infty} \|b^n\|^{1/n} \\
&= \lim_{k \rightarrow \infty} \|b^{2^k}\|^{1/2^k} \\
&= \lim_{k \rightarrow \infty} (\|b\|)^{(2^k)(1/2^k)} \\
&= \|b\|.
\end{aligned}$$

□

**Proposition:** Let  $A$  be a  $*$ -algebra with norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$  such that  $A$  is a  $C^*$ -algebra when equipped with both of these norms. Then,  $\|\cdot\|_1 = \|\cdot\|_2$ .

*Proof.* Let  $a \in A$ . We have

$$\begin{aligned}
\|a\|_1^2 &= \|a^*a\|_1 \\
&= r(a^*a) \\
&= \|a^*a\|_2 \\
&= \|a\|_2^2.
\end{aligned}$$

□

**Exercise:** Let  $a$  and  $b$  be similar elements in a unital  $C^*$ -algebra  $A$ . If  $A$  is normal, prove that  $\|a\| \leq \|b\|$ , with equality if  $b$  is normal as well.

*Proof.* Since  $a$  and  $b$  are similar elements, we must have that  $\sigma(a) = \sigma(b)$ , so  $r(a) = r(b)$ . Thus, we have

$$\begin{aligned}\|a\| &= r(a) \\ &= r(b) \\ &\leq \|b\|.\end{aligned}$$

□

**Proposition:** If  $\varphi: A \rightarrow B$  is a  $*$ -homomorphism between  $C^*$ -algebras, then  $\varphi$  is contractive (i.e.,  $\|\varphi\|_{\text{op}} \leq 1$ ).

*Proof.* Assume  $A$  and  $B$  are unital, as well as  $\varphi$ . Let  $x \in A$  be normal, so  $\varphi(x)$  is normal too. Thus, we have  $\sigma(\varphi(x)) \subseteq \sigma(x)$ , so

$$\begin{aligned}\|\varphi(x)\| &= r(\varphi(x)) \\ &\leq r(x) \\ &= \|x\|.\end{aligned}$$

If  $a \in A$  is arbitrary, then  $a^*a$  is normal, so

$$\begin{aligned}\|\varphi(a)\|^2 &= \|\varphi(a)^* \varphi(a)\| \\ &= \|\varphi(a^*a)\| \\ &\leq \|a^*a\| \\ &= \|a\|^2.\end{aligned}$$

In the general case, we know that a  $*$ -homomorphism extends to the unital  $*$ -homomorphism on the unitization,  $\tilde{\varphi}: \tilde{A} \rightarrow \tilde{B}$ . Since  $A$  and  $B$  are  $C^*$ -algebras, the  $*$ -homomorphism  $\tilde{\varphi}: A^1 \rightarrow B^1$  is contractive, as

$$\begin{aligned}\|\varphi(a)\| &= \|(\varphi(a), 0)\|_1 \\ &= \|\tilde{\varphi}(a, 0)\|_1 \\ &\leq \|(a, 0)\|_1 \\ &= \|a\|.\end{aligned}$$

□

Since the spectrum of any element in a Banach algebra is nonempty, we get the following structural result.

**Theorem (Mazur's Theorem):** Let  $A$  be a unital Banach algebra such that  $GL(A) = A \setminus \{0\}$ . Then,  $\varphi: \mathbb{C} \rightarrow A$ , given by  $\varphi(z) = z1_A$  is an isometric unital algebra isomorphism.

*Proof.* We only need show that  $\varphi$  is onto, as by construction the map is linear, unital, multiplicative, and isometric.

Let  $a \in A$ . We know that  $\sigma(a) \neq \emptyset$ , so there is some  $\lambda \in \sigma(a)$  such that  $a - \lambda 1_A \notin GL(A)$ . Thus,  $a - \lambda 1_A = 0$ , so  $a = \lambda 1_A = \varphi(\lambda)$ . □

**Corollary:** If  $A$  is a unital, commutative, and simple Banach algebra, then  $A \cong \mathbb{C}$ .

*Proof.* Let  $0 \neq a \in A$ . Since  $A$  is commutative,  $\text{ideal}(a) = \{\chi a \mid \chi \in A\}$ . Since  $A$  is simple,  $\text{ideal}(a) = A$ , so  $1_A \in A$ , so  $a$  is invertible. By Mazur's theorem, we have  $A \cong \mathbb{C}$ . □

**Remark:** The above fails without a complete norm. The algebra of rational functions,

$$C(z) = \left\{ \frac{p(z)}{q(z)} \mid p, q \in \mathbb{C}[z], q \neq 0 \right\}$$

is unital, commutative, and every nonzero element is invertible (hence simple), but  $C(z)$  is not isomorphic to  $\mathbb{C}$ .



## The Character Space

Recall that if  $A$  is a  $\mathbb{C}$ -algebra, a character on  $A$  is a nonzero algebra homomorphism  $h: A \rightarrow \mathbb{C}$ .<sup>vi</sup> We set  $\Omega(A)$  to be the set of all characters on  $A$ .

The aim of this subsection will be to establish a correspondence between characters on a unital Banach algebra and maximal ideals in the algebra.

**Proposition:** Let  $A$  be a Banach algebra. Every character on  $A$  is bounded, with  $\|h\|_{\text{op}} \leq 1$ . If  $A$  is unital, then  $\|h\|_{\text{op}} = 1$ .

*Proof.* Suppose  $A$  is unital. Let  $a \in A$ . We know that  $h(a) \in \sigma(a)$  for any  $h \in \Omega(A)$ . Since  $A$  is a Banach algebra, we know that  $|h(a)| \leq \|a\|$ . Thus,  $\|h\|_{\text{op}} \leq 1$ . Since  $h(1_A) = 1$ , we get  $\|h\|_{\text{op}} = 1$ .

If  $A$  is nonunital, then  $\bar{h}: \tilde{A} \rightarrow \mathbb{C}$  is also a character. We see that  $\|\bar{h}\|_{\text{op}} = 1$ , and for  $a \in A$ , we get

$$\begin{aligned} \|h(a)\| &= \|\bar{h}(a, 0)\| \\ &\leq \|(a, 0)\| \\ &= \|a\|, \end{aligned}$$

so  $\|h\|_{\text{op}} \leq 1$ . □

We will endow the set  $\Omega(A) \subseteq B_{A^*} \subseteq A^*$  with the weak\* topology.

**Definition.** Let  $A$  be a Banach algebra with  $\Omega(A) \neq \emptyset$ . The pair  $(\Omega(A), w^*)$  is called the character space of  $A$ .

**Proposition:** Let  $A$  be a Banach algebra.

- (1) If  $A$  is unital, then the character space is compact Hausdorff.
- (2) If  $A$  is nonunital, then the character space is LCH and its one-point compactification is homeomorphic to  $(\Omega(\tilde{A}), w^*)$ .

*Proof.*

- (1) We only need to show that  $\Omega(A) \subseteq S_{A^*}$  is  $w^*$ -closed. Suppose  $(h_\alpha)_\alpha \xrightarrow{w^*} h$ , where  $(h_\alpha)_\alpha \subseteq \Omega(A)$ , and  $h \in A^*$ . If  $a, b \in A$ , then we have  $(h_\alpha(ab))_\alpha \rightarrow h(ab)$ , and also  $(h_\alpha(ab)) = (h_\alpha(a)h_\alpha(b)) \rightarrow h(a)h(b)$ . We show the latter statement, using the fact that  $h$  is contractive. We have

$$\begin{aligned} \|h_\alpha(a)h_\alpha(b) - h(a)h(b)\| &= \|h_\alpha(a)h_\alpha(b) - h_\alpha(a)h(b)\| + \|h_\alpha(a)h(b) - h(a)h(b)\| \\ &\leq \|h_\alpha(a)\| \|h_\alpha(b) - h(b)\| + \|h_\alpha(a) - h(a)\| \|h(b)\| \\ &\leq \|a\| \|h_\alpha(b) - h(b)\| + \|h_\alpha(a) - h(a)\| \|h(b)\| \\ &\rightarrow 0. \end{aligned}$$

Since limits are unique, we have  $h(ab) = h(a)h(b)$ , so  $h$  is multiplicative. Since  $h \neq 0$ , as  $h_\alpha(1_A) = 1 = h(1_A)$ , we have  $h \in \Omega(A)$ . Thus,  $\Omega(A) \subseteq S_{A^*}$  is  $w^*$ -closed, hence  $w^*$ -compact.

- (2) The map  $\Omega(A) \rightarrow \Omega(\tilde{A})$  that sends  $h \mapsto \bar{h}$  is an injection. This map is a homeomorphism onto its range, as  $(h_\alpha)_\alpha \xrightarrow{w^*} h$  if and only if  $(\bar{h}_\alpha)_\alpha \xrightarrow{w^*} \bar{h}$ . We know that  $\Omega(A)$  is LCH. Since the one-point compactification of a LCH is unique up to homeomorphism, we have  $\Omega(A)_\infty \cong \Omega(\tilde{A})$  are homeomorphic.

---

<sup>vi</sup>Such an algebra homomorphism is automatically surjective.

□

**Remark:** If  $A$  is nonunital, the character space is not necessarily  $w^*$ -closed. For instance, if  $A = C_0(\mathbb{R})$ , then the sequence of characters  $\delta_n(f) = f(n)$  converges in  $w^*$  to 0, but 0 is not a character.

**Proposition:** Let  $A$  be a Banach algebra.

- (1) If  $B$  is a subalgebra, then  $\overline{B} \subseteq A$  is a subalgebra.
- (2) If  $I \subseteq A$  is an ideal, then  $\overline{I} \subseteq A$  is an ideal.
- (3) If  $A$  is unital, and  $I \subsetneq A$  is a proper ideal, then  $\overline{I} \subsetneq A$  is a proper ideal.

*Proof.* We will prove (3).

Suppose  $I \subsetneq A$  is a proper ideal. Suppose toward contradiction that  $\overline{I} = A$ . Then,  $1_A \in \overline{I}$ , so there exists  $x \in I$  with  $\|1_A - x\| < 1$ . We have  $x = 1_A - (1_A - x) \in GL(A)$ , so  $1_A = x^{-1}x \in I$ , so  $I = A$ , which is a contradiction. □

**Remark:** It is important for there to be a unit. For instance,  $C_c(\mathbb{R}) \subseteq C_0(\mathbb{R})$  is a proper ideal, but  $\overline{C_c(\mathbb{R})} = C_0(\mathbb{R})$  is not proper.

**Corollary:** Let  $A$  be a unital Banach algebra, and suppose  $M \subseteq A$  is a maximal ideal.

- (1)  $M$  is closed.
- (2) If  $A$  is commutative, then  $A/M \cong \mathbb{C}$  as Banach algebras.

*Proof.*

- (1) Since  $M \subseteq A$  is a proper ideal, as is  $\overline{M}$ , so  $\overline{M} = M$  as  $M$  is maximal.
- (2) Since  $A$  is unital and commutative, and  $M$  is closed,  $A/M$  is a unital and commutative Banach algebra. Since  $M$  is maximal,  $A/M$  is simple, so  $GL(A/M) = A/M \setminus \{0\}$ , so  $A/M \cong \mathbb{C}$  by Mazur's theorem. □

A unital Banach algebra may not have any nontrivial ideals. However, if  $A$  is commutative, and  $A \neq \mathbb{C}$ , then there are a lot of maximal ideals.

**Theorem:** Let  $A$  be a unital and commutative Banach algebra. There is a one to one correspondence between the collection of all maximal ideals in  $A$ ,

$$\mathcal{M}(A) = \{M \subseteq A \mid M \text{ is a maximal ideal}\},$$

and the set of all characters on  $A$ ,  $\Omega(A)$ , given by  $h \leftrightarrow \ker(h)$ .

*Proof.* We consider the map  $\kappa: \Omega(A) \rightarrow \mathcal{M}(A)$ , given by  $h \mapsto \ker(h)$ .

If  $\ker(h_1) = \ker(h_2)$ , then  $h_1 = \alpha h_2$ , but since characters are unital, we must have  $h_1 = h_2$ , hence the map is injective.

Let  $M \in \mathcal{M}(A)$ . Then,  $A/M \cong \mathbb{C}$  as Banach algebras. Consider the sequence of mappings,  $\pi: A \rightarrow A/M$ , followed by  $\varphi: A/M \rightarrow \mathbb{C}$ , where  $\pi$  is the canonical quotient map and  $\varphi$  is the isometric isomorphism of Banach algebras. Then, the algebra homomorphism  $h_M = \varphi \circ \pi$  is nonzero, as  $h(a) \neq 0$  for all  $a \in A/M \neq \emptyset$ . Since  $\ker(h_M) = M$ , the map  $\kappa$  is onto. □

**Remark:** Some textbook authors refer to the space  $(\Omega(A), w^*)$  as the maximal ideal space when  $A$  is a unital and commutative Banach algebra.

**Theorem:** Let  $A$  be a commutative Banach algebra.

(1) If  $A$  is unital, then  $\Omega(A) \neq \emptyset$ . Further, for each  $a \in A$ ,

$$\sigma(a) = \{h(a) \mid h \in \Omega(A)\}.$$

(2) If  $A$  is nonunital, then for each  $a \in A$ ,

$$\sigma(a) = \{h(a) \mid h \in \Omega(A)\} \cup \{0\}.$$

*Proof.*

(1) For any algebra  $A$  and character  $h \in \Omega(A)$ , we have  $h(a) \in \sigma(a)$ .

Let  $\lambda \in \sigma(a)$ , so  $z = a - \lambda 1_A \notin GL(A)$ . Consider the ideal generated by  $z$ ,  $I$ , determined by

$$\text{ideal}(z) = \{xz \mid x \in A\}.$$

Since  $z$  is not invertible,  $1_A \notin I$ , so  $I$  is proper, and thus contained in a maximal ideal  $M$ . There is some  $h \in \Omega(A)$  such that  $\ker(h) = M$ , so  $h|_I = 0$ , and  $h(a - \lambda 1_A) = 0$ , so  $h(a) = \lambda$ .

(2) If  $A$  is nonunital, then we know that

$$\Omega(\tilde{A}) = \{\bar{h} \mid h \in \Omega(A)\} \cup \{\pi\}.$$

If  $a \in A$ , we have

$$\begin{aligned} \sigma(a) &= \sigma(\iota_A(a)) \\ &= \{h(a) \mid h \in \Omega(A)\} \cup \{\pi(\iota_A(a))\} \\ &= \{h(a) \mid h \in \Omega(A)\} \cup \{0\}. \end{aligned}$$

□

**Example.** We have studied the characters of  $C(X)$ , where  $X$  is a compact Hausdorff space. We have shown that the map  $x \mapsto \delta_x$ , where  $\delta_x(f) = f(x)$ , is a bijection. We will show that  $\delta$  is a homeomorphism when  $\Omega(C(X))$  is endowed with the weak\* topology.

If  $(x_\alpha)_\alpha$  is a net in  $X$ , then since  $X$  is compact Hausdorff, it is completely regular, so for all  $f \in C(X)$ ,

$$\begin{aligned} (x_\alpha)_\alpha \rightarrow x &\Leftrightarrow (f(x_\alpha))_\alpha \rightarrow f(x) \\ &\Leftrightarrow (\delta_{x_\alpha}(f)) \rightarrow \delta_x(f) \\ &\Leftrightarrow \delta_{x_\alpha} \xrightarrow{w^*} \delta_x. \end{aligned}$$

Now, we can classify all algebra homomorphisms  $\varphi: C(X) \rightarrow C(Y)$ , when  $X$  and  $Y$  are compact Hausdorff spaces.

Recall that if  $\tau: Y \rightarrow X$  is a continuous function between compact Hausdorff spaces, we get a unital, contractive  $*$ -homomorphism  $T_\tau: C(X) \rightarrow C(Y)$  given by  $T_\tau(f) = f \circ \tau$ , and if  $\tau$  is injective, then  $T_\tau$  is onto, and if  $\tau$  is onto, then  $T_\tau$  is isometric.

**Proposition:** Let  $X$  and  $Y$  be compact Hausdorff spaces, and suppose  $\phi: C(X) \rightarrow C(Y)$  is a unital algebra homomorphism. Then, there exists a continuous function  $\tau: Y \rightarrow X$  such that  $\phi = T_\tau$ . Consequently,  $\phi$  is contractive and  $*$ -preserving. We also have  $\phi$  is injective if and only if it is isometric.

*Proof.* Given  $y \in Y$ , the character  $\delta_y$  composed with  $\phi$  gives a character on  $C(X)$ ,  $\delta_y \circ \phi: C(X) \rightarrow \mathbb{C}$ . Thus, there exists a unique  $x \in X$  such that  $\delta_y \circ \phi = \delta_x$ . We have a map  $\tau: Y \rightarrow X$  defined by  $\tau(y) = x$  such that  $\delta_y \circ \phi = \delta_{\tau(y)}$ .

For all  $f \in C(X)$  and  $y \in Y$ , we have

$$\begin{aligned} f \circ \tau(y) &= f(\tau(y)) \\ &= \delta_{\tau(y)}(f) \\ &= \delta_y \circ \phi(f) \\ &= \phi(f)(y), \end{aligned}$$

so  $f \circ \tau = \phi(f)$ .

To show that  $\tau$  is continuous, let  $(y_\alpha)_\alpha$  be a net in  $Y$  converging to  $y$ . Then,  $(\delta_{y_\alpha})_\alpha \xrightarrow{w^*} \delta_y$  in  $(C(Y))^*$ , so  $(\delta_{y_\alpha} \circ \phi)_\alpha \xrightarrow{w^*} \delta_y \circ \phi$  in  $(C(X))^*$ . Thus,  $(\delta_{\tau(y_\alpha)})_\alpha \xrightarrow{w^*} \delta_{\tau(y)}$ , so  $(\tau(y_\alpha))_\alpha \rightarrow \tau(y)$  in  $X$ .

Since  $\phi = T_\tau$ ,  $\phi$  is a contractive  $*$ -homomorphism, and it is injective if and only if it is isometric.  $\square$

**Example.** We turn our attention to the algebra  $\ell_1(\mathbb{Z})$  (where multiplication is convolution), and examine the character space of this algebra.

Let  $z \in \mathbb{T}$ . Consider  $h_z(f) = \sum_{n \in \mathbb{Z}} f(n)z^n$ . This series is absolutely convergent as  $|z| = 1$  and  $f \in \ell_1(\mathbb{Z})$ . It is the case that  $h_z$  is linear, and  $h_z \neq 0$ , as  $h_z(e_0) = 1$ . If  $f, g \in \ell_1(\mathbb{Z})$ , then

$$\begin{aligned} h_z(f)h_z(g) &= \left( \sum_{k \in \mathbb{Z}} f(k)z^k \right) \left( \sum_{l \in \mathbb{Z}} g(l)z^l \right) \\ &= \sum_{k, l \in \mathbb{Z}} f(k)g(l)z^{k+l} \\ &= \sum_{n, l \in \mathbb{Z}} f(n-l)g(l)z^n & n = k+l \\ &= \sum_{n \in \mathbb{Z}} \left( \sum_{l \in \mathbb{Z}} f(n-l)g(l) \right) z^n \\ &= \sum_{n \in \mathbb{Z}} f \cdot g(n)z^n \\ &= h_z(f \cdot g). \end{aligned}$$

Thus,  $h_z$  is a character. If  $h_{z_1} = h_{z_2}$ , then  $z_1 = h_{z_1}(e_1) = h_{z_2}(e_1) = z_2$ , so we have an injective map  $\mathbb{T} \rightarrow \Omega(\ell_1(\mathbb{Z}))$ , given by  $z \mapsto h_z$ .

We will show this map is onto. Let  $h \in \Omega(\ell_1(\mathbb{Z}))$ , and set  $z = h(e_1)$ . We claim that  $|z| = 1$ . Note that

$$\begin{aligned} 1 &= h(e_0) \\ &= h(e_1 \cdot e_{-1}) \\ &= h(e_1)h(e_{-1}). \end{aligned}$$

We have  $|h(e_1)| \leq \|h\| \|e_1\| = 1$  and  $|h(e_{-1})| \leq \|h\| \|e_{-1}\| = 1$ , so  $|h(e_1)| = 1$ .

If  $f \in \ell_1(\mathbb{Z})$ , we write  $f = \sum_{n \in \mathbb{Z}} f(n)e_1^n$  as a norm-convergent sum, so we have

$$h(f) = h \left( \sum_{n \in \mathbb{Z}} f(n)e_1^n \right)$$

$$\begin{aligned}
&= \sum_{n \in \mathbb{Z}} f(n) h(e_1)^n \\
&= \sum_{n \in \mathbb{Z}} f(n) z^n \\
&= h_z(f).
\end{aligned}$$

We claim the map  $\mathbb{T} \rightarrow \Omega(\ell_1(\mathbb{Z}))$  is a homeomorphism. It suffices to show continuity, since  $\mathbb{T}$  is compact and  $\Omega(\ell_1(\mathbb{Z}))$  is Hausdorff.

Let  $(z_n)_n \rightarrow z$  in  $\mathbb{T}$ . We show that  $(h_{z_n})_n \xrightarrow{w^*} h_z$  in  $\ell_1(\mathbb{Z})^*$ . The set  $\{e_k \mid k \in \mathbb{Z}\}$  is total, and  $(h_{z_n})_n$  is bounded, so we only need show that  $(h_{z_n}(e_k))_n \rightarrow h_z(e_k)$  for a fixed  $k \in \mathbb{Z}$ . Note that  $h_w(e_k) = \sum_{n \in \mathbb{Z}} e_k(n) w^n = w^k$  for  $w \in \mathbb{T}$ , so

$$\begin{aligned}
|h_{z_n}(e_k) - h_z(e_k)| &= |z_n^k - z^k| \\
&\rightarrow 0.
\end{aligned}$$

Thus,  $\Omega(\ell_1(\mathbb{Z})) \cong \mathbb{T}$ .

**Example.** Now, we turn our attention to the character space of the (full) group  $C^*$ -algebra,  $C^*(\Gamma)$ , where  $\Gamma$  is an abelian discrete group.

The unit circle  $\mathbb{T}$  equipped with multiplication is an abelian group. A character on a discrete group  $\Gamma$  is a group homomorphism,  $\chi: \Gamma \rightarrow \mathbb{T}$ . We define

$$\widehat{\Gamma} = \{\chi \mid \chi \text{ is a character on } \Gamma\}.$$

We endow  $\widehat{\Gamma}$  with the subspace topology inherited from  $\prod_{\Gamma} \mathbb{T}$ . We can see that  $\widehat{\Gamma}$  is closed in this topology, hence compact.

If  $\chi: \Gamma \rightarrow \mathbb{T}$  is a character on  $\Gamma$ , we set

$$h_\chi(f) = \sum_{t \in \Gamma} f(t) \chi(t)$$

as a finite sum. Note that  $h_\chi(\delta_t) = \chi(t)$ , where  $t \in \Gamma$ . We claim that  $h_\chi$  is a character on the unital  $*$ -algebra  $\mathbb{C}[\Gamma]$ . It is the case that  $h_\chi$  is linear. For  $f, g \in \mathbb{C}[\Gamma]$ , we compute

$$\begin{aligned}
h_\chi(f \cdot g) &= \sum_{t \in \Gamma} (f \cdot g)(t) \chi(t) \\
&= \sum_{t \in \Gamma} \left( \sum_{s \in \Gamma} f(s) g(s^{-1}t) \right) \chi(t) \\
&= \sum_{s, t \in \Gamma} f(s) g(s^{-1}t) \chi(ss^{-1}t) \\
&= \sum_{s, t \in \Gamma} f(s) \chi(s) g(s^{-1}t) \chi(s^{-1}t) \\
&= \sum_{s, r \in \Gamma} f(s) \chi(s) g(r) \chi(r) \\
&= \left( \sum_{s \in \Gamma} f(s) \chi(s) \right) \left( \sum_{r \in \Gamma} g(r) \chi(r) \right) \\
&= h_\chi(f) h_\chi(g),
\end{aligned}$$

and

$$\begin{aligned}
 h_\chi(f^*) &= \sum_{t \in \Gamma} f^*(t) \chi(t) \\
 &= \sum_{t \in \Gamma} \overline{f(t^{-1})} \chi(t) \\
 &= \sum_{s \in \Gamma} \overline{f(s)} \chi(s^{-1}) \\
 &= \sum_{s \in \Gamma} \overline{f(s) \chi(s)} \\
 &= \overline{\sum_{s \in \Gamma} f(s) \chi(s)} \\
 &= \overline{h_\chi(f)}.
 \end{aligned}$$

By the universal property of  $C^*(\Gamma)$ , the unital  $*$ -homomorphism  $h_\chi$  extends to a unital  $*$ -homomorphism  $h_\chi: C^*(\Gamma) \rightarrow \mathbb{C}$ , a character on the  $C^*$ -algebra  $C^*(\Gamma)$ .

We have a map  $\widehat{\Gamma} \rightarrow \Omega(C^*(\Gamma))$ , given by  $\chi \mapsto h_\chi$ .

The map is injective, as if  $\chi_1, \chi_2 \in \widehat{\Gamma}$  are characters on  $\Gamma$  with  $h_{\chi_1} = h_{\chi_2}$ , then  $\chi_1(t) = h_{\chi_1}(\delta_t) = h_{\chi_2}(\delta_t) = \chi_2(t)$ .

For surjectivity, let  $h \in \Omega(C^*(\Gamma))$ , and define  $\chi_h: \Gamma \rightarrow \mathbb{C}$ , given by  $\chi_h(t) = h(\delta_t)$ . Since  $h$  is multiplicative,  $\chi_h$  is a character on  $\Gamma$ .

We claim that  $h_{\chi_h} = h$ . To see this, if  $t \in \Gamma$ , then

$$\begin{aligned}
 h_{\chi_h}(\delta_t) &= \chi_h(t) \\
 &= h(\delta_t),
 \end{aligned}$$

so by linearity,  $h_{\chi_h} = h$  on  $\mathbb{C}[\Gamma]$ , and by continuity, they agree on  $C^*(\Gamma)$ .

If  $(\chi_\alpha)_\alpha$  is a net of characters in  $\Gamma$  converging to  $\chi \in \widehat{\Gamma}$ , then for all  $t \in \Gamma$ , we have  $h_{\chi_\alpha}(\delta_t) = \chi_\alpha(t) \rightarrow \chi(t) = h_\chi(\delta_t)$ . Thus,  $(h_{\chi_\alpha})_\alpha \xrightarrow{w^*} h_\chi$  in  $\Omega(C^*(\Gamma))$ , and since  $\widehat{\Gamma}$  is compact and  $\Omega(C^*(\Gamma))$  is Hausdorff.

Thus, we have that the character space  $\Omega(C^*(\Gamma))$  is homeomorphic to  $\widehat{\Gamma}$ , the character space of  $\Gamma$ .

### The Gelfand Transform

If  $A$  is unital and commutative, we know that the character space  $\Omega(A)$  is nonempty and maps the spectrum of each element.

**Definition.** Let  $A$  be a commutative Banach algebra with  $\Omega(A) \neq \emptyset$ , and let  $a \in A$ . The map  $\hat{a}: \Omega(A) \rightarrow \mathbb{C}$ , given by  $\hat{a}(h) = h(a)$  is known as the Gelfand transform of  $a$ .

**Proposition:** Let  $A$  be a commutative Banach algebra, with  $\Omega(A) \neq \emptyset$ . Let  $a \in A$ . We have the following:

- (1)  $\hat{a} \in C_0(\Omega(A))$ ;<sup>vii</sup>
- (2) if  $A$  is unital, then  $\text{Ran}(\hat{a}) = \sigma(a)$ , and if  $A$  is nonunital,  $\text{Ran}(\hat{a}) \cup \{0\} = \sigma(a)$ ;
- (3)  $\|\hat{a}\|_\infty = r(a) \leq \|a\|$ ;

---

<sup>vii</sup>Note that if  $\Omega(A)$  is compact, then  $C_0(\Omega(A)) = C(\Omega(A))$ .

- (4) the Gelfand transform of  $A$ ,  $\gamma_A: A \rightarrow C_0(\Omega(A))$ , given by  $\gamma_A(a) = \hat{a}$ , is a contractive algebra homomorphism, and if  $A$  is unital, then  $\gamma_A$  is unital.

*Proof.*

- (1) Continuity follows from the definitions. We see that

$$\begin{aligned} (h_\alpha)_\alpha &\xrightarrow{w^*} h \Rightarrow (h_\alpha(a))_\alpha \rightarrow h(a) \\ &\Rightarrow (\hat{a}(h_\alpha))_\alpha \rightarrow \hat{a}(h). \end{aligned}$$

If  $\varepsilon > 0$ , then the continuity of  $\hat{a}$  implies that  $K_\varepsilon = \{h \in \Omega(A) \mid |\hat{a}(h)| \geq \varepsilon\} \subseteq B_{A^*}$  is weak\*-closed, so it is weak\*-compact. If  $h \in \Omega(A) \setminus K_\varepsilon$ , then  $|\hat{a}(h)| < \varepsilon$ , so  $\hat{a}$  vanishes at infinity.

- (2) If  $A$  is unital, we know that

$$\begin{aligned} \text{Ran}(\hat{a}) &= \{\hat{a}(h) \mid h \in \Omega(A)\} \\ &= \{h(a) \mid h \in \Omega(A)\} \\ &= \sigma(a), \end{aligned}$$

and similarly for the case where  $A$  is nonunital.

- (3) We compute

$$\begin{aligned} \|\hat{a}\|_u &= \sup_{h \in \Omega(A)} |\hat{a}(h)| \\ &= \sup_{h \in \Omega(A)} |h(a)| \\ &= \sup_{\lambda \in \sigma(a)} |\lambda| \\ &= r(a) \\ &\leq \|a\|. \end{aligned}$$

- (4) From (2), we see that  $\gamma_A$  is contractive. To see multiplicativity, we have

$$\begin{aligned} \widehat{ab}(h) &= h(ab) \\ &= h(a)h(b) \\ &= \hat{a}(h)\hat{b}(h) \\ &= \widehat{\hat{a}\hat{b}}(h), \end{aligned}$$

$$\text{so } \gamma_A(ab) = \gamma_A(a)\gamma_A(b).$$

□

We are interested in seeing if the Gelfand map,  $\gamma_A$ , is  $*$ -preserving, in the case where  $A$  is a Banach  $*$ -algebra or  $C^*$ -algebra.

**Proposition:** Let  $A$  be a commutative Banach  $*$ -algebra with  $\Omega(A) \neq \emptyset$ . Let  $\gamma_A: A \rightarrow C_0(\Omega(A))$  be the Gelfand map. The following are equivalent:

- (i)  $\gamma_A$  is  $*$ -preserving;
- (ii) if  $a \in A_{\text{s.a.}}$ , then  $\text{Ran}(\hat{a}) \subseteq \mathbb{R}$ ;
- (iii) every  $h \in \Omega(A)$  is  $*$ -preserving;
- (iv) for every  $h \in \Omega(A)$ ,  $h(A_{\text{s.a.}}) \subseteq \mathbb{R}$ .

If  $A$  satisfies any of these conditions, we say  $A$  is symmetric.

*Proof.* We see that  $\gamma_A$  is  $*$ -preserving if and only if  $\gamma_A(A_{s.a.}) \subseteq C(\Omega(A))_{s.a.} = C(\Omega(A), \mathbb{R})$ . The equivalence of (i) and (ii) follows, and similarly the equivalence between (iii) and (iv).

We only need to show the equivalence between (i) and (iii). Given  $a \in A$ , we have

$$\begin{aligned}\gamma_A(a^*) = \gamma_A(a)^* &\Leftrightarrow \widehat{a^*}(h) = \overline{\widehat{a}(h)} \\ &\Leftrightarrow h(a^*) = \overline{h(a)}.\end{aligned}$$

□

Commutative  $C^*$ -algebras are symmetric, which enables us to realize commutative  $C^*$  algebras as continuous function spaces.

**Proposition:** If  $A$  is a commutative  $C^*$ -algebra, then  $A$  is symmetric.

Additionally,  $\text{Ran}(\gamma_A) \subseteq C_0(\Omega(A))$  is a  $*$ -subalgebra. If  $A$  is unital, then  $\text{Ran}(\gamma_A) \subseteq C(\Omega(A))$  is a unital  $*$ -subalgebra.

*Proof.* Let  $A$  be unital with  $h \in \Omega(A)$ , and let  $a \in A_{s.a.}$ . We will show that  $h(a) \in \mathbb{R}$ .

Write  $h(a) = \alpha + i\beta$ , and we will show that  $\beta = 0$ . Fix  $t \in \mathbb{R}$ , and define  $z = a + it1_A$ . Since  $a = a^*$ ,

$$\begin{aligned}z^*z &= (a + it1_A)^*(a + it1_A) \\ &= a^*a + ita^* - ita + t^21_A \\ &= a^2 + t^21_A.\end{aligned}$$

Notice that

$$\begin{aligned}|h(z)|^2 &= |h(a) + it|^2 \\ &= |\alpha + i\beta + it|^2 \\ &= |\alpha|^2 + |\beta|^2 + 2\beta t + t^2,\end{aligned}$$

and

$$\begin{aligned}|h(z)|^2 &\leq \|z\|^2 \\ &= \|z^*z\| \\ &= \|a^2 + t^21_A\| \\ &\leq \|a\|^2 + t^2.\end{aligned}$$

Thus, we find

$$|\alpha|^2 + |\beta|^2 + 2\beta t \leq \|a\|^2.$$

Since this inequality holds for all  $t \in \mathbb{R}$ , this is only possible if  $\beta = 0$ .

If  $A$  is nonunital, we let  $a \in A_{s.a.}$ , and  $h \in \Omega(A)$ . Then,  $\iota_A(a) \in \widetilde{A}_{s.a.}$ , and  $\bar{h} \in \Omega(\widetilde{A})$ , so

$$\begin{aligned}h(a) &= \bar{h}(\iota_A(a)) \\ &\in \mathbb{R},\end{aligned}$$

so  $A$  is symmetric.

Since  $\gamma_A$  is a  $*$ -homomorphism, we see that  $\text{Ran}(\gamma_A)$  is a  $*$ -subalgebra, which is unital if  $A$  is unital. □



Now, we are interested in understanding when the Gelfand map is isometric. Note that if  $f \in C_0(\Omega)$  for some LCH space  $\Omega$ , then for any  $n \geq 1$ , we have

$$\begin{aligned}\|f\|_u^n &= \left( \sup_{x \in \Omega} |f(x)| \right)^n \\ &= \sup_{x \in \Omega} (|f(x)|)^n \\ &= \sup_{x \in \Omega} |(f(x))^n| \\ &= \|f^n\|_u.\end{aligned}$$

**Proposition:** Let  $A$  be a commutative Banach algebra.

- (1) If  $a \in A$ , then  $\|\gamma_A(a)\|_u = \|a\|$  if and only if  $\|a^{2^k}\| = \|a\|^{2^k}$  for all  $k \in \mathbb{N}$ .
- (2) If  $A$  is a  $C^*$ -algebra, then  $\gamma_A$  is an isometry, and  $\text{Ran}(\gamma_A) \subseteq C_0(\Omega(A))$  is a  $C^*$ -subalgebra. If  $A$  is unital, then  $\text{Ran}(\gamma_A) \subseteq C(\Omega(A))$  is a unital  $C^*$ -subalgebra.

*Proof.*

- (1) If  $\|\gamma_A(a)\|_u = \|a\|$ , then

$$\begin{aligned}\|a^{2^k}\| &\leq \|a\|^{2^k} \\ &= \|\gamma_A(a)\|_u^{2^k} \\ &= \|\gamma(a)^{2^k}\|_u \\ &= \|\gamma_A(a^{2^k})\| \\ &\leq \|a^{2^k}\|,\end{aligned}$$

where we used the aforementioned property of the norm on  $C_0(\Omega)$  and the contractiveness of  $\gamma_A$ .

Thus,  $\|a^{2^k}\| = \|a\|^{2^k}$  for all  $k \in \mathbb{N}$ .

Conversely, suppose  $\|a^{2^k}\| = \|a\|^{2^k}$  for all  $k \in \mathbb{N}$ . By the spectral radius formula, we have

$$\begin{aligned}\|\gamma_A(a)\|_u &= r(a) \\ &= \lim_{n \rightarrow \infty} \|a^n\|^{1/n} \\ &= \lim_{k \rightarrow \infty} \|a^{2^k}\|^{2^{-k}} \\ &= \lim_{k \rightarrow \infty} (\|a\|^{2^k})^{2^{-k}} \\ &= \|a\|.\end{aligned}$$

- (2) Let  $a \in A$ ,  $b = a^*a \in A_{\text{s.a.}}$ . Since  $A$  is symmetric, we get

$$\begin{aligned}\|a\|^2 &= \|a^*a\| \\ &= \|b\| \\ &= \|\gamma_A(b)\| \\ &= \|\gamma_A(a^*a)\| \\ &= \|\gamma_A(a)^*\gamma_A(a)\| \\ &= \|\gamma_A(a)\|^2,\end{aligned}$$

so  $\gamma_A$  is isometric. It is the case that the range of  $\gamma_A$  is a  $C^*$ -subalgebra of  $C_0(\Omega(A))$ .

□

**Corollary:** If  $A$  is a commutative  $C^*$ -algebra, then the characters on  $A$  separate points.

*Proof.* If  $a_1, a_2 \in A$  with  $h(a_1) = h(a_2)$  for all  $h \in \Omega(A)$ , then

$$\begin{aligned}\hat{a}_1(h) = \hat{a}_2(h) &\Leftrightarrow \hat{a}_1 = \hat{a}_2 \\ &\Leftrightarrow \gamma_A(a_1) = \gamma_A(a_2) \\ &\Leftrightarrow a_1 = a_2.\end{aligned}$$

□

**Theorem (Gelfand–Naimark):** Let  $A$  be a commutative  $C^*$ -algebra. The gelfand map  $\gamma_A : A \rightarrow C_0(\Omega(A))$  is an isometric  $*$ -isomorphism. If  $A$  is unital, then  $\gamma_A : A \rightarrow C(\Omega(A))$  is an isometric unital  $*$ -isomorphism.

*Proof.* We only need to show that  $\gamma_A$  is onto. For this, we will use the Stone–Weierstrass theorem.

We start by claiming that  $\text{Ran}(\gamma_A)$  separates points. Suppose  $h_1, h_2 \in \Omega(A)$  are distinct. Then, there exists some  $a \in A$  such that  $h_1(a) \neq h_2(a)$ , so  $\hat{a}(h_1) \neq \hat{a}(h_2)$ .

Next, we show that  $\text{Ran}(\gamma_A)$  has no zeros. If  $h \in \Omega(A)$ , then  $h \neq 0$ , so there is some  $a \in A$  with  $h(a) \neq 0$ , meaning  $\hat{a}(h) \neq 0$ .

Thus, by the Stone–Weierstrass theorem, we have  $\text{Ran}(\gamma_A) = C_0(\Omega(A))$ .

□

**Example.** Let  $\Gamma$  be a discrete abelian group. We have shown that every character  $\chi$  on  $\Gamma$  gives rise to a character  $h_\chi$  on the full group  $C^*$ -algebra,  $C^*(\Gamma)$ , and that the map  $\chi \mapsto h_\chi$  is a homeomorphism.

Composing with the Gelfand map,  $\gamma : C^*(\Gamma) \rightarrow C(\Omega(C^*(\Gamma)))$  gives the evaluation isomorphism,

$$\delta_t \mapsto \text{ev}_t : \chi \mapsto \chi(t).$$

We can give a concrete example. Consider the homeomorphism  $\mathbb{T} \rightarrow \widehat{\mathbb{Z}}$ . We get an isometric  $*$ -isomorphism,  $C(\widehat{\mathbb{Z}}) \rightarrow C(\mathbb{T})$ . Composing with the evaluation isomorphism, we have the isometric  $*$ -isomorphism,

$$\sum_{n \in \mathbb{Z}} c_n \delta_n \mapsto \sum_{n \in \mathbb{Z}} c_n z^n,$$

where the sums are finite.

A useful consequence of the Gelfand–Naimark theorem is spectral permanence.

If  $A$  is a unital subalgebra of a unital algebra  $B$ , and  $a \in A$ , then we can specify  $\sigma_A(a)$  and  $\sigma_B(a)$ , where

$$\begin{aligned}\sigma_B(a) &= \{\lambda \in \mathbb{C} \mid a - \lambda 1_B \notin \text{GL}(B)\} \\ \sigma_A(a) &= \{\lambda \in \mathbb{C} \mid a - \lambda 1_B \notin \text{GL}(A)\}.\end{aligned}$$

It can be seen that  $\sigma_B(a) \subseteq \sigma_A(a)$ , as  $\text{GL}(A) \subseteq \text{GL}(B) \cap A$  always holds.

In a  $C^*$ -algebra, though, this becomes an equality.

**Proposition (Spectral Permanence):** Let  $B$  be a unital  $C^*$ -algebra, and suppose  $A \subseteq B$  is a unital  $C^*$ -algebra of  $B$ .

- (1) If  $z \in A \cap \text{GL}(B)$ , then  $z \in \text{GL}(A)$ .
- (2) For every  $a \in A$ ,  $\sigma_A(a) = \sigma_B(a)$ .

*Proof.* Let  $x \in A \cap GL(B)$  be self-adjoint. We need to show that  $x^{-1} \in A$ . We define  $C = C^*(x, x^{-1}) \subseteq B$  and  $D = C^*(1_B, x) \subseteq A$ .

Since  $x$  is normal, both  $C$  and  $D$  are unital and commutative  $C^*$ -algebras with  $D \subseteq C$ . We will use the following commutative diagram to organize the argument.

$$\begin{array}{ccc} D & \xrightarrow{\iota} & C \\ \gamma_D \downarrow & & \downarrow \gamma_C \\ C(\Omega(D)) & \xrightarrow{\pi} & C(\Omega(C)) \end{array}$$

We have  $\iota$  is inclusion,  $\gamma_C$  and  $\gamma_D$  the isometric  $*$ -isomorphic Gelfand maps, and  $\pi: C(\Omega(D)) \rightarrow C(\Omega(C))$ , defined as  $\pi = \gamma_C \circ \iota \circ \gamma_D^{-1}$ , is a unital and isometric  $*$ -homomorphism. Consequently,  $\text{Ran}(\pi)$  is a unital  $C^*$ -subalgebra of  $C(\Omega(C))$ .

We will show that  $\text{Ran}(\pi)$  separates the points of  $\Omega(C)$ , so we may apply Stone–Weierstrass. Suppose  $h_1, h_2 \in \Omega(C)$  are distinct characters. We claim that  $\hat{x}(h_1) \neq \hat{x}(h_2)$ . If this were not the case, then  $h_1(x) = h_2(x)$ , and

$$\begin{aligned} h_1(x^{-1}) &= \frac{1}{h_1(x)} \\ &= h_2(x^{-1}) \\ &= \frac{1}{h_2(x)}, \end{aligned}$$

so since  $C$  is generated by  $x$  and  $x^{-1}$ , multiplicativity and continuity of  $h_1, h_2$  imply that  $h_1 = h_2$ , which is a contradiction. Since  $\hat{x} \in \text{Ran}(\pi)$ , we have that  $\text{Ran}(\pi)$  separates points, so  $\text{Ran}(\pi) = C(\Omega(C))$ . Since  $\pi$  is surjective, a diagram chase shows that  $\iota$  is surjective, so  $x^{-1} \in D \subseteq A$ .

For the general case, let  $z \in A \cap GL(B)$ , and set  $x = z^*z$ . Since  $GL(B)$  and  $A$  are both closed under involution and multiplication, we see that  $x \in A \cap GL(B)$ . Thus,  $x^{-1} \in A$ . Since  $x^{-1}z^*$  is a left-inverse for  $z$ , and  $z$  is invertible, we must have  $z^{-1} \in A$ .

For the proof of (2), we know that  $\sigma_B(a) \subseteq \sigma_A(a)$ . If  $\lambda \in \rho_B(a)$ , then  $a - \lambda 1_B \in GL(B)$ , so  $a - \lambda 1_B \in GL(A)$ , so  $\lambda \in \rho_A(a)$ . Thus, we get  $\rho_B(a) \subseteq \rho_A(a)$ . Taking complements, we get  $\sigma_A(a) \subseteq \sigma_B(a)$ .  $\square$

**Proposition:** Let  $A$  and  $B$  be  $C^*$ -algebras. A  $*$ -homomorphism,  $\phi: A \rightarrow B$ , is injective if and only if it is isometric.

*Proof.* We assume  $A, B$ , and  $\phi$  are unital. Let  $a \in A$ , and set  $x = a^*a$ . Then,  $\phi(x)$  is a normal element in  $B$ . Set  $C = C^*(x, 1_A) \subseteq A$ , and  $D = C^*(\phi(x), 1_B) \subseteq B$ , which are unital and commutative  $C^*$ -algebras. By continuity, we see that  $\phi|_C(C) \subseteq D$ .

We will use the following commutative diagram to organize the argument.

$$\begin{array}{ccc} C & \xrightarrow{\phi|_C} & D \\ \gamma_C \downarrow & & \downarrow \gamma_D \\ C(\Omega(C)) & \xrightarrow{\pi} & C(\Omega(D)) \end{array}$$

Here,  $\gamma_C$  and  $\gamma_D$  are the isometric  $*$ -isomorphic Gelfand maps, and  $\pi = \gamma_D \circ \phi|_C \circ \gamma_C^{-1}$ . Since  $\pi$  is an injective unital homomorphism between continuous function spaces,  $\pi$  is isometric. We see that

$$\begin{aligned} \|x\| &= \|\gamma_C(x)\| \\ &= \|\pi(\gamma_C(x))\| \end{aligned}$$

$$\begin{aligned}
&= \|\gamma_D(\phi(x))\| \\
&= \|\phi(x)\|,
\end{aligned}$$

so

$$\begin{aligned}
\|a\|^2 &= \|a^*a\| \\
&= \|x\| \\
&= \|\phi(x)\| \\
&= \|\phi(a^*a)\| \\
&= \|\phi(a)^*\phi(a)\| \\
&= \|\phi(a)\|^2,
\end{aligned}$$

meaning  $\phi$  is isometric.

In the non-unital case, we unitize  $\phi$  to yield the unital  $*$ -homomorphism  $\tilde{\phi}: \tilde{A} \rightarrow \tilde{B}$ . Since  $\phi$  is injective, so is  $\tilde{\phi}$ , and  $\tilde{\phi}$  is isometric. Since  $\tilde{\phi} \circ \iota_A = \phi$ , and  $\iota_A$  is isometric, it follows that  $\phi$  is isometric.  $\square$

We may now consider the Gelfand transform of  $\ell_1(\mathbb{Z})$  (where multiplication is given by convolution).

**Example.** We have identified the character space  $\Omega(\ell_1(\mathbb{Z}))$  with  $\mathbb{T}$ . For a given  $z$ , the Gelfand transform is

$$\begin{aligned}
\gamma(f)(h_z) &= \hat{f}(h_z) \\
&= h_z(f) \\
&= \sum_{n \in \mathbb{Z}} f(n)z^n \\
&:= \hat{f}(z).
\end{aligned}$$

Recall that if  $\varphi \in C(\mathbb{T})$ , the Fourier transform of  $\varphi$  is the function

$$\check{\varphi}(n) = \frac{1}{2\pi} \int_0^{2\pi} \varphi(e^{it}) e^{-int} dt.$$

The formal Fourier series of  $\varphi$  is the power series

$$\varphi \sim \sum_{n \in \mathbb{Z}} \check{\varphi}(n) z^n.$$

Suppose  $\varphi \in C(\mathbb{T})$  with  $\varphi = \hat{f}$  for some  $f \in \ell_1(\mathbb{Z})$ . The Fourier coefficient of  $\varphi$  is

$$\begin{aligned}
\check{\varphi}(n) &= \frac{1}{2\pi} \int_0^{2\pi} \varphi(e^{it}) e^{-int} dt \\
&= \frac{1}{2\pi} \int_0^{2\pi} \hat{f}(e^{it}) e^{-int} dt \\
&= \frac{1}{2\pi} \int_0^{2\pi} \left( \sum_{k \in \mathbb{Z}} f(k) e^{ikt} \right) e^{-int} dt \\
&= \frac{1}{2\pi} \sum_{k \in \mathbb{Z}} f(k) \int_0^{2\pi} e^{i(k-n)t} dt \\
&= f(n).
\end{aligned}$$

Thus,  $\check{f} = \check{\varphi} = f$  as functions on  $\mathbb{Z}$ .

We claim that the range of the Gelfand transform is

$$C = \left\{ \varphi \in C(\mathbb{T}) \mid \sum_{n \in \mathbb{Z}} \check{\varphi}(n) z^n \text{ converges absolutely on } \mathbb{T} \right\}.$$

If  $\varphi \in \gamma(\ell_1(\mathbb{Z}))$ , then  $\varphi = \hat{f}$  for some  $f \in \ell_1(\mathbb{Z})$ , and we computed that  $\check{\varphi}(n) = f(n)$  for all  $n \in \mathbb{Z}$ , so  $\sum_{n \in \mathbb{Z}} \check{\varphi}(n) z^n = \sum_{n \in \mathbb{Z}} f(n) z^n$  converges absolutely for all  $z \in \mathbb{T}$ .

Conversely, if the Fourier series of  $\varphi$  converges absolutely, then by taking  $z = 1$ , we get that  $(\check{\varphi}(n))_{n \in \mathbb{Z}} \in \ell_1(\mathbb{Z})$ . Set  $f(n) = \check{\varphi}(n)$ , and observe

$$\begin{aligned} \check{f}(z) &= \hat{f}(h_z) \\ &= h_z(f) \\ &= \sum_{n \in \mathbb{Z}} f(n) z^n \\ &= \sum_{n \in \mathbb{Z}} \check{\varphi}(n) z^n \\ &= \varphi(z). \end{aligned}$$

### Continuous Functional Calculus

Now, we may use the Gelfand–Naimark theorem to apply continuous functions to elements of  $C^*$ -algebras.

Fix a commutative unital Banach algebra  $A$ . If  $a \in A$ , we know that the function  $\hat{a}: \Omega(A) \rightarrow \sigma(a)$ , given by  $\hat{a}(h) = h(a)$  is continuous and onto when  $\Omega(A)$  is endowed with the weak\* topology. By duality, we get an injective unital  $*$ -homomorphism

$$T_{\hat{a}}: C(\sigma(a)) \rightarrow C(\Omega(A)),$$

given by  $T_{\hat{a}} = f \circ \hat{a}$ .

We are interested in the cases where  $T_{\hat{a}}$  is surjective, which requires  $\hat{a}$  to be injective.

**Proposition:** Let  $A$  be a commutative and unital Banach algebra. Let  $a \in A$ . The map  $\hat{a}: \Omega(A) \rightarrow \sigma(a)$  is injective if any of the following conditions hold:

- (i)  $A$  is generated by  $a$  and  $1_A$ ,

$$A = \overline{\{p(a) \mid p \in \mathbb{C}[x]\}}^{\|\cdot\|};$$

- (ii)  $a \in GL(A)$ , and  $A$  is generated as a Banach algebra by  $a$  and  $a^{-1}$ , that is

$$A = \overline{\left\{ \sum_{n=-N}^N c_n a^n \mid N \in \mathbb{N}, c_n \in \mathbb{C} \right\}}^{\|\cdot\|};$$

- (iii)  $A$  is a  $C^*$ -algebra generated by  $a$  and  $1_A$ , that is  $A = C^*(a, 1_A)$ .

*Proof.*

- (i) If  $\hat{a}(h_1) = \hat{a}(h_2)$ , where  $h_1, h_2 \in \Omega(A)$ , then  $h_1(a) = h_2(a)$ . Since  $h_1, h_2$  are multiplicative and linear, we have  $h_1(p(a)) = h_2(p(a))$  for all  $p \in \mathbb{C}[x]$ . Since  $h_1, h_2$  are continuous and agree on a dense subset,  $h_1 = h_2$ .

- (ii) Assuming  $h_1, h_2 \in \Omega(A)$  with  $h_1(a) = h_2(a)$ , since  $a$  is invertible, we know that  $h_i(a) \neq 0$  and  $h_i(a^{-1}) = h_i(a)^{-1}$  for each  $i$ . Thus,  $h_1(a^{-1}) = h_2(a^{-1})$ . Since characters are multiplicative and linear, we see that  $h_1(q(a)) = h_2(q(a))$  for all Laurent polynomials  $q(z) = \sum_{n=-N}^N c_n z^n$ . By continuity,  $h_1 = h_2$ .
- (iii) Characters on a  $C^*$ -algebra are always self-adjoint, so if  $h_1(a) = h_2(a)$ , then  $\overline{h_1(a)} = \overline{h_2(a)}$ . Multiplicativity means  $h_1(w) = h_2(w)$  for all words  $w$ , and by linearity,  $h_1 = h_2$  on  $\text{span}(W)$ . By continuity,  $h_1 = h_2 = C^*(a, 1_A) = \overline{\text{span}}(W)$ .

□

**Corollary:** Let  $A$  be a  $C^*$ -algebra. Suppose  $a \in A$  is normal.

- (1) If  $A$  is unital, and  $C^*(a, 1_A) = A$ , then

$$T_{\hat{a}}(f) = f \circ \hat{a}$$

is a unital isometric  $*$ -isomorphism.

- (2) If  $A$  is nonunital, and  $C^*(a) = A$ , then  $C^*(a, 1_{\tilde{A}}) = \tilde{A}$ , and the unital isometric  $*$ -isomorphism

$$T_{\hat{a}}: C(\sigma(a)) \rightarrow C\left(\Omega\left(\tilde{A}\right)\right)$$

restricts to a  $*$ -isomorphism

$$\{f \in C(\sigma(a)) \mid f(0) = 0\} \rightarrow C_0(\Omega(A)).$$

*Proof.*

- (1) Since the Gelfand transform  $\hat{a}: \Omega(A) \rightarrow \sigma(a)$  is continuous and onto,  $\hat{a}$  is injective, hence a homeomorphism. By duality,  $T_{\hat{a}}: C(\sigma(a)) \rightarrow C(\Omega(A))$  is a unital isometric  $*$ -homomorphism.
- (2) Since  $A$  is nonunital, and  $\tilde{A} = C^*(a, 1_{\tilde{A}})$ , we have that  $0 \in \sigma(a)$ , and  $\Omega(\tilde{A}) = \Omega(A) \cup \{\pi\}$ . We identify  $C_0(\Omega(A))$  with the ideal  $\left\{g \in C\left(\Omega\left(\tilde{A}\right)\right) \mid g(\pi) = 0\right\} \subseteq C\left(\Omega\left(\tilde{A}\right)\right)$ .

Note that if  $f \in C(\sigma(a))$  with  $f(0) = 0$ , then

$$\begin{aligned} T_{\hat{a}}(f)(\pi) &= f \circ \hat{a}(\pi) \\ &= f(\pi(a)) \\ &= f(0) \\ &= 0, \end{aligned}$$

so  $T_{\hat{a}} \in C_0(\Omega(A))$ . Conversely, if  $g(\pi) = 0$  and  $g = T_{\hat{a}}(f)$  for some  $f \in C(\sigma(a))$ , then

$$\begin{aligned} f(0) &= f(\pi(a)) \\ &= f \circ \hat{a}(\pi) \\ &= T_{\hat{a}}(f)(\pi) \\ &= g(\pi) \\ &= 0. \end{aligned}$$

□

**Definition** (Continuous Functional Calculus). Let  $B$  be a unital  $C^*$ -algebra, and suppose  $a \in B$  is normal. Let  $A = C^*(a, 1_B)$  be the commutative unital  $C^*$ -subalgebra  $B$  generated by  $a$ . The functional calculus at  $a$  is the isometric  $*$ -isomorphism  $\phi_a: C(\sigma(a)) \rightarrow A$ , given by  $\phi_a = \gamma_A^{-1} \circ T_{\hat{a}}$ .

We will write  $f(a)$  for  $\phi_a(f)$  for  $f \in C(\sigma(a))$ .

**Theorem** (Properties of the Continuous Functional Calculus): Let  $B$  be a unital  $C^*$ -algebra, and let  $a \in B$  be normal. Set  $A = C^*(a, 1_B)$ , and let

$$\phi_a: C(\sigma(a)) \rightarrow A$$

be the continuous functional calculus. The following hold:

- (1)  $\phi_a(p) = p(a)$  for all polynomials  $\mathbb{C}[x]$  — in particular,  $\phi_a(\iota) = a$  and  $\phi_a(1_{\sigma(a)}) = 1_B$ , where  $\iota: \sigma(a) \rightarrow \mathbb{C}$  is inclusion;
- (2)  $\phi_a(f) = f(a)$  is a normal element in  $A$  for every  $f \in C(\sigma(a))$ ;
- (3)  $\sigma(f(a)) = f(\sigma(a))$  for all  $f \in C(\sigma(a))$  (known as spectral mapping);
- (4) if  $f \in C(\sigma(a))$  and  $g \in C(\sigma(f(a)))$ , then  $g \circ f \in C(\sigma(a))$ , and  $g(f(a)) = g \circ f(a)$ ;
- (5) if  $\pi: B \rightarrow C$  is a unital  $*$ -homomorphism between unital  $C^*$ -algebras, then
  - (i)  $\sigma(\pi(a)) \subseteq \sigma(a)$ ;
  - (ii)  $\pi(a)$  is normal and if  $f \in C(\sigma(a))$ , then  $f(\pi(a)) = \pi(f(a))$ ;
- (6) if  $0 \in \sigma(a)$  and  $f \in C(\sigma(a))$  with  $f(0) = 0$ , or if  $0 \notin \sigma(a)$ , then  $\phi_a(f) \in C^*(a)$ .

*Proof.*

- (1) Since  $\phi_a$  is a unital morphism,  $\phi_a(1_{\sigma(a)}) = 1_B$ . Next, we note that  $T_{\hat{a}}(\iota) = \iota \circ \hat{a} = \hat{a}$ , and  $\gamma_A^{-1}(\hat{a}) = a$ . Thus,  $\phi_a(\iota) = a$ . Since  $\phi_a$  is a homomorphism, we see that  $\phi_a(p) = p(a)$ .
- (2) The image of any normal element under a  $*$ -homomorphism is normal.
- (3) Since  $\phi_a$  is an algebra isomorphism, we know that  $\sigma(\phi_a(f)) = \sigma(f)$  for all  $f \in C(\sigma(a))$ . In particular, we have

$$\begin{aligned} \sigma(f(a)) &= \sigma(\phi_a(f)) \\ &= \sigma(f) \\ &= \text{Ran}(f) \\ &= f(\sigma(a)). \end{aligned}$$

- (4) We note that for any character  $h \in \Omega(A)$  and any continuous  $k \in C(\sigma(a))$ , that  $h(k(a)) = k(h(a))$ . This follows from the definition of the continuous functional calculus:

$$\begin{aligned} h(k(a)) &= h(\phi_a(k)) \\ &= \widehat{\phi_a(k)}(h) \\ &= \gamma_A(\phi_a(k))(h) \\ &= (\gamma_A \circ \phi_a)(k)(h) \\ &= T_{\hat{a}}(k)(h) \\ &= k \circ \hat{a}(h) \\ &= k(h(a)). \end{aligned}$$

We know that  $f(a)$  is normal, so  $f(a)$  admits the continuous functional calculus,  $\phi_{f(a)}: C(\sigma(f(a))) \rightarrow C^*(f(a), 1_B)$ . Thus, if  $g: \sigma(f(a)) \rightarrow \mathbb{C}$  is continuous, then  $g(f(a)) = \phi_{f(a)}(g)$  is a valid expression. Since  $\sigma(f(a)) = f(\sigma(a))$ , we have  $g \circ f: \sigma(a) \rightarrow \mathbb{C}$  is well-defined and continuous, defined by  $g \circ f(a) = \phi_a(g \circ f)$ .

Let  $h \in \Omega(A)$  be arbitrary. Using the above expression with  $k = g \circ f$ , we get

$$h(g \circ f(a)) = g \circ f(h(a)).$$

Replacing  $a$  with  $f(a)$  in the expression  $h(k(a)) = k(h(a))$ , and setting  $k = g \in C(\sigma(f(a)))$ , we get

$$\begin{aligned} h(g(f(a))) &= g(h(f(a))) \\ &= g(f(h(a))) \\ &= g \circ f(h(a)). \end{aligned}$$

Thus, we get  $h(g(f(a))) = h(g \circ f(a))$ . Since both  $g \circ f(a)$  and  $g(f(a))$  belong to  $A$ , and since characters separate points, we must have  $g \circ f(a) = g(f(a))$ .

- (5) Since normal elements are preserved under  $*$ -homomorphisms, we have that  $\pi(a)$  is normal in  $C$ . The  $C^*$ -subalgebra,  $A_1 = C^*(\pi(a), 1_C) \subseteq C$  is thus unital and commutative. We thus have  $\sigma(\pi(a)) \subseteq \sigma(a)$ . Dualizing the inclusion map  $\iota: \sigma(\pi(a)) \rightarrow \sigma(a)$ , we get the restriction map  $\rho: C(\sigma(a)) \rightarrow C(\sigma(\pi(a)))$ .

The unital  $*$ -homomorphism  $\pi: A \rightarrow A_1$  induces an adjoint map  $\pi^*: \Omega(A_1) \rightarrow \Omega(A)$ , given by  $\pi^*(h) = h \circ \pi$ . Since  $\pi^*$  consists of a character composed with a unital  $*$ -homomorphism, it is also a character. We see that  $\pi^*$  is continuous between these character spaces. If  $(h_\alpha)_\alpha \rightarrow h$  in  $\Omega(A_1)$ , then

$$\begin{aligned} (h_\alpha)_\alpha \xrightarrow{w^*} h &\Rightarrow (h_\alpha(z))_\alpha \rightarrow h(z) & z \in A_1 \\ &\Rightarrow (h_\alpha(\pi(x)))_\alpha \rightarrow h(\pi(x)) & x \in A \\ &\Rightarrow (h_\alpha \circ \pi(x))_\alpha \rightarrow h \circ \pi(x) \\ &\Rightarrow \pi^*(h_\alpha) \xrightarrow{w^*} \pi^*(h). \end{aligned}$$

Dualizing  $\pi^*: \Omega(A_1) \rightarrow \Omega(A)$ , we get the unital  $*$ -homomorphism  $T_{\pi^*}: C(\Omega(A)) \rightarrow C(\Omega(A_1))$ . The following diagram expresses the results we have so far.

$$\begin{array}{ccccc} C(\sigma(a)) & \longrightarrow & C(\Omega(A)) & \longrightarrow & A \\ \downarrow & & \downarrow & & \downarrow \\ C(\sigma(\pi(a))) & \longrightarrow & C(\Omega(A_1)) & \longrightarrow & A_1 \end{array}$$

Here, the top row expresses the continuous functional calculus on  $a$ , and the bottom row is the continuous functional calculus on  $\pi(a)$ . We claim that both squares in this diagram commute.

For  $f \in C(\sigma(a))$  and  $h \in \Omega(A_1)$ , we have

$$\begin{aligned} T_{\pi^*} \circ T_{\hat{a}}(f)(h) &= T_{\pi^*}(f \circ \hat{a})(h) \\ &= f \circ \hat{a} \circ \pi^*(h) \\ &= f \circ \hat{a} \circ (h \circ \pi) \\ &= f(h(\pi(a))) \\ &= f \circ \widehat{\pi(a)}(h) \\ &= T_{\widehat{\pi(a)}} \circ \rho(f)(h), \end{aligned}$$

so the left square commutes. For the right square, it is enough to verify that  $\gamma_{A_1} \circ \pi = T_{\pi^*} \circ \gamma_A$ . We have

$$\begin{aligned} \gamma_{A_1} \circ \pi &= G_{\pi^*} \circ \gamma_A \Leftrightarrow \gamma_{A_1} \circ \pi(a) = T_{\pi^*} \circ \gamma_A(a) \\ &\Leftrightarrow \widehat{\pi(a)} = \hat{a} \circ \pi^* \\ &\Leftrightarrow \widehat{\pi(a)}(h) = \hat{a} \circ \pi^*(h) \\ &\Leftrightarrow h(\pi(a)) = \hat{a}(h \circ \pi) \end{aligned}$$



$$\Leftrightarrow h(\pi(a)) = h(\pi(a)).$$

Since the above diagram commutes, we have  $\phi_{\pi(a)} \circ \rho = \pi \circ \phi_a$ .

If  $f \in C(\sigma(a))$ , then

$$\begin{aligned} f(\pi(a)) &= \phi_{\pi(a)} \circ \rho(f) \\ &= \pi \circ \phi_a(f) \\ &= \pi(f(a)). \end{aligned}$$

(6) We may uniformly approximate  $f$  by a sequence of functions  $p_n: \sigma(a) \rightarrow \mathbb{C}$ , where each  $p_n$  is of the form

$$p_n(z) = \sum_{k,l=1}^m c_{k,l} z^k \bar{z}^l,$$

with no constant term. For each  $n \geq 1$ , we see that  $p_n(a) \in C^*(a)$ , as each  $p_n$  contains no constant term. Using the result from (1), and the fact that each  $\phi_a$  is continuous, we get

$$\begin{aligned} \phi_a(f) &= \phi_a\left(\lim_{n \rightarrow \infty} p_n\right) \\ &= \lim_{n \rightarrow \infty} \phi_a(p_n) \\ &= \lim_{n \rightarrow \infty} p_n(a) \\ &\in C^*(a). \end{aligned}$$

□

When we want to carry out the continuous functional calculus in non-unital algebras, we need a slightly modified approach. If  $B$  is nonunital and  $a \in B$  is normal, then  $\phi_a: C(\sigma(a)) \rightarrow C^*(a, 1_{\bar{B}})$  is our continuous functional calculus. Note that  $0 \in \sigma(a)$ , so  $\phi_a(f) \in C^*(a)$  if  $f \in C(\sigma(a))$  with  $f(0) = 0$ .

**Corollary:** If  $B$  is a  $C^*$ -algebra, and  $a$  is normal, then there is an isometric  $*$ -isomorphism

$$\psi_a: C_0(\sigma(a) \setminus \{0\}) \rightarrow C^*(a)$$

that satisfies  $\psi_a(\iota) = a$ , where  $\iota: \sigma(a) \setminus \{0\} \rightarrow \mathbb{C}$  is inclusion.

*Proof.* If  $0 \notin \sigma(a)$ , then we set  $\psi_a = \phi_a$ , and apply the previous theorem.

Suppose  $0 \in \sigma(a)$ . We identify

$$C_0(\sigma(a) \setminus \{0\}) = \{f \in C(\sigma(a)) \mid f(0) = 0\}.$$

We restrict  $\psi_a = \phi_a|_{C_0(\sigma(a) \setminus \{0\})}$ , and apply the previous theorem. □

We can provide an alternative proof of a previous result.

**Proposition:** Let  $\pi: A \rightarrow B$  be a  $*$ -homomorphism between  $C^*$ -algebras.

- (1) If  $a \in A$  is normal and  $\pi$  is injective, then  $\sigma(\pi(a)) = \sigma(a)$ .
- (2) The map  $\pi$  is injective if and only if it is isometric.

*Proof.* We assume that  $A, B, \pi$  are unital.

(1) We already know that  $\sigma(\pi(a)) \subseteq \sigma(a)$ .

Suppose the inclusion is not strict. By Urysohn's lemma, we may find a nonzero  $f \in C(\sigma(a))$  such that  $f|_{\sigma(\pi(a))} = 0$ . If  $\phi_a: C(\sigma(a)) \rightarrow C^*(1, a)$  is the continuous functional calculus at  $a$ , then  $\phi_a(f) = f(a) \neq 0$ .

Since  $\pi$  is injective, we get  $0 \neq \pi(f(a)) = f(\pi(a)) = 0$ , which is a contradiction.

(2) It is clear that isometric maps are injective.

Suppose  $\pi$  is injective. Let  $x \in A$ , and set  $a = x^*x$ . Since  $a$  is normal,  $\sigma(a) = \sigma(\pi(a))$ , hence  $r(a) = r(\pi(a))$ . Thus, we have

$$\begin{aligned} \|\pi(a)\| &= r(\pi(a)) \\ &= r(a) \\ &= \|a\|. \end{aligned}$$

Thus, via the  $C^*$  identity, we get

$$\begin{aligned} \|x\|^2 &= \|x^*x\| \\ &= \|a\| \\ &= \|\pi(a)\| \\ &= \|\pi(x^*x)\| \\ &= \|\pi(x)^*\pi(x)\| \\ &= \|\pi(x)\|^2. \end{aligned}$$

□

We are interested in understanding the spectra of elements in  $C^*$ -algebras through the functional calculus.

**Proposition:** Let  $B$  be a  $C^*$ -algebra, and let  $a \in B$  be normal. The following hold:

- (1)  $a \in B_{s.a.}$  if and only if  $\sigma(a) \subseteq \mathbb{R}$ ;
- (2)  $a \in \mathcal{P}(B)$  if and only if  $\sigma(a) \subseteq \{0, 1\}$ ;
- (3) if  $B$  is unital, then  $a \in \mathcal{U}(B)$  if and only if  $\sigma(a) \subseteq \mathbb{T}$ .

*Proof.* If  $B$  is not unital, we unitize and consider the calculus  $\phi_a: C(\sigma(a)) \rightarrow C^*(a, 1_{\tilde{B}})$ , recalling that  $\sigma(a) = \sigma_{\tilde{B}}(a)$ . We assume  $B$  is unital, and let  $f = \text{id}_{\sigma(a)}$ .

(1) Since  $\phi_a$  is injective, self-adjoint, and since  $\phi_a(f) = a$ , we have

$$\begin{aligned} a^* = a &\Leftrightarrow \phi_a(f) = \phi_a(f)^* \\ &\Leftrightarrow \phi_a(f) = \phi_a(f^*) \\ &\Leftrightarrow f = f^* \\ &\Leftrightarrow f(\lambda) = \overline{f(\lambda)} \\ &\Leftrightarrow \lambda = \bar{\lambda} \\ &\Leftrightarrow \sigma(a) \subseteq \mathbb{R}. \end{aligned}$$

(2) Similarly,

$$a = a^2 \Leftrightarrow \phi_a(f) = \phi_a(f)^2$$

$$\begin{aligned}
&\Leftrightarrow \phi_a(f) = \phi_a(f^2) \\
&\Leftrightarrow f = f^2 \\
&\Leftrightarrow f(\lambda) = f(\lambda)^2 \\
&\Leftrightarrow \lambda = \lambda^2 \\
&\Leftrightarrow \sigma(a) \subseteq \{0, 1\}.
\end{aligned}$$

(3) Since  $\phi_a$  is unital, we have

$$\begin{aligned}
a^*a = 1_B &\Leftrightarrow \phi_a(f)^* \phi_a(f) = \phi_a(\mathbb{1}_{\sigma(a)}) \\
&\Leftrightarrow \phi_a(f^*f) = \phi_a(\mathbb{1}_{\sigma(a)}) \\
&\Leftrightarrow f^*f = \mathbb{1}_{\sigma(a)} \\
&\Leftrightarrow \overline{f(\lambda)}f(\lambda) = 1 \\
&\Leftrightarrow |\lambda|^2 = 1 \\
&\Leftrightarrow |\lambda| \in 1 \\
&\Leftrightarrow \sigma(a) \subseteq \mathbb{T}.
\end{aligned}$$

□

An important fact is that in any unital algebra, the inverse of an invertible element  $a \in GL(A)$  can be obtained using the functional calculus.

**Fact.** Let  $A$  be a unital  $C^*$ -algebra. Suppose  $a \in A$  is normal and invertible. Then,  $a^{-1} = g(a)$ , where  $g(z) = z^{-1}$ . Moreover,

$$\sigma(a^{-1}) = \{z^{-1} \mid z \in \sigma(a)\}.$$

*Proof.* Since  $a$  is invertible,  $0 \notin \sigma(a)$ , so  $g$  is defined and continuous on  $\sigma(a)$ . Then, where  $\iota: \sigma(a) \hookrightarrow \mathbb{C}$  is inclusion, we have

$$\begin{aligned}
\phi_a(g)a &= \phi_a(g)\phi_a(\iota) \\
&= \phi_a(g\iota) \\
&= \phi_a(\mathbb{1}_{\sigma(a)}) \\
&= 1_A.
\end{aligned}$$

Similarly,  $a\phi_a(g) = 1_A$ . Since inverses are unique, we have  $a^{-1} = \phi_a(g)$ . □

**Exercise:** Let  $A$  be a unital  $C^*$ -algebra, and suppose  $a \in A$  is normal and invertible. Prove that  $a^{-1} \in C^*(a)$ , and  $C^*(a) = C^*(a, 1_A)$ .

**Solution:** We only need to show the first statement, as the second statement follows directly.

Note that we have an isometric  $*$ -isomorphism  $\psi_a: C_0(\sigma(a) \setminus \{0\}) \rightarrow C^*(a)$ . Since  $\sigma(a) \setminus \{0\} = \sigma(a)$ , and  $\sigma(a)$  is compact, we must have an isometric  $*$ -isomorphism between  $C(\sigma(a))$  and  $C^*(a)$ .

Since  $0 \notin \sigma(a)$ , we have  $g(z) = z^{-1} \in C(\sigma(a))$ , so  $a^{-1} \in C^*(a)$ .

**Exercise:** Let  $T \in \mathcal{B}(\mathcal{H})$  be a normal operator that is not a scalar multiple of the identity. Prove that there are self-adjoint operators  $a, b \in C^*(T, I_{\mathcal{H}})$  such that  $ab = 0$ .

Another application of the continuous functional calculus is in the proof of the spectral theorem for compact normal operators.

**Theorem:** If  $T$  is a compact normal operator acting on an infinite-dimensional Hilbert space, there is a (possibly finite) sequence  $(\lambda_n)_n$  in  $c_0$  consisting of distinct nonzero scalars and a (possibly finite) sequence  $(P_n)_n$  of mutually orthogonal projections such that

$$T = \sum_{n \geq 1} \lambda_n P_n$$

is an operator norm-convergent sum.

*Proof.* We know that  $\sigma(T)$  is finite, or we have  $\|T\|_{\text{op}} = |\lambda_1| > |\lambda_2| > \dots \rightarrow 0 = \lambda_0$ .

For  $n \geq 1$ , we have shown that  $\lambda_n$  is isolated, so  $\delta_n = \mathbb{1}_{\{\lambda_n\}} \in C_0(\sigma(T) \setminus \{0\})$ . We have a uniformly convergent series

$$\iota = \sum_{n \geq 1} \lambda_n \delta_n$$

in  $C_0(\sigma(T) \setminus \{0\})$ , where  $\iota$  is inclusion. Applying the isometric  $*$ -isomorphism  $\psi_a: C_0(\sigma(T) \setminus \{0\}) \rightarrow C^*(T)$ , we get the operator norm-convergent sum

$$T = \sum_{n \geq 1} \lambda_n P_n,$$

where  $P_n = \psi_a(\delta_n) \in C^*(T) \subseteq \mathcal{K}(\mathcal{H})$ . Note that compact projections are finite-rank, and for  $n \neq m$ , we have

$$\begin{aligned} P_n P_m &= \psi_a(\delta_n) \psi_a(\delta_m) \\ &= \psi_a(\delta_n \delta_m) \\ &= \psi_a(0) \\ &= 0. \end{aligned}$$

□

We can also use the continuous functional calculus to determine isomorphism classes of universal  $C^*$ -algebras.

**Example.** We have looked at the  $C^*$ -algebra generated by a unitary,  $C^*(u)$ . With the continuous functional calculus, we can show that  $C^*(u)$  is  $*$ -isomorphic to any  $C^*$ -algebra generated by a unitary with full spectrum.

Let  $A$  be a unital  $C^*$ -algebra, and suppose  $w \in \mathcal{U}(A)$ , with  $\sigma(w) = \mathbb{T}$ .<sup>viii</sup> Let  $\phi_w: C(\mathbb{T}) \rightarrow C^*(w)$  and  $\phi_u: C(\sigma(u)) \rightarrow C^*(u)$  be the continuous functional calculi at  $w \in A$  and  $u \in C^*(u)$  respectively.

Since  $\sigma(u) \subseteq \mathbb{T}$ , we have the restriction unital  $*$ -homomorphism  $\rho: C(\mathbb{T}) \rightarrow C(\sigma(u))$ , given by  $\rho(f) = f|_{\sigma(u)}$ . Moreover, by the universal property of  $C^*(u)$ , we have a surjective  $*$ -homomorphism  $\psi: C^*(u) \rightarrow C^*(w)$ .

The unital  $*$ -homomorphism  $\varphi = \phi_u \circ \rho \circ \phi_w^{-1}: C^*(w) \rightarrow C^*(u)$  sends  $w \mapsto u$ . Thus,  $\varphi \circ \psi$  agrees on a dense  $*$ -subalgebra of  $C^*(u)$ , and by continuity, on  $C^*(u)$ , so  $\psi$  is injective. Thus, we have

$$\begin{aligned} C^*(u) &\cong C(\mathbb{T}) \\ &\cong C^*(V) \\ &\cong C^*(\mathbb{Z}), \end{aligned}$$

where  $V$  is the right bilateral shift on  $\ell_2(\mathbb{Z})$ .

<sup>viii</sup>We call these Haar unitaries.

## Ordering and Positive Elements

Recall that we say an element of a  $*$ -algebra  $a \in A$  is positive (in the algebraic sense) if  $a = x^*x$  for some  $x \in A$ . We denote  $A_+ \subseteq A_{s.a.}$  to be the set of positive elements as a subset of the set of all self-adjoint elements.

However, if  $A$  is a  $C^*$ -algebra, this set of positive elements goes further, and forms a closed generating cone in  $A_{s.a.}$  that induces an ordering on the self-adjoint elements in  $A$ . This will require some work.

Given a LCH space  $\Omega$ , we have looked at the closed cone of positive elements,  $C_0(\Omega)_+$  inside the  $\mathbb{R}$ -space of  $C_0(\Omega, \mathbb{R})$ . A function  $f \in C_0(\Omega)$  is positively-valued if and only if it is positive in the algebraic sense.

Similarly, the set of positive operators

$$\mathbb{B}(\mathcal{H})_+ = \{T \in \mathbb{B}(\mathcal{H})_{s.a.} \mid \langle T(\xi), \xi \rangle \geq 0, \forall \xi \in \mathcal{H}\}$$

forms a norm-closed cone in  $\mathbb{B}(\mathcal{H})_+$ , which induces an ordering on the self-adjoint elements of  $\mathbb{B}(\mathcal{H})_{s.a.}$ . Positivity in  $\mathbb{B}(\mathcal{H})_{s.a.}$  is described spatially, but it is not obvious that this definition agrees with the algebraic definition.

Any element in a  $*$ -algebra  $A$  can be written as  $a = h + ik$ , where  $h, k \in A_{s.a.}$ . If  $A$  is a normed  $*$ -algebra, then  $\|h\| \leq \|a\|$  and  $\|k\| \leq \|a\|$ . This is the Cartesian decomposition of  $a$ . We are aware that every real-valued continuous function can be written as the difference of two positive continuous functions. We start by generalizing to  $C^*$ -algebras.

**Proposition:** Let  $A$  be a  $C^*$ -algebra and let  $h \in A_{s.a.}$ . There exist unique positive elements  $p, q \in A_+$  such that

- (a)  $h = p - q$ ;
- (b)  $pq = 0$ ;
- (c)  $\sigma(p), \sigma(q) \subseteq [0, \infty)$ .

Moreover,  $\|h\| = \max(\|p\|, \|q\|)$ . We denote these elements  $h_+ = p$  and  $h_- = q$ .

*Proof.* Assume  $A$  is unital. Let  $\phi_h: C(\sigma(h)) \rightarrow C^*(h, 1_A)$  be the continuous functional calculus at  $h$ . Since  $\sigma(h) \subseteq \mathbb{R}$ , we consider the continuous functions

$$\begin{aligned} f(t) &= \max(t, 0) \\ g(t) &= \max(-t, 0). \end{aligned}$$

Note that  $fg = 0$ , and  $\text{id}_{\sigma(h)} = f - g$ . We set  $p = \phi_h(f)$  and  $q = \phi_h(g)$ . Since  $f$  and  $g$  are positive, their  $*$ -homomorphic images are positive as well. Moreover,

$$\begin{aligned} \sigma(p) &= \sigma(\phi_h(f)) \\ &= \phi_h(\sigma(h)) \\ &\subseteq [0, \infty), \end{aligned}$$

and similarly for  $\sigma(q)$ . We also have

$$\begin{aligned} pq &= \phi_h(f)\phi_h(g) \\ &= \phi_h(fg) \\ &= \phi_h(0) \\ &= 0, \end{aligned}$$

and

$$\begin{aligned}
 h &= \phi_h(\text{id}_{\sigma(h)}) \\
 &= \phi_h(f - g) \\
 &= \phi_h(f) - \phi_h(g) \\
 &= p - q.
 \end{aligned}$$

Since  $\phi_h$  is isometric, we also have

$$\begin{aligned}
 \|h\| &= \|\phi_h(\text{id}_{\sigma(h)})\| \\
 &= \|\text{id}_{\sigma(h)}\| \\
 &= \max(\|f\|_u, \|g\|_u) \\
 &= \max(\|\phi_h(f)\|, \|\phi_h(g)\|) \\
 &= \max(\|p\|, \|q\|).
 \end{aligned}$$

We will now show uniqueness. Suppose  $x, y \in A_+$  are such that  $h = x - y$ ,  $xy = 0$ , and  $\sigma(x), \sigma(y) \subseteq [0, \infty)$ . By induction, we have  $h^n = x^n + (-y)^n$  for all  $n \geq 1$ , so  $p(h) = p(x) + p(-y)$  for all polynomials  $p$  without constant terms.

Since  $f(0) = 0$ , there is a sequence of polynomials converging uniformly to  $f$  on the compact set  $K = \sigma(h) \cup \sigma(x) \cup \sigma(-y)$ . Applying the functional calculus at  $h, x$ , and  $y$ , we get

$$\begin{aligned}
 p &= f(h) \\
 &= \lim_{n \rightarrow \infty} p_n(h) \\
 &= \lim_{n \rightarrow \infty} (p_n(x) + p_n(-y)) \\
 &= f(x) + f(-y).
 \end{aligned}$$

Since  $\sigma(x) \subseteq [0, \infty)$ , and  $f(t) = t$  on  $[0, \infty)$ , we must have  $f(x) = x$ . Also, since  $\sigma(-y) = -\sigma(y) \subseteq (-\infty, 0]$ , and  $f(t) = 0$  on  $(-\infty, 0]$ , we must have  $f(-y) = 0$ . Thus,  $p = x$ , and  $q = p - h = y$ .

If  $A$  does not have a unit, we consider the functional calculus  $\phi_h: C(\sigma(h)) \rightarrow C^*(h, 1_{\tilde{A}})$ . Since  $0 \in \sigma(h)$  and  $f(0) = g(0) = 0$ , we must have  $p, q \in C^*(h) \subseteq A$ .  $\square$

**Lemma:** Let  $A$  be a  $C^*$ -algebra with  $y \in A_{s.a.}$ . Then,  $\sigma(y^2) \subseteq [0, \infty)$ .

*Proof.* We assume  $A$  is unital. By spectral mapping, we have  $\sigma(y^2) = (\sigma(y))^2 \subseteq [0, \infty)$ , as  $\sigma(y) \subseteq \mathbb{R}$ .  $\square$

**Lemma:** Let  $A$  be a  $C^*$ -algebra.

- (1) If  $\sigma(a) \subseteq [0, \infty)$ , and  $t \geq 0$ , then  $\sigma(ta) \subseteq [0, \infty)$ .
- (2) Assume  $A$  is unital, let  $a \in A_{s.a.}$ , and let  $t \geq 0$  be such that  $\|a\| \leq t$ . Then,  $\sigma(a) \subseteq [0, \infty)$  if and only if  $\|t1_A - a\| \leq t$ .
- (3) Let  $a, b \in A_{s.a.}$  with  $\sigma(a), \sigma(b) \subseteq [0, \infty)$ . Then,  $\sigma(a + b) \subseteq [0, \infty)$ .

*Proof.*

- (1) By spectral mapping, we have  $\sigma(ta) = t\sigma(a)$ .
- (2) Since  $a = a^*$ , and  $\|a\| \leq t$ , we have  $\sigma(a) \subseteq [-t, t]$ . Let  $\phi_a: C(\sigma(a)) \rightarrow C^*(a, 1_A)$  be the continuous functional calculus at  $a$ . Letting  $\iota$  be inclusion, we see that

$$\begin{aligned}
 \|t1_A - a\| &= \|\phi_a(t1_{\sigma(a)} - \iota)\| \\
 &= \|t1_{\sigma(a)} - \iota\|.
 \end{aligned}$$

This is only less than or equal to  $t$  if, for all  $\lambda \in \sigma(a)$ ,  $|t - \lambda| \leq t$ , or  $\sigma(a) \subseteq [0, 2t]$ .

(3) Assuming  $A$  is unital, we see that  $\| \|a\|1_A - a \| \leq \|a\|$  and  $\| \|b\|1_A - b \| \leq \|b\|$ . Thus,

$$\begin{aligned} \|(\|a\| + \|b\|)1_A - (a + b)\| &\leq \| \|a\|1_A - a \| + \| \|b\|1_A - b \| \\ &\leq \|a\| + \|b\|. \end{aligned}$$

Taking  $t = \|a\| + \|b\|$ , we see that  $\|a + b\| \leq t$ , so  $\sigma(a + b) \subseteq [0, \infty)$ .

□

**Theorem:** Let  $A$  be a  $C^*$ -algebra with  $a \in A$ . The following are equivalent:

- (a)  $a \in A_{s.a.}$  and  $\sigma(a) \subseteq [0, \infty)$ ;
- (b)  $a$  is normal and  $\sigma(a) \subseteq [0, \infty)$ ;
- (c)  $a \in A_+$ .

*Proof.* It is obvious that (i) implies (ii).

To see (ii) implies (iii), we assume  $A$  is unital. Let  $\phi_a: C(\sigma(a)) \rightarrow C^*(a, 1_A)$  be the continuous functional calculus at  $a$ . Since  $\sigma(a) \subseteq [0, \infty)$ , the function  $f(t) = \sqrt{t}$  is well-defined, self-adjoint, and continuous on  $\sigma(a)$ . Set  $b = \phi_a(f)$ . Then, letting  $\iota$  be inclusion, we have

$$\begin{aligned} a &= \phi_a(\iota) \\ &= \phi_a(f^2) \\ &= \phi_a(f)\phi_a(f) \\ &= b^2. \end{aligned}$$

We see that  $b$  is self-adjoint, as it is the  $*$ -homomorphic image of a self-adjoint element,<sup>ix</sup> we have  $a = b^*b \in A_+$ .

If  $A$  is nonunital, let  $\phi_a: C(\sigma(a)) \rightarrow C^*(a, 1_{\tilde{A}})$  be the continuous functional calculus at  $a$ . Since  $f(0) = 0$ , we have  $b = \phi_a(f) \in C^*(a)$ .

To see (iii) implies (i), assume  $a = b^*b$  for some  $b \in A$ . It is the case that  $a$  is self-adjoint, so we need to show that  $\sigma(a) \subseteq [0, \infty)$ . We may assume  $A$  is unital. Write  $a = p - q$  to be the decomposition of  $a$  where  $\sigma(p), \sigma(q) \subseteq [0, \infty)$  and  $p, q \in A_+$ , and set  $c = bq$ .

We have

$$\begin{aligned} c^*c &= (bq)^*(bq) \\ &= qb^*bq \\ &= qa q \\ &= q(p - q)q \\ &= -q^3, \end{aligned}$$

as  $pq = qp = 0$ . Thus, by spectral mapping, we have

$$\begin{aligned} \sigma(c^*c) &= \sigma(-q^3) \\ &= -\sigma(q)^3 \\ &\subseteq (-\infty, 0]. \end{aligned}$$

---

<sup>ix</sup>The self-adjoint element is in  $C(\sigma(a))$ , and the  $*$ -homomorphism is the continuous functional calculus.

Since  $0 \in \sigma(c^*c)$ , we see that  $\sigma(cc^*) \subseteq (-\infty, 0]$ , and by spectral mapping,  $\sigma(-cc^*) \subseteq [0, \infty)$ .

On the other hand, we may write the cartesian decomposition  $c = h + ik$ , where  $h, k \in A_{s.a.}$ , and find

$$\begin{aligned} c^*c + cc^* &= (h + ik)^*(h + ik) + (h + ik)(h + ik)^* \\ &= 2(h^2 + k^2), \end{aligned}$$

so  $c^*c = 2h^2 + 2k^2 + (-cc^*)$ . Since  $\sigma(2h^2), \sigma(2k^2), \sigma(-cc^*) \subseteq [0, \infty)$ , we have  $\sigma(c^*c) \subseteq [0, \infty)$ . Thus,  $\sigma(c^*c) = \{0\}$ . As  $c$  is normal, we have

$$\begin{aligned} \|c\|^2 &= \|c^*c\| \\ &= r(c^*c) \\ &= 0, \end{aligned}$$

so  $-q^3 = c^*c = 0$ . Applying the continuous functional calculus with  $f = -t^{1/3}$ , we have  $q = 0$ , meaning  $a = p$ . Thus,  $\sigma(a) \subseteq [0, \infty)$ .  $\square$

We will reconcile some definitions for bounded operators on Hilbert spaces now.

**Proposition:** Let  $\mathcal{H}$  be a Hilbert space, and let  $T \in \mathcal{B}(\mathcal{H})$ . The following are equivalent:

- (i)  $\langle T(\xi), \xi \rangle \geq 0$  for all  $\xi \in \mathcal{H}$ ;
- (ii)  $T = T^*$  and  $\sigma(T) \subseteq [0, \infty)$ ;
- (iii)  $T = S^*S$  for some  $S \in \mathcal{B}(\mathcal{H})$ .

*Proof.* We only need to prove (i) implies (ii), as we have proven the equivalence between (ii) and (iii).

If  $\langle T(\xi), \xi \rangle \geq 0$  for all  $\xi \in \mathcal{H}$ , then  $T = T^*$ , as  $\langle T(\xi), \xi \rangle \in \mathbb{R}$ , so  $\sigma(T) \subseteq \mathbb{R}$ . If  $\lambda < 0$ , then using the fact that  $T$  is self-adjoint, we have

$$\begin{aligned} \|(T - \lambda I)(\xi)\|^2 &= \langle T(\xi) - \lambda\xi, T(\xi) - \lambda\xi \rangle \\ &= \|T(\xi)\|^2 - 2\lambda\langle T(\xi), \xi \rangle + \lambda^2\|\xi\|^2 \\ &\geq \lambda^2\|\xi\|^2, \end{aligned}$$

so  $T - \lambda I$  is bounded below, hence invertible, so  $\lambda \in \rho(T)$ . Thus,  $\sigma(T) \subseteq [0, \infty)$ .  $\square$

**Corollary:** Let  $A$  be a  $C^*$ -algebra. The collection of positive elements,  $A_+$ , is a generating norm-closed cone in  $A_{s.a.}$ .

*Proof.* If  $a, -a \in A_+$ , then  $a = a^*$  and  $\sigma(a) \subseteq [0, \infty)$  and  $\sigma(-a) \subseteq (-\infty, 0]$ . Thus,  $\sigma(a) = -\sigma(-a) \subseteq (-\infty, 0]$ , so  $\sigma(a) = \{0\}$ , so  $\|a\| = r(a) = 0$ , so  $a = 0$ . Thus,  $A_+ \subseteq A_{s.a.}$  is a cone.

The fact that  $A_+$  generates  $A_{s.a.}$  follows from the fact that any element in  $A_{s.a.}$  can be written as a difference of two positive elements.

We show that  $A_+$  is closed. Let  $A$  be unital, and let  $(a_n)_n \rightarrow a \in A$  be a sequence in  $A_+$ . Then,

$$\begin{aligned} a &= \lim_{n \rightarrow \infty} (a_n)_n \\ &= \lim_{n \rightarrow \infty} (a_n^*)_n \\ &= \left( \lim_{n \rightarrow \infty} (a_n)_n \right)^* \\ &= a^*, \end{aligned}$$



so  $a$  is self-adjoint.

Let  $C > 0$  be such that  $\|a_n\| \leq C$  for all  $n$ . It follows that  $\|a\| \leq C$ , so we have

$$\begin{aligned} \|C1_A - a\| &= \lim_{n \rightarrow \infty} \|C1_A - a_n\| \\ &\leq C \end{aligned}$$

so  $a \in A_+$ .

If  $A$  lacks a unit, then  $A \subseteq \tilde{A}$  is closed and  $A_+ = A \cap \tilde{A}_+$ . □

Now, we shall define an ordering on  $A_{s.a.}$  with the cone of positive elements  $A_+$ .

**Proposition:** Let  $A$  be a  $C^*$ -algebra. There is an ordering on  $A_{s.a.}$  defined by  $x \leq y$  if  $y - x \in A_+$ . The ordering satisfies the following:

- (1) if  $x \leq y$ , then for any  $z \in A$ ,  $z^*xz \leq z^*yz$ ;
- (2) if  $A$  is unital,  $x \in A_{s.a.}$ , and  $s, t \in \mathbb{R}$ , then  $s1_A \leq x \leq t1_A$  if and only if  $\sigma(x) \subseteq [s, t]$ ;
- (3) if  $A$  is unital and  $x \in A_{s.a.}$ , then  $-\|x\|1_A \leq x \leq \|x\|1_A$ ;
- (4) for  $a \geq 0$ , we have  $a \leq 1_A$  if and only if  $\|a\| \leq 1$ ;
- (5) if  $0 \leq a \leq b$ , then  $\|a\| \leq \|b\|$ ;
- (6) if  $a, b \in A$ , then  $0 \leq b^*a^*ab \leq \|a\|^2b^*b$ ;
- (7) any  $*$ -homomorphism between  $C^*$ -algebras is positive (hence order preserving).

*Proof.* Since  $A_+$  is a cone, we know that the ordering  $x \leq y$  if  $y - x \in A_+$  is indeed an ordering.

- (1) If  $x \leq y$ , then  $y - x = a^*a$  for some  $a \in A$ . Consequently,

$$\begin{aligned} z^*yz - z^*xz &= z^*(y - x)z \\ &= z^*a^*az \\ &= (az)^*az \\ &\in A_+. \end{aligned}$$

- (2) Note that  $s1_A - x$  is self-adjoint. Thus, we have

$$\begin{aligned} s1_A \leq x &\Leftrightarrow x - s1_A \in A_+ \\ &\Leftrightarrow \sigma(x - s1_A) \subseteq [0, \infty) \\ &\Leftrightarrow \sigma(x) - s \subseteq [0, \infty) \\ &\Leftrightarrow \sigma(x) \subseteq [s, \infty). \end{aligned}$$

Similarly, we have  $x \leq t1_A$  if and only if  $\sigma(x) \subseteq (-\infty, t]$ . Thus,  $\sigma(x) \subseteq [s, t]$ .

- (3) This follows from the fact that  $\sigma(x) \subseteq [-\|x\|, \|x\|]$ . Since  $\|x\|1_A - x$  is self-adjoint, and  $\sigma(x) \subseteq [-\|x\|, \|x\|]$ , we must have

$$\begin{aligned} \sigma(\|x\|1_A - x) &= \|x\| - \sigma(x) \\ &\subseteq [0, 2\|x\|] \\ &\subseteq [0, \infty), \end{aligned}$$

so  $\|x\|1_A - x \in A_+$ , and  $x \leq \|x\|1_A$ . Similarly,  $-\|x\|1_A \leq x$ .

- (4) Taking  $s = 0$  and  $t = 1$  in (2), we get  $0 \leq a \leq 1$  if and only if  $\sigma(a) \subseteq [0, 1]$  if and only if  $r(a) = 1$  if and only if  $\|a\| = 1$ .
- (5) We have  $0 \leq a \leq b \leq \|b\|1_A$ , so  $\sigma(a) \subseteq [0, \|b\|]$ . Since  $a$  is normal,  $\|a\| \in \sigma(a)$ , so  $\|a\| \leq \|b\|$ .
- (6) Assume  $A$  is unital. The  $C^*$ -identity gives  $a^*a \leq \|a\|^2 1_A$ . Applying conjugation, we get  $b^*a^*ab \leq \|a\|^2 b^*b$ . If  $A$  is nonunital, we consider these as elements of  $\tilde{A}$ , replacing  $1_A$  with  $1_{\tilde{A}}$ .
- (7) If  $a = x^*x \in A_+$ , then  $\phi(a) = \phi(x^*x) = \phi(x)^*\phi(x) \in B_+$ . If  $x \leq y$  in  $A_{s.a.}$ , then  $y - x \in A_+$ , so

$$\begin{aligned}\phi(y) - \phi(x) &= \phi(y - x) \\ &\in B_+, \end{aligned}$$

so  $\phi(x) \leq \phi(y)$  in  $B_{s.a.}$ .

□

Just as the square root of a positive number  $t$  is the unique positive number  $s$  such that  $s^2 = t$ , we have a similar concept with  $C^*$ -algebras.

**Proposition:** Let  $A$  be a  $C^*$ -algebra. If  $x \in A_+$ , there is a unique  $y \in A_+$  such that  $y^2 = x$ . Moreover,  $y \in C^*(x)$ .

*Proof.* We start by assuming  $A$  is unital. Consider the continuous functional calculus  $\phi_x: C(\sigma(x)) \rightarrow C^*(x, 1_A)$ . Since  $\sigma(x) \subseteq [0, \infty)$ , we know the function  $g(t) = \sqrt{t}$  is continuous on  $\sigma(x)$ . Set  $y = \phi_x(g)$ . Since  $g$  is positive in  $C(\sigma(x))$ ,  $y$  is positive in  $A$ , and

$$\begin{aligned}y^2 &= \phi_x(g)^2 \\ &= \phi_x(g^2) \\ &= \phi_x(\iota) \\ &= x. \end{aligned}$$

This shows existence. Note that  $g(0) = 0$ , so  $y \in C^*(x)$ .

We will now show uniqueness. Suppose  $a \in A_+$  is such that  $a^2 = x$ . The continuous functional calculus at  $a$  gives  $f(a) = x$ , where  $f(t) = t^2$ . Thus, we get

$$\begin{aligned}y &= g(x) \\ &= g(f(a)) \\ &= g \circ f(a) \\ &= \iota(a) \\ &= a. \end{aligned}$$

If  $A$  is nonunital, then we consider  $x \in \tilde{A}_+$ , and use the above argument to produce  $y \in \tilde{A}_+$  with  $y^2 = x$ . However, since  $y \in C^*(x)$ , we have  $y \in A_+$ , and since  $A \cap \tilde{A}_+ = A_+$ , we get uniqueness. □

**Definition.** Let  $A$  be a  $C^*$ -algebra.

- (1) If  $x \in A_+$ , the unique positive element  $y \in A_+$  such that  $y^2 = x$  is called the positive square root of  $x$ , denoted  $y = x^{1/2}$ .
- (2) The absolute value of  $a \in A$  is given by  $|a| = (a^*a)^{1/2}$ .

**Exercise:** Let  $A$  be a  $C^*$ -algebra.

- (a) Prove that  $|a| \in C^*(a)$ .
- (b) Let  $h \in A_{s.a.}$  have decomposition  $h = h_+ - h_-$ . Prove that  $|h| = h_+ + h_-$ .

(c) Let  $a \in A$  be normal. Prove that  $|a| = f(a)$ , where  $f(z) = |z| \in C(\sigma(a))$ .

**Solution:**

(a) It is obvious that  $a^*a \in C^*(a)$ , and since the square root is an element of  $C^*(a)$ , we have  $|a| \in C^*(a)$ .

(b) Note that

$$\begin{aligned} |h| &= (h^*h)^{1/2} \\ &= \left( h_+^2 - h_+h_- - h_-h_+ + h_-^2 \right)^{1/2} \\ &= \left( h_+^2 + h_-^2 \right)^{1/2} \\ &= h_+ + h_-. \end{aligned}$$

(c) We have  $f(z) = |z| = (\bar{z}z)^{1/2}$ , so, using the continuous functional calculus, we have

$$\begin{aligned} f(a) &= (a^*a)^{1/2} \\ &= |a|. \end{aligned}$$

**Exercise:** Let  $\mathcal{H}$  be a Hilbert space, and suppose  $x, y \in \mathcal{H}$ . Prove that the rank one operator  $\theta_{x,y}$  has absolute value

$$|\theta_{x,y}| = \frac{\|x\|}{\|y\|} \theta_{y,y}.$$

**Solution:** We have

$$\theta_{x,y}^* \theta_{x,y} = \theta_{y,x} \theta_{x,y}.$$

**Exercise:** If  $A$  is a unital  $C^*$ -algebra, and  $a \in GL(A) \cap A_+$ , prove that  $a^{-1} \in A_+$ .

**Solution:** Note that we have  $0 \leq a \leq \|a\|$  as  $a$  is positive. However, since  $a$  is also invertible, there is some  $t > 0$  such that  $\sigma(a) \subseteq [t, \|a\|]$ . Applying the continuous functional calculus on  $g(t) = 1/t$ , we get  $\sigma(a^{-1}) \subseteq [\|a\|^{-1}, t^{-1}]$ , so  $a^{-1} \in A_+$ .

**Exercise:** Let  $A$  be a unital  $C^*$ -algebra, and suppose  $a \in GL(A) \cap A_+$ . Prove that  $a^{1/2} \in GL(A)$  and  $(a^{1/2})^{-1} = (a^{-1})^{1/2} = g(a)$ , where  $g(t) = \frac{1}{\sqrt{t}}$ .

**Solution:** Note that since  $a \in GL(A) \cap A_+$ , there is some  $t > 0$  such that  $\sigma(a) \subseteq [t, \|a\|]$ . Thus,  $\sigma(a^{1/2}) \subseteq [t^{1/2}, \|a\|^{1/2}]$ , so  $a^{1/2} \in GL(A)$ .

We apply the continuous functional calculus with  $g(t) = t^{-1/2}$  to yield

$$(a^{-1})^{1/2} = (a^{1/2})^{-1}.$$

**Lemma:** Let  $A$  be a unital  $C^*$ -algebra.

(1) If  $a \in A_{s.a.}$  is such that  $a + \varepsilon 1_A \geq 0$  for all  $\varepsilon > 0$ , then  $a \geq 0$ .

(2) If  $x, y \in A_{s.a.}$  are such that  $x \leq y + \varepsilon 1_A$  for all  $\varepsilon > 0$ , then  $x \leq y$ .

*Proof.* We will prove (1). Set  $a = y - x$ , and suppose there is a  $t \in \sigma(a)$  with  $t < 0$ . Set  $\varepsilon = -t/2 > 0$ . By spectral mapping, we have

$$\begin{aligned} t/2 &= t + \varepsilon \\ &\in \sigma(a) + \varepsilon \\ &= \sigma(a + \varepsilon 1_A) \\ &\subseteq [0, \infty), \end{aligned}$$

which is a contradiction. Thus,  $\sigma(a) \subseteq [0, \infty)$ , hence  $y - x \geq 0$ . □

**Proposition:** Let  $A$  be a  $C^*$ -algebra.

- (1) If  $a, b \in A_+$  with  $ab = ba$ , then  $ab \in A_+$ .
- (2) If  $a, b \in A_+$  with  $a \leq b$  and  $ab = ba$ , then  $a^2 \leq b^2$ .
- (3) Let  $A$  be unital. If  $a \in GL(A)$  with  $a \geq 1_A$ , then  $0 \leq a^{-1} \leq 1_A$ .
- (4) Let  $A$  be unital, and suppose  $0 \leq a \leq b$  with  $a \in GL(A)$ . Then,  $b$  is invertible, and  $0 \leq b^{-1} \leq a^{-1}$ .
- (5) If  $0 \leq a \leq b$ , then  $0 \leq a^{1/2} \leq b^{1/2}$ .
- (6) If  $a \in A_{s.a.}$ , then  $-|a| \leq a \leq |a|$ .

*Proof.*

- (1) Since  $a = a^*$ , we know that  $C^*(a)$  is the norm-closure of  $\{p(a) \mid p \in \mathbb{C}[x]\}$ . By continuity of multiplication,  $b$  commutes with every member of  $C^*(a)$ , including  $a^{1/2}$ . Finally, we have

$$\begin{aligned} ab &= a^{1/2} a^{1/2} b \\ &= a^{1/2} b a^{1/2} \\ &\geq 0. \end{aligned}$$

- (2) Since  $a$  and  $b$  commute, so do the positive elements  $b - a$  and  $b + a$ . Thus, we have

$$\begin{aligned} 0 &\leq (b - a)(b + a) \\ &= b^2 - ba + ab - a^2 \\ &= b^2 - a^2. \end{aligned}$$

- (3) Suppose  $a = a^* \geq 1$ . We know that  $\sigma(a)$  is compact and a subset of  $[1, \|a\|]$ . It follows that  $a$  is invertible and  $a^{-1}$  is also self-adjoint. Since  $a^{-1} = g(a)$ , where  $g(t) = 1/t$  is continuous on  $\sigma(a)$ . Thus, we find

$$\begin{aligned} \sigma(a^{-1}) &= \{t^{-1} \mid t \in \sigma(a)\} \\ &\subseteq [\|a\|^{-1}, 1]. \end{aligned}$$

Thus, we find  $0 \leq a^{-1} \leq 1$ .

- (4) Since  $a$  is invertible,  $0 \notin \sigma(a)$ , so  $\sigma(a) \subseteq [t, \|a\|]$  for some  $t > 0$ . Thus,  $t1_A \leq a \leq \|a\|1_A$ , so  $\sigma(b) \subseteq [t, \|b\|]$ , so  $0 \notin \sigma(b)$ , so  $b \in GL(A)$ .

We know that  $a^{-1}$  and  $b^{-1}$  are positive. Set  $C = (a^{-1})^{1/2} = (a^{1/2})^{-1}$ . Note that  $c$  commutes with  $a$ . Thus, we get

$$\begin{aligned} 1_A &= a^{-1}a \\ &= c^2a \\ &= cac \\ &\leq cbc, \end{aligned}$$

so we get

$$\begin{aligned} a^{1/2}b^{-1}a^{1/2} &= (cbc)^{-1} \\ &\leq 1_A. \end{aligned}$$

Multiplying both sides by  $c$  preserves the inequality, so we get

$$\begin{aligned} b^{-1} &= c \left( a^{1/2} b^{-1} a^{1/2} \right) \\ &\leq c^2 \\ &= a^{-1}. \end{aligned}$$

(5) Let  $x, y \geq 0$  be such that  $x^2 \leq y^2$ . We will show that  $x \leq y$ . Set  $x = a^{1/2}$  and  $y = b^{1/2}$ .

We start by assuming  $y$  is invertible. In that case,  $y^{-1}$  is positive, so we get

$$\begin{aligned} 0 &\leq y^{-1} x^2 y^{-1} \\ &\leq y^{-1} y^2 y^{-1} \\ &= 1, \end{aligned}$$

so  $\|y^{-1} x^2 y^{-1}\| \leq 1$ . By the  $C^*$ -identity, we have

$$\begin{aligned} \|xy^{-1}\|^2 &= \|(xy^{-1})^* xy^{-1}\| \\ &= \|y^{-1} x^2 y^{-1}\| \\ &\leq 1, \end{aligned}$$

so  $\|xy^{-1}\| \leq 1$ . Since  $xy^{-1}$  and  $y^{-1/2}xy^{-1/2}$  are similar elements, we have  $\|y^{-1/2}xy^{-1/2}\| \leq \|xy^{-1}\| \leq 1$ , so we have  $0 \leq y^{-1/2}xy^{-1/2} \leq 1$ . Thus,  $x \leq y$ .

For the general case, if we fix  $\varepsilon > 0$  and set  $y_\varepsilon = y + \varepsilon 1_A$ , then  $y_\varepsilon$  is invertible and  $y^2 \leq y_\varepsilon^2$ , so  $x^2 \leq y_\varepsilon^2$ , so  $x \leq y_\varepsilon$ . Since  $\varepsilon > 0$  is arbitrary, we have that  $x \leq y$ .

(6) Write  $a = a_+ - a_-$ , and note that  $|a| = a_+ + a_-$ . Thus,  $|a| - a = 2a_- \geq 0$ , so  $|a| \geq a$ . Similarly,  $a \geq -|a|$ .

□

**Proposition:** Let  $A$  be a  $C^*$ -algebra.

- (1) The square root map  $A_+ \rightarrow A_+$ ,  $a \mapsto a^{1/2}$ , is continuous.
- (2) The absolute value map,  $A \rightarrow A_+$ ,  $a \mapsto |a|$ , is continuous.

**Solution:**

- (1) Let  $(a_n)_n \rightarrow a$  in  $A_+$ , and let  $C > 0$  be such that  $\|a\| \leq C$  and  $\|a_n\| \leq C$  for all  $n \geq 1$ . By Weierstrass's approximation theorem, there is a sequence of polynomials  $(p_k)_k$  converging uniformly to  $f(t) = \sqrt{t}$  on  $[0, C]$ . Using the continuous functional calculus, we get

$$\begin{aligned} \|a^{1/2} - a_n^{1/2}\| &= \|\phi_a(f) - \phi_{a_n}(f)\| \\ &\leq \|\phi_a(f) - \phi_a(p_k)\| + \|\phi_a(p_k) - \phi_{a_n}(p_k)\| + \|\phi_{a_n}(p_k) - \phi_{a_n}(f)\| \\ &= \|\phi_a(f - p_k)\| + \|\phi_a(p_k) - \phi_{a_n}(p_k)\| + \|\phi_{a_n}(p_k - f)\| \\ &= 2\|f - p_k\| + \|p_k(a) - p_k(a_n)\|, \end{aligned}$$

which holds for all  $n, k \in \mathbb{N}$ . Given  $\varepsilon > 0$ , find  $K$  such that  $\|f - p_K\| \leq \varepsilon/3$ . Since  $(a_n)_n \rightarrow a$ , we have  $(p_K(a_n))_n \rightarrow p_K(a)$ . Thus, there is  $N$  such that  $\|p_K(a) - p_K(a_n)\| < \varepsilon/3$  for all  $n \geq N$ . Thus, for  $n \geq N$ , we get  $\|a^{1/2} - a_n^{1/2}\| < \varepsilon$  for  $n \geq N$ .

- (2) If  $(a_n)_n \rightarrow a$ , then  $(a_n^* a_n) \rightarrow a^* a$ , so

$$\begin{aligned} (|a_n|)_n &= \left( (a_n^* a_n)^{1/2} \right)_n \\ &\rightarrow (a^* a)^{1/2} \\ &= |a|. \end{aligned}$$

## Approximate Identities

Currently, whenever we have been faced with a nonunital  $C^*$ -algebra, we have chosen to unitize. However, it is often useful to work in the given algebra, or to use ideals and take quotients. For this purpose, we have to work with an approximate unit (or approximate identity), which is an increasing sequence or net of positive contractions that approximate a multiplicative unit.

**Definition.** Let  $A$  be a Banach algebra (unital or not). An approximate identity (or approximate unit) for  $A$  is a net  $(e_\alpha)_\alpha$  in  $A$  such that

- for all  $\alpha$ ,  $\|e_\alpha\| \leq 1$ ;
- for all  $x \in A$ ,  $\|x - xe_\alpha\| \rightarrow 0$  and  $\|x - e_\alpha x\| \rightarrow 0$ .

If  $A$  is a  $C^*$ -algebra, we also require

- $e_\alpha \geq 0$  for all  $\alpha$ ;
- $(e_\alpha)_\alpha$  is increasing — if  $\alpha \leq \beta$ , then  $e_\alpha \leq e_\beta$ .

**Example.** Consider  $A = C_0(\mathbb{R})$ , which is nonunital. For each  $n \geq 1$ , we find a continuous function  $f_n \in C_0(\mathbb{R})$  such that  $f$  is 1 on  $[-n, n]$ , 0 on  $[-n-1, n+1]$ , and linear elsewhere. If  $g \in C_0(\mathbb{R})$  and  $\varepsilon > 0$ , there is  $m \geq 1$  with  $|g(t)| < \varepsilon$  outside  $[-m, m]$ . For  $n \geq m$ , we see that  $\|g - gf_n\| < \varepsilon$ , so  $(f_n)_n$  is an approximate unit for  $C_0(\mathbb{R})$ .

**Example.** Consider  $\mathbb{K}(\ell_2)$ . For each  $n \geq 1$ , we consider

$$P_n = \sum_{k=1}^n \theta_{e_k, e_k},$$

which is the orthogonal projection onto  $\text{span}(\{e_1, \dots, e_n\})$ . The sequence  $(P_n)_n$  is increasing, and each  $P_n$  is positive with norm 1. We will show that  $\|T - TP_n\|_{\text{op}} \rightarrow 0$  for any  $T \in \mathbb{K}(\ell_2)$ .

Let  $\varepsilon > 0$ . Since  $T$  is compact and  $(e_k)_k \rightarrow 0$  weakly, we know that  $\|T(e_k)\| \rightarrow 0$ . Find  $N$  such that  $\|T(e_k)\| < \varepsilon$  for all  $k \geq N$ . For any  $\xi = \sum_{k \geq 1} \langle \xi, e_k \rangle e_k$ , in  $B_{\mathcal{H}}$ , we use Bessel's inequality to find, provided  $n \geq N$ , that

$$\begin{aligned} \|(T - TP_n)(\xi)\| &= \left\| \sum_{k \geq 1} \langle \xi, e_k \rangle T(e_k) - \sum_{k \geq 1} TP_n(e_k) \right\| \\ &= \left\| \sum_{k \geq 1} \langle \xi, e_k \rangle T(e_k) - \sum_{k=1}^n \langle \xi, e_k \rangle T(e_k) \right\| \\ &= \left\| \sum_{k > n} \langle \xi, e_k \rangle T(e_k) \right\| \\ &\leq \sum_{k > n} |\langle \xi, e_k \rangle| \|T(e_k)\| \\ &\leq \varepsilon \left( \sum_{k > n} |\langle \xi, e_k \rangle| \right) \\ &\leq \varepsilon. \end{aligned}$$

Since  $\xi$  was arbitrary, it follows that  $\|T - TP_n\|_{\text{op}} \leq \varepsilon$  for  $n \geq N$ , so  $\|T - TP_n\|_{\text{op}} \rightarrow 0$ .

Since  $T^*$  is also compact, we have  $\|T - P_n T\| \rightarrow 0$ . Thus,  $(P_n)_n$  is an approximate identity.

**Exercise:** Let  $(F_\alpha)_\alpha$  be an increasing net of finite rank projections on a Hilbert space  $\mathcal{H}$  with  $(F_\alpha)_\alpha \xrightarrow{\text{SOT}} I_{\mathcal{H}}$ . Prove that  $(F_\alpha)_\alpha$  is an approximate unit for  $\mathbb{K}(\mathcal{H})$ .

**Solution:** For any  $\xi \in \mathcal{H}$ , we have that

$$\|F_\alpha(\xi) - \xi\| \rightarrow 0.$$

Thus, for any  $T \in \mathbb{K}(\mathcal{H})$ , we have

$$\begin{aligned} \|TF_\alpha(\xi) - T(\xi)\| &= \|T(F_\alpha(\xi) - \xi)\| \\ &\leq \|T\|_{\text{op}} \|F_\alpha(\xi) - \xi\| \\ &\rightarrow 0. \end{aligned}$$

Similarly, by taking adjoints, we find that  $(F_\alpha)_\alpha$  is an approximate identity for  $\mathbb{K}(\mathcal{H})$ .

Indeed, every  $C^*$ -algebra admits an approximate unit. To do this, we create a family of operator-monotone continuous maps to apply the continuous functional calculus to.

**Definition.** A continuous function  $f: [0, \infty) \rightarrow [0, \infty)$  is called operator monotone if  $f(a) \leq f(b)$  whenever  $a \leq b$  for positive elements  $a, b$  in a  $C^*$ -algebra.

For example, we have shown that  $f(t) = t^{1/2}$  is operator monotone.

**Lemma:** Let  $n = 1, 2, \dots$ , and consider

$$f_n(t) = \frac{nt}{1 + nt}.$$

Then, the following hold.

- (1) For all  $n$ ,  $f_n$  is continuous,  $f_n(0) = 0$ , and  $0 \leq f_n \leq 1$ .
- (2) For all  $n$ ,  $f_n(t) = 1 - (1 + nt)^{-1} = t(1/n + t)^{-1}$ .
- (3) For all  $n \leq m$ ,  $f_n \leq f_m$ .
- (4) For all  $n$ ,  $f_n$  is operator-monotone.
- (5) If  $n \leq m$  and  $a, b$  are positive elements of a unital  $C^*$ -algebra with  $a \leq b$ , then  $0 \leq f_n(a) \leq f_m(b) \leq 1$ .

*Proof.* We will prove (4) and (5).

To see (4), we let  $A$  be a  $C^*$ -algebra with  $a \in A_+$ . We see that  $f_n(a) \in C^*(a)$ , as  $f_n(0) = 0$ . Thus, we assume  $A$  is unital. Fix  $n \in \mathbb{N}$ . If  $a \leq b$ , then  $1_A + na \leq 1_A + nb$ , and since  $1_A + na$  is invertible, we get  $(1_A + na)^{-1} \geq (1_A + nb)^{-1}$ . Thus, we have  $1_A - (1_A + na)^{-1} \leq 1_A - (1_A + nb)^{-1}$ . Using the functional calculus at  $a$  and  $b$ , we get

$$\begin{aligned} f_n(a) &= \phi_a(f_n) \\ &= 1_A - (1_A + na)^{-1} \\ &\leq 1_A - (1_A + nb)^{-1} \\ &= \phi_b(f_n) \\ &= f_n(b). \end{aligned}$$

To see (5), since the functional calculus preserves ordering, we have  $f_n(a) \leq f_m(a)$  and  $0 \leq f_m(b) \leq 1_A$ . Thus, we get  $0 \leq f_n(a) \leq f_m(a) \leq f_m(b) \leq 1_A$ .  $\square$

**Proposition:** Let  $A$  be a unital  $C^*$ -algebra, and let  $I \subseteq A$  be a closed left ideal. There is an increasing net  $(e_\alpha)_\alpha$  of positive contractions in  $I$  such that

$$\|x - xe_\alpha\| \rightarrow 0$$

for all  $x \in I$ .

*Proof.* Let  $\mathcal{F}$  be the set of all finite subsets of  $I$  ordered by inclusion. For each  $\alpha \in \mathcal{F}$ , we write  $n_\alpha = \text{card}(\alpha)$ . Given  $\alpha = \{x_1, \dots, x_{n_\alpha}\} \in \mathcal{F}$ , we set

$$v = \sum_{j=1}^{n_\alpha} x_j^* x_j.$$

We have that  $v_\alpha \in I$ , and  $v_\alpha \geq 0$  for each  $\alpha \in \mathcal{F}$ . We set  $e_\alpha = f_{n_\alpha}(v_\alpha)$ , where  $f_{n_\alpha}$  is defined as in the previous lemma.

Since  $I$  is a left ideal, each  $e_\alpha$  belongs to  $I$ , and if  $\alpha \leq \beta$  in  $\mathcal{F}$ , we have  $n_\alpha \leq n_\beta$ , and  $v_\alpha \leq v_\beta$ , so  $e_\alpha \leq e_\beta$ .

Given  $x \in \alpha \in \mathcal{F}$ , we see that  $x^*x \leq v_\alpha$ , so

$$\begin{aligned} 0 &\leq (1_A - e_\alpha)x^*x(1_A - e_\alpha) \\ &\leq (1 - e_\alpha)v_\alpha(1 - e_\alpha) \\ &= (1_A - f_{n_\alpha}(v_\alpha))^2 v_\alpha \\ &\leq \frac{1_A}{4n_\alpha}. \end{aligned}$$

Thus, for all  $x \in \alpha$ , we have

$$\begin{aligned} \|x - xe_\alpha\|^2 &= \|x(1_A - e_\alpha)\|^2 \\ &= \|(1_A - e_\alpha)x^*x(1_A - e_\alpha)\| \\ &\leq \frac{1}{4n_\alpha}, \end{aligned}$$

where we used the  $C^*$ -identity. Thus, given  $x \in I$  and  $\varepsilon > 0$ , we find a finite set  $\beta \subseteq I$  such that  $x \in \beta$  and  $\frac{1}{2\sqrt{n_\beta}} < \varepsilon$ . If  $\alpha \geq \beta$ , then  $x \in \alpha$  and  $n_\alpha \geq n_\beta$  meaning

$$\begin{aligned} \|x - xe_\alpha\| &\leq \frac{1}{2\sqrt{n_\alpha}} \\ &\leq \frac{1}{2\sqrt{n_\beta}} \\ &\leq \varepsilon. \end{aligned}$$

□

**Corollary:** If  $I \subseteq A$  is a closed two-sided ideal in a unital  $C^*$ -algebra  $A$ , then  $I$  is self-adjoint.

*Proof.* Since  $I$  is a closed left-ideal, we have a net  $(e_\alpha)_\alpha$  satisfying  $\|x - xe_\alpha\| \rightarrow 0$ . Let  $x \in I$ . Note that  $e_\alpha x^*$  is also in  $I$  as  $I$  is a right ideal. Thus,

$$\begin{aligned} \|x^* - e_\alpha x^*\| &= \|(x - xe_\alpha)^*\| \\ &= \|x - xe_\alpha\| \\ &\rightarrow 0, \end{aligned}$$

so  $x^* \in I$ .

□

**Theorem:** Every  $C^*$ -algebra  $A$  admits an approximate unit. If  $A$  is separable, then  $A$  admits a sequential approximate unit

*Proof.* If  $A$  is unital, then  $e_n = 1_A$  is an approximate unit. We may assume  $A$  is nonunital.



Embedding  $A$  into its unitization,  $A \subseteq \tilde{A}$  is a closed two-sided ideal. Thus, we get a net  $(e_\alpha)_\alpha$  of positive contractions in  $A$  such that  $\|x - xe_\alpha\| \rightarrow 0$  and  $\|x - e_\alpha x\| \rightarrow 0$ . This is our desired approximate unit.

If  $A$  is separable, and  $(e_\alpha)_\alpha$  is an approximate unit for  $A$ , we find an increasing sequence of finite subsets  $F_1 \subseteq F_2 \subseteq F_3 \subseteq \dots \subseteq A$  such that  $F = \bigcup_{k \geq 1} F_k$  is norm-dense in  $A$ . Since  $F_1$  is finite, there is  $\alpha_1$  with  $\|x - xe_{\alpha_1}\| < 1$  for all  $\alpha \geq \alpha_1$  and  $x \in F_1$ .

We then find  $\alpha_2 \geq \alpha_1$  with  $\|x - xe_{\alpha_2}\| < 1/2$  for all  $\alpha \geq \alpha_2$  and  $x \in F_2$ , and continue finding an increasing sequence  $(\alpha_n)_n$  with  $\|x - xe_{\alpha_n}\| < 1/n$  for all  $\alpha \geq \alpha_n$  and  $x \in F_n$ .

Set  $e_n = e_{\alpha_n}$ . We see that  $\|x - xe_n\| \rightarrow 0$  for all  $x \in F$ . For any  $x \in A$ , we find a sequence  $(x_n)_n \rightarrow x$  contained in  $F$ , and use a  $\varepsilon/3$  argument to find that

$$\begin{aligned} \|x - xe_n\| &\leq \|x - x_n\| + \|x_n - x_n e_n\| + \|x_n e_n - xe_n\| \\ &\rightarrow 0. \end{aligned}$$

Thus,  $(e_n)_n$  is an increasing sequence of positive contractions that is an approximate unit for  $A$ . □

**Exercise:** Let  $A$  be a unital  $C^*$ -algebra, and suppose  $I \subseteq A$  is a closed ideal with approximate identity  $(e_\alpha)_\alpha$ . Set  $f_\alpha = 1_A - e_\alpha$ . Prove that for every  $x \in I$ ,

$$(e_\alpha x e_\alpha)_\alpha \rightarrow x$$

and

$$(f_\alpha x f_\alpha)_\alpha \rightarrow 0.$$

**Solution:** We see that

$$\begin{aligned} \|e_\alpha x e_\alpha - x\| &\leq \|e_\alpha x e_\alpha - e_\alpha x\| + \|e_\alpha x - x\| \\ &\leq \|e_\alpha\| \|x e_\alpha - x\| + \|e_\alpha x - x\| \\ &\leq \|x e_\alpha - x\| + \|e_\alpha x - x\| \\ &\rightarrow 0. \end{aligned}$$

Additionally, we have that

$$\begin{aligned} \|(1 - e_\alpha)x(1 - e_\alpha)\| &= \|(1 - e_\alpha)(x - xe_\alpha)\| \\ &\leq \|1 - e_\alpha\| \|x - xe_\alpha\| \\ &\rightarrow 0. \end{aligned}$$

The existence of an approximate unit can also allow us to understand the structure of ideals and quotients.

**Exercise:** Let  $A$  be a  $C^*$ -algebra. Let  $a \in A$ . Show that the closed ideal generated by  $a$  has the picture

$$J(a) = \overline{\text{span}} \left( \left\{ \sum_{j=1}^n x_j a y_j \mid n \in \mathbb{N}, x_j, y_j \in A \right\} \right).$$

**Solution:** Consult Rainone.

**Exercise:** Let  $I \subseteq A$  be a closed ideal in a  $C^*$ -algebra  $A$ , and suppose  $J \subseteq I$  is a closed ideal in  $I$ . Prove that  $J$  is an ideal in  $A$  as well.

When we work with  $C^*$ -algebras, we require essential ideals to be closed.

**Proposition:** Let  $A$  be a  $C^*$ -algebra. Suppose  $I \subseteq A$  is a closed ideal. The following are equivalent:

- (i) for every nonzero closed ideal  $J \subseteq A$ ,  $J \cap I \neq \{0\}$ ;
- (ii) if  $a \in A$  and  $aI = \{0\}$ , then  $a = 0$ ;

(iii) if  $a \in A$  and  $Ia = \{0\}$ , then  $a = 0$ .

*Proof.* The implication (ii) if and only if (iii) follows from taking adjoints.

Suppose  $a \neq 0$  and  $aI = \{0\}$ . Then, for all  $x, y \in A$  and  $z \in I$ , we have  $(xay)z = xa(yz) = 0$ , since  $yz \in I$ . Thus,  $J(a)I = \{0\}$ , so  $J(a) \cap I = \{0\}$ , which contradicts our hypothesis since  $J(a) \neq 0$ .

Let  $J$  be any nonzero closed ideal in  $A$ , and suppose  $a \neq 0$  belongs to  $J$ . Then,  $\{0\} \neq aI \subseteq JI = J \cap I$ .  $\square$

**Exercise:** Let  $\mathcal{H}$  be a Hilbert space. Show that  $\mathcal{K}(\mathcal{H}) \subseteq \mathcal{B}(\mathcal{H})$  is an essential ideal.

**Solution:** We are aware that any nonzero closed ideal  $J \subseteq \mathcal{B}(\mathcal{H})$  contains  $\mathcal{K}(\mathcal{H})$ , so necessarily we must have  $\mathcal{K}(\mathcal{H}) \cap J \neq 0$ .

**Exercise:** Let  $\Omega$  be a LCH space. Show that  $C_0(\Omega) \subseteq C_b(\Omega)$  is an essential ideal.

**Solution:** Note that since  $\Omega$  is LCH, by Urysohn's lemma we know that  $C_0(\Omega)$  separates points. Thus, it is the case that if  $fC_0(\Omega) = \{0\}$ , we must have  $f = 0$ , so  $C_0$  is an essential ideal.

Now, we can define a quotient  $C^*$ -algebra, as well as the norm on the quotient  $C^*$ -algebra, using the approximate unit.

**Theorem:** Let  $A$  be a (unital)  $C^*$ -algebra, and suppose  $I \subseteq A$  is a closed ideal.

(1) If  $(e_\alpha)_\alpha$  is an approximate identity for  $I$ , the quotient norm for a coset  $a + I \in A/I$  is given by

$$\begin{aligned} \|a + I\| &= \lim_{\alpha} \|a - ae_\alpha\| \\ &= \lim_{\alpha} \|a - e_\alpha a\|. \end{aligned}$$

(2) The quotient  $A/I$  equipped with the quotient norm is again a (unital)  $C^*$ -algebra.

*Proof.*

(1) We may assume  $A$  is unital by unitizing. We see that  $\|a + I\| = \text{dist}_I(a) \leq \|a - ae_\alpha\|$ , as  $ae_\alpha \in I$ . Given  $\varepsilon > 0$ , we can find  $x \in I$  such that  $\|a - x\| < \|a + I\| + \varepsilon/2$ . For every  $\alpha$ , we have

$$\begin{aligned} \|a - ae_\alpha\| &= \|a - x + x - xe_\alpha + xe - \alpha - ae_\alpha\| \\ &= \|(a - x)(1_A - e_\alpha) + x - xe_\alpha\| \\ &\leq \|a - x\| + \|x - xe_\alpha\| \\ &< \|x - xe_\alpha\| + \|a + I\| + \varepsilon/2. \end{aligned}$$

Since  $(xe_\alpha) \rightarrow x$ , we have  $\alpha_0$  such that  $\|x - xe_\alpha\| < \varepsilon/2$  for all  $\alpha \geq \alpha_0$ . Thus, for all  $\alpha \geq \alpha_0$ , we have

$$\begin{aligned} \|a + I\| &\leq \|a - ae_\alpha\| \\ &\leq \|a + I\| + \varepsilon, \end{aligned}$$

so  $\|a + I\|_{A/I} = \lim_{\alpha} \|a - ae_\alpha\|$ . The second limit follows from taking adjoints.

(2) We only need to show that  $A/I$  equipped with the quotient norm satisfies the  $C^*$ -identity. Since the norm is submultiplicative, and the adjoint is isometric, it suffices to show that

$$\|a + I\|^2 \leq \|a^*a + I\|.$$

Let  $(e_\alpha)_\alpha$  be an approximate unit for  $I$ , and set  $f_\alpha = 1_A - e_\alpha$ . Given  $x \in I$ , we have

$$\begin{aligned} \|a + I\|^2 &= \left( \lim_{\alpha} \|a - ae_\alpha\| \right)^2 \\ &= \lim_{\alpha} \|a - ae_\alpha\|^2 \end{aligned}$$

$$\begin{aligned}
&= \|af_\alpha\|^2 \\
&= \lim_\alpha \|f_\alpha a^* a f_\alpha\| \\
&= \lim_\alpha \|f_\alpha(a^* a - \chi)f_\alpha + f_\alpha \chi f_\alpha\| \\
&\leq \sup_\alpha \|f_\alpha(a^* a - \chi)f_\alpha\| + \lim_\alpha \|f_\alpha \chi f_\alpha\| \\
&\leq \|a^* a - \chi\|,
\end{aligned}$$

where we use  $f_\alpha \chi f_\alpha \rightarrow 0$  and  $\|f_\alpha\| \leq 1$ . Taking the infimum, we get

$$\|a + I\|^2 \leq \|a^* a + I\|.$$

□

**Corollary:** Let  $\varphi: A \rightarrow B$  be a  $*$ -homomorphism between  $C^*$ -algebras. Then,  $\text{Ran}(\varphi) \subseteq B$  is a  $C^*$ -subalgebra.

*Proof.* We know that  $\text{Ran}(\varphi)$  is a  $*$ -subalgebra, so we must show that it is closed.

Since  $\varphi$  is contractive, the  $*$ -ideal  $\ker(\varphi) \subseteq A$  is closed, so  $A/\ker(\varphi)$  is a  $C^*$ -algebra. By the first isomorphism theorem,  $\phi: A/\ker(\varphi) \rightarrow B$  is injective, hence isometric. Since the isometric image of a complete space is complete, it is closed, and we know that  $\text{Ran}(\varphi) = \text{Ran}(\phi)$ . □

## Representations, States, and the GNS Construction

Up until now, we have defined  $C^*$ -algebras (and their properties) abstractly, with some vague references to concrete examples of  $C^*$ -algebras like  $\mathbb{B}(\mathcal{H})$ . It turns out that every abstract  $C^*$ -algebra is isometrically  $*$ -isomorphic to some subalgebra of  $\mathbb{B}(\mathcal{H})$ . This will require a lot of machinery.

Representations of  $C^*$ -algebras are defined as they are for  $*$ -algebras.

**Definition.** Let  $A$  be a  $C^*$ -algebra.

- (1) A representation of  $A$  is a pair  $(\pi, \mathcal{H})$ , where  $\mathcal{H}$  is a Hilbert space and  $\pi: A \rightarrow \mathbb{B}(\mathcal{H})$  is a  $*$ -homomorphism. If  $A$  is unital and  $\pi(1_A) = I_{\mathcal{H}}$ , we say the representation is unital.
- (2) If  $(\pi, \mathcal{H})$  is a representation of  $A$ , a closed subspace  $\mathcal{K} \subseteq \mathcal{H}$  is called reducing for  $(\pi, \mathcal{H})$  if  $\mathcal{K}$  reduces  $\pi(A)$ .
- (3) A representation  $(\pi, \mathcal{H})$  is called irreducible if  $\{0\}$  and  $\mathcal{H}$  are the only reducing spaces for  $(\pi, \mathcal{H})$ . Otherwise, we say  $(\pi, \mathcal{H})$  is reducible.
- (4) A representation  $(\pi, \mathcal{H})$  of  $A$  is called cyclic if there is a vector  $\xi_\pi \in \mathcal{H}$  such that  $[\pi(A)\xi_\pi] = \mathcal{H}$ . We call  $\xi_\pi$  a cyclic vector for  $\pi$ .
- (5) A representation  $(\pi, \mathcal{H})$  is called nondegenerate if  $\pi(A)$  acts nondegenerately on  $\mathcal{H}$  — that is,  $\ker(\pi(A)) = \{0\}$ , or  $[\pi(A)\mathcal{H}] = \mathcal{H}$ .
- (6) Two representations  $(\pi, \mathcal{H})$  and  $(\rho, \mathcal{K})$  are called unitarily equivalent if there exists a unitary  $U: \mathcal{H} \rightarrow \mathcal{K}$  such that

$$U\pi(a)U^* = \rho(a)$$

for all  $a \in A$ .

**Remark:** If  $\pi: A \rightarrow \mathbb{B}(\mathcal{H})$  is a representation of a  $C^*$ -algebra  $A$ , then  $\pi$  is faithful if and only if  $\pi$  is injective if and only if  $\pi$  is isometric, as  $\pi$  is a  $*$ -homomorphism between  $C^*$ -algebras.

If  $\mathcal{K} \subseteq \mathcal{H}$  is reducing for  $\pi: A \rightarrow \mathbb{B}(\mathcal{H})$ , we can speak of the reduced representation  $\pi|_{\mathcal{K}}: A \rightarrow \mathbb{B}(\mathcal{K})$ , where  $\pi|_{\mathcal{K}}(a) = \pi(a)|_{\mathcal{K}}$ . Additionally, since  $\mathcal{K}^\perp$  reduces  $\pi(A)$ , we also have  $\pi|_{\mathcal{K}^\perp}$  is a reduced representation.