

Introduction

Oh hey, it's another one of those textbook notes that I never complete. I've decided to try something different in order to develop my understanding of measure theory. One of the primary for understanding measure theory is Gerald B. Folland's *Real Analysis and Applications* — and one of the benefits it has over a lot of other texts is that it has a significant number of exercises. I'm going to try to do them all — I'll start with Chapters 1–3, and if that goes well enough, continue up through whatever chapter I end up having to tap out at. Interspersed, I will include various notes. I figure that in order to make a subject like measure theory really stick, I need to deal with it consistently.

Contents

Introduction	1
Chapter 1	1
Section 1.2	1

Chapter 1

Section 1.2

Exercise (Exercise 1): A family of sets $\mathcal{R} \subseteq \mathcal{P}(X)$ is called a ring if it is closed under finite unions and differences. A ring that is closed under countable unions is called a σ -ring.

- (a) Rings (σ -rings) are closed under finite (countable) intersections.
- (b) If \mathcal{R} is a ring (σ -ring), then \mathcal{R} is an algebra (σ -algebra) if and only if $X \in \mathcal{R}$.
- (c) If \mathcal{R} is a σ -ring, then $\{E \subseteq X \mid E \in \mathcal{R} \text{ or } E^c \in \mathcal{R}\}$ is a σ -algebra.
- (d) If \mathcal{R} is a σ -ring, then $\{E \subseteq X \mid E \cap F \in \mathcal{R} \text{ for all } F \in \mathcal{R}\}$ is a σ -algebra.

Proposition (Proposition 1.2): The Borel σ -algebra, $\mathcal{B}_{\mathbb{R}}$, is generated by each of the following:

- (a) the open intervals, $\mathcal{E}_1 = \{(a, b) \mid a < b\}$;
- (b) the closed intervals, $\mathcal{E}_2 = \{[a, b] \mid a < b\}$;
- (c) the half-open intervals, $\mathcal{E}_3 = \{(a, b] \mid a < b\}$ or $\mathcal{E}_4 = \{[a, b) \mid a < b\}$;
- (d) the open rays, $\mathcal{E}_5 = \{(a, \infty) \mid a \in \mathbb{R}\}$ or $\mathcal{E}_6 = \{(-\infty, a) \mid a \in \mathbb{R}\}$;
- (e) the closed rays, $\mathcal{E}_7 = \{[a, \infty) \mid a \in \mathbb{R}\}$ or $\mathcal{E}_8 = \{(-\infty, a] \mid a \in \mathbb{R}\}$.

Proof. The elements for \mathcal{E}_j for $j \neq 3, 4$ are open or closed, and the elements of $\mathcal{E}_3, \mathcal{E}_4$ are G_δ sets — for instance,

$$(a, b] = \bigcap_{n=1}^{\infty} \left(a, b + \frac{1}{n} \right).$$

Thus, $\sigma(\mathcal{E}_j) \subseteq \mathcal{B}_{\mathbb{R}}$ for each j . On the other hand, every open set in \mathbb{R} is a countable union of open intervals, so $\mathcal{B}_{\mathbb{R}} \subseteq \sigma(\mathcal{E}_1)$. Thus, $\mathcal{B}_{\mathbb{R}} = \sigma(\mathcal{E}_1)$. \square