

Problem 2

The standard topology on \mathbb{R} is generated by the metric on \mathbb{R} ; therefore, we may show that $[a, b]$ is sequentially compact.

If $(x_n)_n \subseteq [a, b]$ is any sequence, then by Bolzano–Weierstrass, there is a subsequence $(x_{n_k})_k \rightarrow x \in \mathbb{R}$. Furthermore, since $[a, b]^c = (-\infty, a) \cup (b, \infty)$ is open, we must have $x \in [a, b]$, meaning that $[a, b]$ is sequentially compact, hence compact.

Problem 17

Write

$$\sum_{i=1}^m |x_i + y_i|^p = \sum_{i=1}^m |x_i| |x_i + y_i|^{p-1} + |y_i| |x_i + y_i|^{p-1}.$$

Then, by applying Hölder's Inequality, we have

$$\begin{aligned} \sum_{i=1}^m |x_i| |x_i + y_i|^{p-1} &\leq \left(\sum_{i=1}^m |x_i|^p \right)^{1/p} \left(\sum_{i=1}^m |x_i + y_i|^{(p-1)q} \right)^q \\ \sum_{i=1}^m |y_i| |x_i + y_i|^{p-1} &\leq \left(\sum_{i=1}^m |y_i|^p \right)^{1/p} \left(\sum_{i=1}^m |x_i + y_i|^{(p-1)q} \right)^q. \end{aligned}$$

Since $(p-1)q = p$, we then have

$$\sum_{i=1}^m |x_i + y_i|^p \leq \left(\sum_{i=1}^m |x_i + y_i|^p \right)^q \left(\left(\sum_{i=1}^m |x_i|^p \right)^{1/p} + \left(\sum_{i=1}^m |y_i|^p \right)^{1/p} \right)^q,$$

and dividing, we get

$$\left(\sum_{i=1}^m |x_i + y_i|^p \right)^{1/p} \leq \left(\sum_{i=1}^m |x_i|^p \right)^{1/p} + \left(\sum_{i=1}^m |y_i|^p \right)^{1/p}.$$

Problem 19

(i) We see that

$$\sup_{i \in \mathbb{N}} |x_i| = 0$$

if and only if $|x_i| \leq 0$ for all i , meaning that $(x_i)_i$ is the zero sequence. Similarly,

$$\begin{aligned} \|\alpha x\| &= \sup_{i \in \mathbb{N}} |\alpha x_i| \\ &= |\alpha| \sup_{i \in \mathbb{N}} |x_i| \\ &= |\alpha| \|x\|. \end{aligned}$$

Finally,

$$\|x + y\| = \sup_{i \in \mathbb{N}} |x_i + y_i|$$

$$\begin{aligned}
&\leq \sup_{i \in \mathbb{N}} (|x_i| + |y_i|) \\
&\leq \sup_{i \in \mathbb{N}} |x_i| + \sup_{j \in \mathbb{N}} |y_j| \\
&= \|x\| + \|y\|,
\end{aligned}$$

meaning that $\|\cdot\|$ is a bona fide norm.

(ii) Let $B = \{x \in X \mid \|x\| \leq 1\}$. Let $(x_n)_n \subseteq B$ converge to $x \in \ell_\infty$ in the ℓ_∞ norm.

Note that for all n , $\sup_{i \in \mathbb{N}} |x_n(i)| \leq 1$, meaning that since

$$\sup_{i \in \mathbb{N}} |x(i) - x_n(i)| \rightarrow 0,$$

we have that

$$|x(i) - x_n(i)| \rightarrow 0$$

for each i , so

$$x_n(i) \rightarrow x(i)$$

for all i . Thus, $|x(i)| \leq 1$ for all i , meaning $\sup_{i \in \mathbb{N}} |x(i)| \leq 1$, so $\|x\| \in B$.

(iii) Let $\varepsilon = 1/2$, and consider the collection $(e_n)_n$ of sequences in ℓ_∞ consisting of 1 at position n and zero elsewhere. Then, $(e_n)_n \subseteq B$, but since $\sup_{i \in \mathbb{N}} |e_n(i) - e_m(i)| = 1$ for all $n \neq m$, we cannot have balls of radius $1/2$ cover the family $(e_n)_n$ with finitely many such balls, meaning that B is not totally bounded.

Problem 20

(i) We see that $d(x, y) = 0$ if and only if $x_n = y_n$ for each n , since each d_n is a metric; therefore, $d(x, y) = 0$ if and only if $x = y$.

Furthermore, we have that for all $x = (x_n)_n$, $y = (y_n)_n$, and $z = (z_n)_n$, $\frac{1}{2^n} d(x_n, z_n) \leq \frac{1}{2^n} d(x_n, y_n) + \frac{1}{2^n} d(y_n, z_n)$. Therefore, we get

$$\begin{aligned}
d(x, z) &= \sum_{n=1}^{\infty} \frac{1}{2^n} d(x_n, z_n) \\
&\leq \sum_{n=1}^{\infty} \frac{1}{2^n} d(x_n, y_n) + \frac{1}{2^n} d(y_n, z_n) \\
&\leq d(x, y) + d(y, z).
\end{aligned}$$

Since $d(x_n, y_n), d(y_n, z_n) \leq 1$, these sums must converge, so $d(x, y)$ is indeed a metric.

(ii) We will show that a sequence $(y_n)_n \subseteq X$ converges to $y \in X$ with the given distance metric if and only if it does so pointwise. This will show that the metric d induces the topology of pointwise convergence, which is exactly the topology τ_{prod} .

To start, let $(y_n)_n \rightarrow y$ in the given distance metric. Then, for all $\varepsilon > 0$, there is some $N \in \mathbb{N}$ such that for all $n \geq N$, we have

$$d(y_n, y) = \sum_{j=1}^{\infty} \frac{1}{2^j} d_j(y_n(j), y(j))$$

$$< \varepsilon,$$

so we see that for each j , $d(y_n(j), y(j)) < \varepsilon$, meaning that $y_n(j) \rightarrow y(j)$ for each j .

Let $(y_n)_n \rightarrow y$ pointwise. If $\varepsilon > 0$, convergence of series gives some J such that $\sum_{j=J+1}^{\infty} \frac{1}{2^j} < \varepsilon/2$, meaning that

$$\sum_{j=J+1}^{\infty} \frac{1}{2^j} d_j(y_n(j), y(j)) < \varepsilon/2$$

For $j = 1, \dots, J$, we find N_1, \dots, N_J such that for all $n \geq N_j$, $d_j(y_n(j), y(j)) < \varepsilon/2$. Therefore, for $n \geq \max(N_1, \dots, N_J)$, we have

$$\begin{aligned} d(y_n, y) &= \sum_{j=1}^{\infty} \frac{1}{2^j} d_j(y_n(j), y(j)) \\ &= \sum_{j=1}^J \frac{1}{2^j} d_j(y_n(j), y(j)) + \sum_{j=J+1}^{\infty} \frac{1}{2^j} d_j(y_n(j), y(j)) \\ &< \sum_{j=1}^J \frac{\varepsilon}{2^{j+1}} + \frac{\varepsilon}{2} \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon. \end{aligned}$$

Therefore, $(y_n)_n \rightarrow y$ in our given distance metric.

Since convergence of sequences in our given distance metric is given by pointwise convergence, the induced topologies must be equal, so $\tau_d = \tau_{\text{prod}}$.

(iii) We prove that X is complete if and only if X_n is complete for all n .

To see this, note that $(y_n)_n \subseteq X$ is Cauchy if and only if $(y_n(j))_n \subseteq X_j$ is Cauchy for each j , as for all $\varepsilon > 0$ and $m, n \geq N$ with $d(y_n, y_m) < \varepsilon$, then

$$\sum_{j=1}^{\infty} \frac{1}{2^j} d_j(y_n(j), y_m(j)) < \varepsilon,$$

meaning this holds for all j , and in the reverse direction, we use the same $\frac{\varepsilon}{2}$ method from part (ii).

The sequence $(y_n)_n$ thus converges in X if and only if every $y_n(j)$ converges in X_j (as τ_d is the topology of pointwise convergence), meaning that X is complete if and only if each X_j is complete.

Problem 21

Let X be complete, and let $(C_n)_n \subseteq P(X)$ be nonempty, decreasing, closed sets with $\text{diam}(C_n) \rightarrow 0$.

Let $(x_n)_n$ be defined by $x_n \in C_n$ for each n . Then, for any $\varepsilon > 0$, we may find C_N such that $\text{diam}(C_N) < \varepsilon$, meaning that for all $n, m \geq N$, we have that $x_n, x_m \in C_N$, so $d(x_n, x_m) < \varepsilon$, meaning that $(x_n)_n$ is Cauchy. Since X is complete, $(x_n)_n \rightarrow x$ for some $x \in X$. This point must be in all such C_n , meaning that

$$\bigcap_{n=1}^{\infty} C_n = \{x\}.$$

Now, let X be a metric space such that for any $(C_n)_n \subseteq \mathcal{P}(X)$ nonempty, decreasing, and closed with $\text{diam}(C_n) \rightarrow 0$, there is some $x \in X$ with $\bigcap_{n=1}^{\infty} C_n = \{x\}$. Let $(x_n)_n$ be a Cauchy sequence in X .

Define a family of closed sets by

$$C_n = \overline{\{x_n, x_{n+1}, \dots\}}.$$

We note the following:

- each of the C_n is closed;
- $C_n \supseteq C_{n+1}$ by construction, since $\{x_n, x_{n+1}, \dots\} \supseteq \{x_{n+1}, x_{n+2}, \dots\}$, and closures respect set inclusion;
- $\text{diam}(C_n) \rightarrow 0$, as $(x_n)_n$ is Cauchy, so if $\varepsilon > 0$, there is some N such that for all $n, m \geq N$, $d(x_n, x_m) < \varepsilon$, meaning that the diameter of the closure of the set $\{x_N, x_{N+1}, \dots\}$ is no more than ε .

Therefore, there is some $x \in X$ such that

$$\bigcap_{n=1}^{\infty} C_n = \{x\},$$

meaning that $(x_n)_n \rightarrow x$, and X is complete.