

## Complex Numbers

A complex number is an ordered pair of real numbers,  $(a, b) = a + bi$ . A vector in  $\mathbb{R}^2$  is also an ordered pair,  $(a, b)$  of real numbers.

Indeed, vector addition and scalar multiplication on complex numbers are defined just as with  $\mathbb{R}^2$ . However, unlike vectors in  $\mathbb{R}^2$ , there is also an operation  $\cdot$ . We desire for  $(0, 1) \cdot (0, 1) = (-1, 0)$ ; essentially,  $i^2 = -1$ . We say that  $i$  is a square root of  $-1$ ; every complex number except 0 has two square roots.

$$\begin{aligned}(a, b) \cdot (c, d) &= (a + bi) + (c + di) \\ &:= a(c) + adi + bci + bd(i^2) \\ &:= (ac - bd) + (ad + bc)i \\ &= (ac - bd, ad + bc)\end{aligned}$$

Thus,  $\mathbb{R}^2$  with the operations  $+$  and the above defined complex multiplication is known as  $\mathbb{C}$ . We write as  $a + bi$  instead of  $(a, b)$ .

Given  $z = (a + bi) \in \mathbb{C}$ , we write  $\operatorname{Re}(z) = a$  and  $\operatorname{Im}(z) = b$ . If  $\operatorname{Im}(z) = 0$ , then  $z \in \mathbb{R} \times \{0\} \subset \mathbb{C}$ . However, many people say that  $\mathbb{R} \subseteq \mathbb{C}$ , even if  $\mathbb{C}$  isn't defined as such.

## Reciprocals of Complex Numbers

Let  $z \in \mathbb{C}$ , where  $z \neq 0$ . Then,  $\exists w \in \mathbb{C}$  such that  $zw = 1$ .

Let  $w = c + di$ . We want to show that  $zw = 1$ .

$$(a + bi) + (c + di) = (ac - bd) + (ad + bc)i$$

with the condition that

$$\begin{aligned}ac - bd &= 1 \\ ad + bc &= 0.\end{aligned}$$

Thus, let  $w = c + di$ , with  $a, b \neq 0$

$$\begin{aligned}c &= \frac{a}{a^2 + b^2} \\ d &= \frac{-b}{a^2 + b^2}\end{aligned}$$

For every  $z \neq 0$ , with  $z = a + bi$ , the *reciprocal* of  $z$  is defined as  $\frac{1}{z} = \frac{a}{a^2 + b^2} + \frac{-b}{a^2 + b^2}i$ . Then, for  $w \in \mathbb{C}$ , we define

$$\frac{w}{z} := w \left( \frac{1}{z} \right).$$

## Properties of Complex Numbers

Let  $z = a + bi \in \mathbb{C}$ . Then, the (Euclidean) norm (or absolute value) of  $z$  is defined as

$$|z| = \sqrt{a^2 + b^2}.$$

The conjugate of  $z = a + bi$  is  $\bar{z} = a - bi$ .

$$(i) \quad z\bar{z} = |z|^2$$

$$(ii) \quad \overline{(\bar{z})} = z$$

$$(iii) \overline{(z + w)} = \bar{z} + \bar{w}$$

$$(iv) \overline{zw} = \bar{z} \cdot \bar{w}$$

$$(v) z + \bar{z} = 2\operatorname{Re}(z), \text{ so } \operatorname{Re}(z) = \frac{z + \bar{z}}{2}$$

$$(vi) z - \bar{z} = 2i\operatorname{Im}(z), \text{ so } \operatorname{Im}(z) = \frac{z - \bar{z}}{2i}$$

## Polar Representation

Let  $z = a + bi$  (or  $z = (a, b)$ ). Then,  $|z| = \sqrt{a^2 + b^2}$  is the *radius*, and the *argument* is found by  $\theta = \arctan(b/a)$  for  $a \neq 0$ . Therefore, the full polar representation is as follows:

$$z = |z| (\cos \theta + i \sin \theta). \quad \theta \in [0, 2\pi)$$

If  $z = 0$ , then  $|z| = 0$ , and  $\arg z$  is undefined.

For example, we can find  $\arg i$  in  $[\pi, 3\pi)$  as  $\frac{5\pi}{2}$ .

For  $z_1$  and  $z_2$  in polar form, we have:

$$|z_1 z_2| = |z_1| |z_2| \quad (1)$$

$$\arg(z_1 z_2) = \arg z_1 + \arg z_2 \pmod{2\pi} \quad (2)$$

Proof of (1):

$$\begin{aligned} |z_1 z_2|^2 &= (z_1 z_2) \overline{(z_1 z_2)} \\ &= z_1 z_2 \bar{z}_1 \bar{z}_2 \\ &= z_1 \bar{z}_1 z_2 \bar{z}_2 \\ &= |z_1|^2 |z_2|^2 \end{aligned}$$

Since  $|z| \geq 0$ , we get  $|z_1 z_2| = |z_1| |z_2|$ .

Let  $z = 2(\cos \pi/6 + i \sin \pi/6)$ , and let  $f : \mathbb{C} \rightarrow \mathbb{C}$  defined as  $f(w) = zw$ . Then,  $f$  rotates  $w$  by  $\pi/6$  and scales  $w$  by 2.

**Theorem:** For  $n \in \mathbb{N}$ , if  $z = r(\cos \theta + i \sin \theta)$ , then  $z^n = r^n(\cos(n\theta) + i \sin(n\theta))$ .

**Proof:** Induct on  $n$ . For the base case, we know that  $n = 1$  satisfies this property. For  $n > 1$ , we have:

$$\begin{aligned} z^{n+1} &= (z^n)(z) \\ &= (r^n(\cos(n\theta) + i \sin(n\theta))) r(\cos \theta + i \sin \theta) \\ &= (r^n)(r) (\cos(n\theta + \theta) + i \sin(n\theta + \theta)) && \text{Polar Representation Definition} \\ &= r^{n+1}(\cos((n+1)\theta) + i \sin((n+1)\theta)) \end{aligned}$$

We can use this technique to find the “roots of unity.” For example, to find all  $z$  such that  $z^3 = 1$ , we use our

technique:

$$\begin{aligned}
 z^3 &= 1 \\
 |z| &= 1 \\
 \arg z^3 &= 0 \\
 3 \arg z &= 0 \pmod{2\pi} \\
 \arg z &= \frac{k2\pi}{3} \\
 &= 0, \frac{2\pi}{3}, \frac{4\pi}{3} \\
 z_1 &= 1 \\
 z_2 &= (\cos 2\pi/3 + i \sin 2\pi/3) \\
 z_3 &= (\cos 4\pi/3 + i \sin 4\pi/3)
 \end{aligned}$$

We can see that  $z_2^2 = z_3$ .

For the  $n$  case, we find  $z_2 = \cos(2\pi/n) + i \sin(2\pi/n)$ , and  $z_k = z_2^{k-1}$ .

## Exponential, Logarithm, and Trigonometric Functions in $\mathbb{C}$

### Exponential

Let  $z = a + bi$ . We define  $e^{a+bi}$  as follows:

$$e^{a+bi} = e^a (\cos b + i \sin b)$$

Recall that for every nonzero complex number,  $z = |z| (\cos \theta + i \sin \theta)$ , where  $\theta = \arg z$ . Thus,

$$\begin{aligned}
 z &= |z| e^{i\theta} \\
 &= |z| e^{i \arg z}.
 \end{aligned}$$

The function  $e^z$  has some properties similar to the function  $e^x$  in real numbers, and some properties varying with the real numbers.

$$\begin{aligned}
 e^z e^w &= e^{z+w} \\
 e^z &\neq 0
 \end{aligned}$$

However, there are some differences:

$$\begin{aligned}
 |e^{i\theta}| &= 1 \\
 e^{a+bi} &= e^a
 \end{aligned}
 \quad \forall \theta$$

From these properties, we find Euler's equation:

$$e^{i\pi} + 1 = 0$$

Additionally,  $e^z$  is periodic, while  $f(x) = e^x$  is injective:

$$\begin{aligned}
 e^{z+2n\pi} &= e^z (\cos(2n\pi) + i \sin 2n\pi) \\
 &= e^z
 \end{aligned}$$

When examining the function  $f : \mathbb{C} \rightarrow \mathbb{C} \setminus \{0\}$ ,  $z \mapsto e^z$ , we find that the following happen:

- $f(\mathbb{R}) = (0, \infty)$  — we apply  $f(x) = e^x$ .
- $f(a + bi) = e^a e^{bi}$  —  $e^a$  is rotated by  $b$ .
- $f(\mathbb{R} + bi)$  is expressed as the line along  $b$  radians through the origin.
- Therefore,  $f(A_0) = \mathbb{C} \setminus \{0\}$ , where  $A_0 = \{a + bi \mid a \in \mathbb{R}, b \in [0, 2\pi)\}$ .

## Logarithm

Recall that for a function  $f : A \rightarrow B$ ,  $f^{-1}$  is a function if  $f$  is injective. However, for any  $f$ , it is the case that  $f^{-1}(b)$  does exist, defined as follows:

$$f^{-1}(b) = \{a \mid f(a) = b\}.$$

For the function  $f(z) = e^z$ ,  $f$  is not one to one, so for  $w = e^z$ ,  $f^{-1}(w) = \{z' \in \mathbb{C} \mid e^{z'} = w\}$ . We can find this as  $f^{-1}(w) = \{z + 2n\pi i \mid n \in \mathbb{Z}\}$ .

We define  $\log(w) := \{z \in \mathbb{C} \mid e^z = w\}$ . For a fixed  $\theta \in \mathbb{R}$ , we define  $\log_{A_\theta}(w) := \{z \mid e^z = w, z \in A_\theta\}$ .

Let  $z = 1 + \frac{5\pi}{2}i$ . Then,

$$\log_{A_{-\pi}} e^z = 1 + \frac{\pi}{2}i$$

Let  $w \in \mathbb{C} \setminus \{0\}$ . To find  $\log w$  (all values), then

$$\begin{aligned} z &\in \log w \\ e^z &= w \\ &= |w|e^{i \arg w} \\ e^{a+bi} &= |w|e^{i \arg w} \\ e^a e^{ib} &= |w|e^{i \arg w}. \end{aligned}$$

Therefore,  $a = \ln |w|$  and  $b = \arg w$ . Additionally, the following hold, for  $z_1, z_2 \in \mathbb{C}$ :

$$\log_{A_\theta}(z_1 z_2) = \log_{A_\theta}(z_1) + \log_{A_\theta}(z_2) + 2n\pi i$$

## Cosine and Sine

$$\begin{aligned} e^{ib} &= \cos b + i \sin b \\ e^{-ib} &= \cos b - i \sin b \\ \cos z &:= \frac{e^{iz} + e^{-iz}}{2} \\ \sin z &:= \frac{e^{iz} - e^{-iz}}{2i} \end{aligned}$$

## Complex Powers

Recall that for  $s, t \in \mathbb{R}$ ,  $s^t = e^{t \ln s}$ , where  $s > 0$ . For  $z, w \in \mathbb{C}$ ,  $z^w = e^{w \log z}$ , where  $z \neq 0$ .

$$\begin{aligned} (-2)^i &= e^{i \log(-2)} \\ &= e^{i(\ln(2) + i\pi)} \\ &= e^{i \ln 2 - (\pi + 2\pi n)} \\ &= e^{-\pi + 2\pi n + i \ln 2} \end{aligned}$$

This has *infinitely* many values.

Let  $\alpha = u + vi$ . Then,

$$\begin{aligned} z^\alpha &= e^{\alpha \log z} \\ &= e^{(u+vi)(\ln |z| + i \arg z)} \\ &= e^{(u \ln |z| - v \arg z)} e^{i(v \ln |z| + u \arg z)} \end{aligned}$$

Since  $\arg z = \theta + 2\pi n$  for some real  $\theta \in [0, 2\pi)$ ,

$$= e^{u \ln z} e^{-v(\theta + 2\pi n)} e^{i v \ln |z|} e^{i u(\theta + 2\pi n)}$$

Therefore, complex exponentiation is single-valued if  $\alpha \in \mathbb{R}$ . If  $\alpha \in \mathbb{Z}$ , then  $z^\alpha$  has only one value; if  $\alpha \in \mathbb{Q}$ , where  $\alpha = \frac{p}{q}$  and  $\gcd(p, q) = 1$ , then  $z^\alpha$  takes  $q$  distinct values, which are the  $q$ th-roots.

## Continuous Functions with Complex Domains

Let  $z \in \mathbb{C}$ , let  $r > 0$ .

- The set  $D(z; r) := \{w \in \mathbb{C} \mid |z - w| < r\}$  is the  $r$ -neighborhood of  $z$ .
- A subset  $A \subseteq \mathbb{C}$  is open if  $(\forall z \in A) (\exists r > 0) \ni D(z; r) \subseteq A$ .

For example, if  $A = \{z \mid \operatorname{Re}(z) > 0\}$ , we can find  $r$  equal to half the magnitude of the real component of  $z$  for any  $z \in A$ , meaning  $A$  is open.

Meanwhile, if  $A = \{z \mid \operatorname{Re}(z) \geq 0\}$ , this is not the case. If  $z = 0$ , then  $\nexists r > 0$  such that  $D(z; r) \subseteq A$ , as any open ball of radius  $r$  will have some element in  $\overline{A}$ .

- A subset  $B \subseteq \mathbb{C}$  is closed if  $\overline{B} \subseteq \mathbb{C}$  is open.

For example,  $A = \emptyset$  is open, by vacuous truth, so  $\overline{A} = \mathbb{C}$  is closed. Similarly, since  $\mathbb{C}$  is open,  $\emptyset$  is closed.

Meanwhile,  $A = \{x + iy \mid -1 \leq x < 1\}$  is neither open nor closed.

## Limits

Let  $A \subseteq \mathbb{C}$ ,  $f : A \rightarrow \mathbb{C}$ ,  $z_0 \in \mathbb{C}$ . Then,

$$\lim_{z \rightarrow z_0} f(z) = \ell$$

means both of the following hold:

- for some  $r > 0$ ,  $D(z_0; r) \setminus \{z_0\} \subseteq \operatorname{dom}(f)$ ,
- $\forall \varepsilon > 0, \exists \delta > 0$  such that  $f(D(z_0; \delta) \setminus \{z_0\}) \subseteq D(\ell; \varepsilon)$ .

For example, if

$$f(z) = \begin{cases} z & z \in \mathbb{C} \setminus \mathbb{R} \\ 3i & z \in \mathbb{R} \end{cases}$$

Then,  $\lim_{z \rightarrow 0} f(z)$  does not exist, as there is no  $\ell$  that satisfies both conditions. Specifically, if  $\ell = 3i$ , and we set  $\varepsilon = 1$ , then a disc of any radius around 0 has some  $z \in \mathbb{C} \setminus \mathbb{R}$  that maps to itself. Similarly, if we set  $\ell = 0$ , then there is a real number in a disc of any radius around 0.

**Note:**  $f$  does not have to be defined at  $z_0$  for the limit to be defined at  $z_0$ .

Let  $A \subseteq \mathbb{C}$  be open,  $f : A \rightarrow \mathbb{C}$ , and  $z_0 \in A$ . We say  $f$  is continuous at  $z_0$  if  $\lim_{z \rightarrow z_0} f(z) = f(z_0)$ . We say  $f$  is continuous on  $A$  if  $\forall z_0 \in A$ ,  $f$  is continuous at  $z_0$ .

We will show that  $f : \mathbb{C} \rightarrow \mathbb{C}$ ,  $z \mapsto 3z$  is continuous.

**Scratch Work:** We want  $\delta$  such that  $f(D(z_0; \delta)) \subseteq D(3z_0; \varepsilon)$ . Let  $z \in D(z_0; \delta)$ , meaning  $f(z) = 3z$ . We want  $3z \in D(3z_0; \varepsilon)$ , meaning we want  $|3z - 3z_0| < \varepsilon$ , or  $|z - z_0| < \frac{\varepsilon}{3}$ .

**Proof:** Let  $\varepsilon > 0$ . Set  $\delta = \frac{\varepsilon}{3}$ . We show  $f(D(z_0; \delta)) \subseteq D(f(z_0); \varepsilon)$ . Let  $z \in D(z_0; \delta)$ . Then,  $|z - z_0| < \delta = \varepsilon/3$ , meaning  $3|z - z_0| < \varepsilon$ , meaning  $|3z - 3z_0| < \varepsilon$ , so  $|f(z) - f(z_0)| < \varepsilon$ . Therefore,  $f(z) \in D(f(z_0); \varepsilon)$ . Since  $f$  is continuous at arbitrary  $z_0$ ,  $f$  is continuous on  $\mathbb{C}$ .

## Sequences

A sequence  $z_1, z_2, \dots \in \mathbb{C}$ . A sequence converges to  $z_0 \in \mathbb{C}$  if

$$(\forall \varepsilon > 0)(\exists M \in \mathbb{N}) \ni \forall z_{n>M}, |z_n - z_0| < \varepsilon$$

In words, for any radius around  $z_0$ , we can find  $z_n$  arbitrarily close to  $z_0$  for sufficiently large  $n$ . We write  $z_n \rightarrow z_0$  if this is the case.

Let  $f : \mathbb{C} \rightarrow \mathbb{C}$ . Then,  $f$  is continuous on  $\mathbb{C}$  if and only if the following equivalent conditions are met:

- (i) the inverse image of every open set is open ( $f^{-1}(B) := \{a \in \mathbb{C} \mid f(a) \in B\}$ );
- (ii) the inverse image of every closed set is closed;
- (iii) for every sequence  $(z_n)_n$  such that  $(z_n)_n \rightarrow z_0$ ,  $f(z_n) \rightarrow f(z_0)$ .

Let

$$f(z) = \begin{cases} 0 & z = 0 \\ 1 & z \neq 0 \end{cases}.$$

This function is not continuous. We will check that (i)–(iii) fail.

- (i) Let  $B = D(0; 1)$ . Then,  $f^{-1}(B) = \{0\}$ , which is not open set.
- (ii) Let  $B = \text{cl}(D(1; 0.5))$ . Then,  $f^{-1}(B) = \mathbb{C} \setminus \{0\}$ , which is not closed.
- (iii) Let  $z_n = \frac{1}{n}$ . Then,  $(z_n)_n \rightarrow 0$ , but  $f(z_n) = 1$  for all  $n$ , meaning  $f(z_n) \rightarrow 1 \neq f(0)$ .

To show limit divergence, recall the definition of limit convergence:

$$\lim_{n \rightarrow \infty} z_n = z_0 \Leftrightarrow (\forall \varepsilon > 0)(\exists M \in \mathbb{N}) \ni \forall z_{n>M}, |z_n - z_0| < \varepsilon.$$

Let  $z_1, \dots \in \mathbb{C}$  be a sequence. Then,  $\lim_{n \rightarrow \infty} z_n = \infty$  means

$$(\forall M > 0)(\exists N \in \mathbb{N}) \ni \forall n > N, |z_n| > M.$$

In words,  $|z_n|$  is arbitrarily large for sufficiently large  $n$ .

## Connected Sets

Let  $a, b \in \mathbb{C}$ . A path from  $a$  to  $b$  is a continuous function  $p : [0, 1] \rightarrow \mathbb{C}$  such that  $p(0) = a$  and  $p(1) = b$ . Let  $S \subseteq \mathbb{C}$ . If  $p([0, 1]) \subseteq S$ , then  $p$  is a path in  $S$ .

We say  $S$  is *path-connected* if for any  $s, t \in S$ , there is a path in  $S$  from  $s$  to  $t$ .

Every set that is path-connected is connected, but not necessarily the other way around — if  $A$  is open and path connected, then  $A$  is connected.

An open, path-connected subset of  $\mathbb{C}$  is known as a region, or a domain.

Let  $A = \mathbb{R} \times \{0\}$  (or the  $x$  axis in  $\mathbb{C}$ ).  $A$  is not a region, as  $A$  is not an open set, even if  $A$  is path-connected.

$A \subseteq \mathbb{C}$  is bounded if there exists  $r > 0$  such that  $A \subseteq D(0; r)$ .  $A = \mathbb{R} \times \{0\}$  is not bounded.

If  $A \subseteq \mathbb{C}$ , then  $A$  is compact if  $A$  is closed and bounded. There are various properties of compact sets that make them particularly amenable towards analysis.

**Extreme Value Theorem:** Every real-valued continuous function on a compact domain attains its maximum and minimum values.

**Uniform Continuity Theorem:** Elaborated below.

## Uniform Continuity

Recall that if  $f : A \rightarrow \mathbb{C}$ ,  $f$  is continuous if  $\forall a \in A, \lim_{z \rightarrow a} f(z) = f(a)$ .

$$(\forall a \in A)(\forall \epsilon > 0)(\exists \delta_a > 0) \ni f(D(a; \delta_a)) \subseteq D(f(a); \epsilon) \quad \delta \text{ depends on } a$$

When  $f$  is uniformly continuous, there is one value of  $\delta$ , dependent on  $\epsilon$ , that applies for every value of  $a$ .

$$(\forall \epsilon > 0)(\exists \delta_\epsilon > 0) \ni (\forall a \in A), f(D(a; \delta_\epsilon)) \subseteq D(f(a); \epsilon)$$

## Riemann Sphere

Let  $S^2 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$ . Let  $N = (0, 0, 1)$  denote the north pole. Then, there is a continuous bijection from  $S^2 \setminus \{N\} \rightarrow \mathbb{C}$ .

We can visualize this by picking a random point on the sphere and drawing a line from the north pole through the sphere to this point, and finding the point that intersects the plane.

Consider the sequence  $z_n = n^2 i$  for  $n = 1, 2, \dots$ . We can see that, on the projection from  $z_n$  to the sphere, all the values of  $p$  converge to  $N$ . Therefore, we write  $\lim_{n \rightarrow \infty} z_n = \infty$ , where  $\infty$  corresponds to  $N$  on  $S^2$ .

We can define  $\bar{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$  to be the complex plane that includes the “point at infinity” (from the projection on  $S^2$  that corresponds to the north pole).

## Analytic Functions

Let  $f : A \subseteq \mathbb{C} \rightarrow \mathbb{C}$  where  $A$  is open. Let  $z_0 \in A$ . We say  $f$  is differentiable at  $z_0$  if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists.

## Rules of Differentiation

- $(f + g)' = f' + g'$
- $(fg)' = f'g + fg'$
- $\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{(g)^2}$
- $(f \circ g)' = g'(f' \circ g)$
- For  $n \in \mathbb{Z}$ ,  $(z^n)' = nz^{n-1}$

Let  $f(z) = \bar{z}$ . We will find this value by directly applying the definition of the derivative.

$$\begin{aligned} f'(z_0) &= \lim_{z \rightarrow z_0} \frac{\bar{z} - \bar{z}_0}{z - z_0} \\ &= \lim_{z \rightarrow z_0} \frac{\overline{z - z_0}}{z - z_0}. \end{aligned}$$

Let's approach  $z_0$  from the horizontal direction. Suppose  $z = z_0 + t$  for some  $t \in \mathbb{R}$ . Then,

$$\lim_{z \rightarrow z_0} \frac{\overline{z_0 + t} - \bar{z}_0}{z_0 + t - z_0} = 1.$$

Let's approach  $z_0$  from the horizontal direction. Suppose  $z = z_0 + ti$  for some  $t \in \mathbb{R}$ . Then,

$$\begin{aligned} \lim_{z \rightarrow z_0} \frac{\overline{z_0 + ti} - \bar{z}_0}{z_0 + ti - z_0} &= \frac{-ti}{ti} \\ &= -1. \end{aligned}$$

Since  $1 \neq -1$ , we find that the limit does not exist.

We see that complex-differentiability is a strong condition.

Suppose that  $f'(z_0) = 2i$ , meaning

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = 2i.$$

If  $z$  is close to  $z_0$ , then  $f(z) - f(z_0) \approx 2i(z - z_0)$ . Pictorially, we can visualize this as, for  $z_0$  sufficiently close to  $z$ , the vector  $z_0 - z$  is akin to a counterclockwise rotation and a scaling by 2. This is applicable for *all*  $z$  in sufficient proximity to  $z_0$ .

Specifically, we can see that the complex differentiable function is *angle-preserving*. The technical name for  $f$  is that  $f$  is *conformal*.

## Analytic Function

Let  $f : A \subseteq \mathbb{C} \rightarrow \mathbb{C}$ . If  $f$  is differentiable at every  $z_0 \in A$ , we say  $f$  is *analytic* on  $A$ .

If  $f$  is analytic on  $A$ , then  $f$  is infinitely differentiable on  $A$ .

If  $f$  is analytic on  $A$  and  $f'(z_0) \neq 0$  for some  $z_0 \in A$ , then  $f$  is conformal at  $z_0 \in A$ .



## Cauchy-Riemann Theorem

Given a function  $f(x, y) : \mathbb{R}^2 \rightarrow \mathbb{R}$ . Recall that we can take partial derivatives,  $\frac{\partial f}{\partial x}$ , and directional derivative  $\frac{\partial f}{\partial u}$  for some unit vector  $u$ .

However, for  $\mathbb{C}$ , there is only one derivative,  $f'(z_0)$ , meaning that regardless of direction,  $f'(z_0)$  exists and has one value. We can contextualize  $f(z) = f(x + yi) = u(x, y) + iv(x, y)$ , where  $u(x, y) \in \mathbb{R}$  and  $v(x, y) \in \mathbb{R}$ . Then,

$$\frac{\partial u}{\partial x} \neq \frac{\partial u}{\partial y}$$

and

$$\frac{\partial v}{\partial x} \neq \frac{\partial v}{\partial y}$$

but

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y}.$$

We can see this by first letting  $z = z_0 + \delta x$ .

$$\begin{aligned} f'(z_0) &= \lim_{z \rightarrow z_0} \frac{f(z_0 + \delta x) - f(z_0)}{z_0 + \delta x - z_0} \\ &= \lim_{z \rightarrow z_0} \frac{u(x_0 + \delta x, y_0) + iv(x_0 + \delta x, y_0) - (u(x_0, y_0) + iv(x_0, y_0))}{\delta x} \\ &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \end{aligned}$$

and in the  $y$  direction,

$$\begin{aligned} f'(z_0) &= \frac{1}{i} \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} \\ &= -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y}. \end{aligned}$$

We set these two values equal to find

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \\ \frac{\partial v}{\partial x} &= -\frac{\partial u}{\partial y}, \end{aligned}$$

which are the Cauchy-Riemann equations. The corresponding theorem states that if  $f'(z_0)$  exists, then the Cauchy-Riemann equations must hold.

For example, if  $f(z) = \bar{z}$ , with  $f(x + yi) = x - yi$ , we have  $u(x, y) = x$  and  $v(x, y) = -y$ . Then,

$$\begin{aligned} \frac{\partial u}{\partial x} &= 1 \\ \frac{\partial v}{\partial x} &= -1, \end{aligned}$$

meaning  $f$  is not complex-differentiable.

If  $f : A \rightarrow \mathbb{C}$  satisfies the Cauchy-Riemann equations at every  $z_0 \in A$ , then  $f$  is analytic on  $A$ .

If  $f : A \subseteq \mathbb{C} \rightarrow \mathbb{C}$  is analytic on  $A$ , then we know  $f'$  and  $f''$  are continuous. From multivariable calculus, we know that  $u_{xy} = u_{yx}$  if both are continuous. So,

$$\begin{aligned} u_{xy} &= \frac{\partial}{\partial y}(u_x) \\ &= \frac{\partial}{\partial y}(v_y) \\ &= v_{yy} \\ u_{yx} &= \frac{\partial}{\partial x}(u_y) \\ &= \frac{\partial}{\partial x}(-v_x) \\ &= -v_{xx} \end{aligned}$$

Therefore,  $v_{xx} + v_{yy} = 0$ . Similarly,  $u_{xx} + u_{yy} = 0$ .

If  $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}$  If  $\varphi_{xx} + \varphi_{yy} = 0$ , then we say  $\varphi$  is a harmonic function. Therefore, if  $f$  is an analytic function, then both the real and imaginary parts of  $f$  are harmonic.

Let  $A \subseteq \mathbb{R}^2$ . If  $u : A \rightarrow \mathbb{R}$  and  $v : A \rightarrow \mathbb{R}$ . Then,  $u$  and  $v$  are harmonic conjugates if  $u + iv$  is an analytic function. Additionally,  $u$  and  $v$  are harmonic conjugates if and only if they satisfy the Cauchy-Riemann equations.

We may ask if there exists an analytic function  $f$  such that  $\operatorname{Re}(f) = x^3 - 3xy^2 + y$ . Then,

$$\begin{aligned} v_y &= u_x = 3x^2 - 3y^2 \\ -v_x &= u_y = 1 - 6xy. \end{aligned}$$

Therefore, we find  $v = -x + 3x^2y - y^3 + c$  through integration. Therefore, we have

$$\begin{aligned} f(z) &= (x^3 - 3xy^2 + y) + i(3x^2y - y^3 - x + c) \\ &= (x - iy)^3 + y - ix + ic \\ &= z^3 + i(-iy + x) + ic \\ &= \bar{z}^3 + i(\bar{z} + c) \end{aligned}$$

Recall from from multivariable calculus that  $\nabla u \perp$  contour lines of  $u$ . Similarly,  $\nabla v \perp$  contour lines of  $v$ . Then, using the Cauchy-Riemann equations, we find

$$\begin{aligned} \nabla u \cdot \nabla v &= (-u_x u_y) + u_x u_y \\ &= 0, \end{aligned}$$

meaning the gradients are orthogonal to each other, meaning the contours of  $u$  are perpendicular to the contours of  $v$ .

## Inverse Functions

Let  $f : A \subseteq \mathbb{C} \rightarrow \mathbb{C}$ . Let  $z_0 \in A$ . If  $f$  is analytic on  $A$  and  $f'(z_0) \neq 0$ , then  $f$  is one to one on some neighborhood of  $z_0$ . Then,  $f^{-1} : f(N) \rightarrow N$  is analytic on  $f(N)$ , and

$$(f^{-1})'(f(z_0)) = \frac{1}{f'(z_0)}.$$

## Derivatives of Elementary Functions

Specifically, we will be working with complex exponentiation, complex trigonometric functions, and complex logarithms.

## Complex Exponential

$$\frac{d}{dz} e^z = e^z,$$

since, letting  $z = x + iy$ ,

$$\begin{aligned} e^z &= e^x e^{iy} \\ &= e^x (\cos(y) + i \sin(y)). \\ \frac{d}{dz} e^z &= \frac{\partial}{\partial x} e^z && \text{treating } y \text{ as constant} \\ &= e^x (\cos(y) + i \sin(y)) \\ &= e^{x+iy} \\ &= e^z. \end{aligned}$$

We know that  $e^z$  is continuous on  $\mathbb{C}$ , but this doesn't imply differentiability at every  $z_0 \in \mathbb{C}$ . We can verify by checking the Cauchy-Riemann equations, where  $u(x, y) = e^x \cos(y)$  and  $v(x, y) = e^x \sin(y)$ . Then,

$$\begin{aligned} \frac{\partial u}{\partial x} &= e^x \cos(y) \\ &= \frac{\partial v}{\partial y} \\ \frac{\partial v}{\partial y} &= -e^x \sin(y) \\ &= -\frac{\partial u}{\partial x}. \end{aligned}$$

If a function is analytic on  $\mathbb{C}$ , then  $f$  is known as entire.

## Complex Logarithm

We might ask where  $\log z$  is analytic. Let  $f(z) = e^z$ . Then,  $\log z = f^{-1}(z)$ ; since  $f$  is not one to one, we restrict the domain of  $f$  to  $A_\theta = \{z \mid \operatorname{Im}(z) \in [\theta, \theta + 2\pi)\}$  for any  $\theta$ .

Since  $f|_{A_\theta}$  is one to one, then

$$\left(f|_{A_\theta}\right)^{-1} = \log_{A_\theta}.$$

Fixing  $\theta$ , set  $g = f|_{A_\theta}$ . Then,

$$g^{-1}(g(z)) = z.$$

Because  $g$  is analytic on  $A_\theta$ ,  $g^{-1}$  is analytic on  $A_\theta$ . By chain rule, we have

$$\begin{aligned} \frac{d}{dz} (g^{-1}(g(z))) &= \frac{d}{dz} z \\ g^{-1'}(g(z)) &= \frac{1}{g'(z)} && g'(z) \neq 0 \\ g^{-1}(w) &= \frac{1}{g'(z)} && w = e^z \\ &= \frac{1}{e^z} \\ &= \frac{1}{w}. \end{aligned}$$

Therefore,  $\frac{d}{dw} \log_{A_\theta}(z) = \frac{1}{z}$ . Therefore,  $\text{dom}(\log_{A_\theta}) = \text{ran}(e^z_{A_\theta}) = \mathbb{C} \setminus \{0\}$ . However,  $\log_{A_0}$  (setting  $\theta = 0$ ) is not even continuous on  $\mathbb{C} \setminus \{0\}$ !

Specifically, at  $z = 0$ ,  $e^z = 1$ . Travelling around the unit circle counterclockwise in the image, we see that the preimage of these points travels along the imaginary axis. Approaching 1 "from the bottom," we find that the preimage of the points approaches  $2\pi$  in the domain. However, they ought to be approaching 0. Therefore, the limit doesn't exist.

However, notice that the domain is not open! To fix this, we will let  $B_\theta = \{z \in \mathbb{C} \mid \text{Im}(z) \in (\theta, \theta + 2\pi)\}$ .

Our log function *is* when  $e^z$  is restricted to  $B_\theta$ . Then,  $\log_{B_\theta}$  is analytic on  $\mathbb{C} \setminus \{re^{i\theta} \mid r \geq 0\}$ . When  $\theta = -\pi$ , then  $\log_{B_{-\pi}}$  is the principle branch of  $\log z$ .

Then, the domain is  $\mathbb{C} \setminus \{z \mid z = x + 0i, x < 0\}$  and the range is  $B_{-\pi}$ .

## Powers

Let  $\alpha \in \mathbb{C}$ . We might ask

$$\frac{d}{dz} \alpha^z$$

$$\frac{d}{dz} z^\alpha.$$

Recall that  $a^b = e^{b \log a}$ . Specifically,  $a^b = e^{b(\ln|a| + i \arg a)}$ .

$$\frac{d}{dz} \alpha^z = \frac{d}{dz} e^{z \log \alpha}$$

Fix  $\theta$ . Then,

$$= \frac{d}{dz} e^{z \log_{A_\theta} \alpha}$$

$$= \log_{A_\theta} \alpha e^{z \log_{A_\theta} \alpha} = \alpha^z \log_{A_\theta} \alpha. \quad \text{assuming analytic domain}$$

Specifically, as long as  $\alpha \notin \{re^{i\theta} \mid r \geq 0\}$ ,  $z \log_{A_\theta} \alpha$  is analytic, meaning  $e^{z \log_{A_\theta} \alpha}$  is analytic (composition of analytic functions).

$$z^\alpha = e^{\alpha \log z}$$

$$= e^{\alpha \log_{B_\theta} z}$$

$$= e^{\alpha \log_{B_\theta} z} \frac{\alpha}{z}$$

$$= \alpha z^{\alpha-1}.$$

Specifically, this holds for  $z \notin \{re^{i\theta} \mid r \geq 0\}$ .

We know that  $\frac{d}{dz} \log_{B_{-\pi}}(z) = \frac{1}{z}$ . The domain of  $\log_{B_{-\pi}}$  is  $\mathbb{C} \setminus (-\infty, 0]$ .