

These are some notes I have taken from Gerald B. Folland's *A Course in Abstract Harmonic Analysis*, with some other textbooks for various sources.

Basic Properties of Topological Groups

Definition: A *topological group* is a group G with a topology such that the operation

$$\begin{aligned} m: G \times G &\rightarrow G \\ (x, y) &\mapsto xy \end{aligned}$$

is continuous with respect to the product topology on $G \times G$ and the operation

$$\begin{aligned} i: G &\rightarrow G \\ x &\mapsto x^{-1} \end{aligned}$$

is continuous with respect to the topology on G .

For a topological group G , we denote the unit element as 1_G , and we set

$$\begin{aligned} Ax &= \{yx \mid y \in A\} \\ xA &= \{xy \mid y \in A\} \\ A^{-1} &= \{y^{-1} \mid y \in A\} \\ AB &= \{xy \mid x \in A, y \in B\} \end{aligned}$$

for all subsets $A, B \subseteq G$ and elements $x \in G$.

Definition: A subset $A \subseteq G$ is called *symmetric* if $A = A^{-1}$.

Proposition: Let G be a topological group.

- (i) The topology of G is invariant under translations and inversion; that is, if U is open, then xU, Ux, U^{-1}, AU, UA are open for any $x \in G$ and subset $A \subseteq G$.
- (ii) For every neighborhood U of 1_G , there is a symmetric neighborhood V of 1_G such that $VV \subseteq U$.
- (iii) If H is a subgroup of G , so is \overline{H} .
- (iv) Every open subgroup of G is closed.
- (v) If A and B are compact sets in G , so is AB .

Proof.

- (i) This is equivalent to the separate continuity of $(x, y) \mapsto xy$ and $x \mapsto x^{-1}$; furthermore,

$$\begin{aligned} AU &= \bigcup_{x \in A} xU \\ UA &= \bigcup_{x \in A} Ux. \end{aligned}$$

- (ii) Since $(x, y) \mapsto xy$ is continuous at 1_G , then for every neighborhood U of 1_G , there are neighborhoods $W_1, W_2 \subseteq U$. We may take $V = W_1 \cap W_2 \cap W_1^{-1} \cap W_2^{-1}$.
- (iii) For $x, y \in \overline{H}$, there are nets $(x_\alpha)_\alpha \rightarrow x$ and $(y_\alpha)_\alpha \rightarrow y$; since $(x_\alpha y_\alpha) \rightarrow xy$ and $(x_\alpha^{-1})_\alpha \rightarrow x^{-1}$ by continuity of the operations, we have $xy, x^{-1} \in \overline{H}$.

- (iv) If H is open, then so are all the cosets xH ; since $G \setminus H$ is the union of all the cosets of H except for H itself, $G \setminus H$ is open, so H is closed.
- (v) Since $A \times B$ is compact, and AB is the continuous image of $A \times B$ under $(x, y) \mapsto xy$, we have AB is compact.

□

Now, if H is a subgroup of G , we let G/H be the space of left cosets of H , and $q: G \rightarrow G/H$ is the canonical quotient map, we may impose the quotient topology on G/H , meaning that $U \subseteq G/H$ is open if and only if $q^{-1}(U)$ is open. Thus, q maps open sets in G to open sets in G/H , as if $V \subseteq G$ is open, $q^{-1}(q(V)) = VH$ is also open, so $q(V)$ is open.

Proposition: Let H be a subgroup of a topological group G .

- (i) If H is closed, then G/H is Hausdorff.
- (ii) If G is locally compact, so is G/H .
- (iii) If H is normal, then G/H is a topological group.

Proof.

- (i) If $\bar{x} = q(x)$ and $\bar{y} = q(y)$ are distinct points in G/H , and since H is closed, xHy^{-1} is a closed set that does not contain 1_G . There is a symmetric neighborhood U of 1_G such that $UU \cap xHy^{-1} = \emptyset$; since $U = U^{-1}$ and $H = HH$ (H is a subgroup), we have $1_G \notin UxH(Uy)^{-1} = (UxH)(UyH)^{-1}$, so $UxH \cap UyH = \emptyset$. Therefore, $q(Ux)$ and $q(Uy)$ are disjoint neighborhoods of \bar{x} and \bar{y} .
- (ii) If U is a compact neighborhood of 1_G , $q(Ux)$ is a compact neighborhood of $q(x)$ in G/H .
- (iii) If $x, y \in G$, and U is a neighborhood of G/H , continuity of multiplication in G implies that there are neighborhoods V of x and W of y such that $VW \subseteq q^{-1}(U)$. We see that $q(V)$ and $q(W)$ are neighborhoods of $q(x)$ and $q(y)$ such that $q(V)q(W) \subseteq U$, meaning multiplication is continuous in G/H . Similarly, inversion is continuous.

□

Corollary: If G is T1, then G is Hausdorff, and if G is not T1, then $\overline{\{1_G\}}$ is a closed normal subgroup, and $G/\overline{\{1_G\}}$ is a Hausdorff topological group.

Proof. Since singletons are closed in any T1 space, the first assertion follows from part (i) in the previous proposition by taking $H = \{1_G\}$.

To see the second assertion, we note that $\overline{\{1_G\}}$ is a subgroup, and it is the smallest closed subgroup of G ; it is normal, as otherwise we would obtain a smaller closed subgroup by intersection with one of the conjugates, meaning the result follows from parts (i) and (iii) in the previous proposition by taking $H = \overline{\{1_G\}}$.

□

Thus, without loss of generality, we may assume that a topological group is Hausdorff (else take $G/\overline{\{1_G\}}$), and when we talk about locally compact groups, we are talking about topological groups that are locally compact and Hausdorff.

Proposition: Every locally compact group G has a subgroup G_0 that is open, closed, and σ -compact.

Proof. Let U be a symmetric compact neighborhood of 1_G , let $U_n = \prod_{i=1}^n U$, and let

$$G_0 = \bigcup_{n=1}^{\infty} U_n.$$

Then, G_0 is the group generated by U , so it is a subgroup; G_0 is open since U_{n+1} is a neighborhood of U_n for all n , and so G_0 is closed as all open subgroups are closed. Finally, since each U_n is a finite product of compact subsets of G , G_0 is σ -compact.

□

We thus see that G_0 is the disjoint union of cosets of G_0 , meaning G is a disjoint union of σ -compact spaces. In particular, if G is connected, then G is necessarily σ -compact.

Definition: Let $f: G \rightarrow \mathbb{C}$ be a function. The *translates* of f via $y \in G$ are defined by

$$\begin{aligned} L_y f(x) &= f(y^{-1}x) \\ R_y f(x) &= f(xy). \end{aligned}$$

Note that the maps $y \mapsto L_y$ and $y \mapsto R_y$ are group homomorphisms.

The function f is called left/right uniformly continuous if

$$\begin{aligned} \|L_y f - f\|_u &\rightarrow 0 \\ \|R_y f - f\|_u &\rightarrow 0 \end{aligned}$$

as $y \rightarrow 1_G$ respectively.

Proposition: If $f \in C_c(G)$, then f is left and right uniformly continuous.

Proof. We will prove this for $R_y f$.

If $f \in C_c(G)$, and $\varepsilon > 0$, then for every $x \in K = \text{supp}(f)$, there is a neighborhood U_x of 1_G such that

$$|f(xy) - f(x)| < \frac{1}{2}\varepsilon$$

for any $y \in U_x$. Similarly, there is a symmetric neighborhood V_x of 1_G such that $V_x V_x \subseteq U_x$; the sets xV_x cover K , so there exist $x_1, \dots, x_n \in K$ such that $K \subseteq \bigcup_{j=1}^n x_j V_{x_j}$.

Let $V = \bigcap_{j=1}^n V_{x_j}$. If $x \in K$, then there is some j such that $x_j^{-1}x \in V_{x_j}$, so $xy = x_j(x_j^{-1}x)y \in x_j U_{x_j}$, so

$$\begin{aligned} |f(xy) - f(x)| &\leq |f(xy) - f(x_j)| + |f(x_j) - f(x)| \\ &< \frac{1}{2}\varepsilon + \frac{1}{2}\varepsilon \\ &= \varepsilon, \end{aligned}$$

for any $y \in V$, meaning that $\|R_y f - f\|_u < \varepsilon$. Similarly, if $xy \in K$, then $|f(xy) - f(x)| < \varepsilon$; meanwhile, if $x, xy \notin K$, then $f(x) = f(xy) = 0$, so we are done. \square

Haar Measure

Definition: We define a subset of $C_c(G)$ to be

$$C_c^+(G) = \{f \in C_c(G) \mid f \geq 0, f \neq 0\}.$$

Definition: A left/right Haar measure on G is a nonzero Radon measure μ on G such that $\mu(xE) = \mu(E)$ for every Borel $E \subseteq G$ and all $x \in G$.