

## Chapter 2 Problems

### 2.3

#### Cylindrical Coordinates

Starting with our expression of  $\vec{r}$ , we have

$$\begin{aligned}\vec{r} &= \rho \cos \phi \hat{i} + \rho \sin \phi \hat{j} + z \hat{k} \\ d\vec{r} &= \frac{\partial \vec{r}}{\partial \rho} d\rho + \frac{\partial \vec{r}}{\partial \phi} d\phi + \frac{\partial \vec{r}}{\partial z} dz.\end{aligned}$$

Calculating each partial derivative,

$$\begin{aligned}\frac{\partial \vec{r}}{\partial \rho} &= \cos \phi \hat{i} + \sin \phi \hat{j} \\ \hat{\rho} &= \frac{\frac{\partial \vec{r}}{\partial \rho}}{\left\| \frac{\partial \vec{r}}{\partial \rho} \right\|} \\ &= \cos \phi \hat{i} + \sin \phi \hat{j}, \\ \frac{\partial \vec{r}}{\partial \phi} &= \rho (-\sin \phi \hat{i} + \cos \phi \hat{j}) \\ \hat{\phi} &= \frac{\frac{\partial \vec{r}}{\partial \phi}}{\left\| \frac{\partial \vec{r}}{\partial \phi} \right\|} \\ &= -\sin \phi \hat{i} + \cos \phi \hat{j}\end{aligned}$$

implying

$$\frac{\partial \vec{r}}{\partial \phi} = \rho \hat{\phi},$$

and finally, we have

$$\frac{\partial \vec{r}}{\partial z} = \hat{k}.$$

The above calculations yield

$$d\vec{r} = (d\rho) \hat{\rho} + (\rho d\phi) \hat{\phi} + \hat{k} dz.$$

#### Spherical Coordinates

Starting with our expression of  $\vec{x}$ <sup>I</sup>

$$\begin{aligned}\vec{x} &= r \sin \phi \sin \theta \hat{i} + r \cos \phi \sin \theta \hat{j} + r \cos \theta \hat{k} \\ d\vec{x} &= \frac{\partial \vec{x}}{\partial r} dr + \frac{\partial \vec{x}}{\partial \phi} d\phi + \frac{\partial \vec{x}}{\partial \theta} d\theta,\end{aligned}$$

---

<sup>I</sup>I am using  $\vec{x}$  instead of  $\vec{r}$  because  $r$  is already used in the expression of the spherical coordinates.

Evaluating each partial derivative, we have

$$\begin{aligned}\frac{\partial \vec{x}}{\partial r} &= \sin \phi \sin \theta \hat{i} + \cos \phi \sin \theta \hat{j} + \cos \theta \hat{k} \\ \hat{r} &= \frac{\frac{\partial \vec{x}}{\partial r}}{\left\| \frac{\partial \vec{x}}{\partial r} \right\|} \\ &= \sin \phi \sin \theta \hat{i} + \cos \phi \sin \theta \hat{j} + \cos \theta \hat{k}, \\ \frac{\partial \vec{x}}{\partial \phi} &= -r \sin \phi \sin \theta \hat{i} + r \cos \phi \sin \theta \hat{j} \\ \hat{\phi} &= \frac{\frac{\partial \vec{x}}{\partial \phi}}{\left\| \frac{\partial \vec{x}}{\partial \phi} \right\|} \\ &= -\sin \phi \sin \theta \hat{i} + \cos \phi \sin \theta \hat{j}\end{aligned}$$

implying

$$\frac{\partial \vec{x}}{\partial \phi} = r \sin \theta \hat{\phi},$$

and finally, we have

$$\begin{aligned}\frac{\partial \vec{x}}{\partial \theta} &= r \cos \phi \cos \theta \hat{i} + r \sin \phi \cos \theta \hat{j} - r \sin \theta \hat{k} \\ \hat{\theta} &= \frac{\frac{\partial \vec{x}}{\partial \theta}}{\left\| \frac{\partial \vec{x}}{\partial \theta} \right\|} \\ &= \cos \phi \cos \theta \hat{i} + \sin \phi \cos \theta \hat{j} - \sin \theta \hat{k},\end{aligned}$$

implying

$$\frac{\partial \vec{x}}{\partial \theta} = r \hat{\theta}.$$

The above calculations yield

$$d\vec{x} = (dr) \hat{r} + (r \sin \theta d\phi) \hat{\phi} + (r d\theta) \hat{\theta}.$$

## 2.8

Let

$$\vec{a} = r_a \cos \phi_a \sin \theta_a \hat{i} + r_a \sin \phi_a \sin \theta_a \hat{j} + r_a \cos \theta_a \hat{k} \quad \vec{b} = r_b \cos \phi_b \sin \theta_b \hat{i} + r_b \sin \phi_b \sin \theta_b \hat{j} + r_b \cos \theta_b \hat{k}.$$

Then,

$$\begin{aligned}\cos \gamma &= \frac{\vec{a} \cdot \vec{b}}{\left\| \vec{a} \right\| \left\| \vec{b} \right\|} \\ &= \frac{1}{r_a r_b} (r_a r_b (\sin \theta_a \sin \theta_b (\cos \phi_a \cos \phi_b + \sin \phi_a \sin \phi_b) + \cos \theta_a \cos \theta_b)) \\ &= \cos \theta_a \cos \theta_b + \sin \theta_a \sin \theta_b \cos (\phi_a - \phi_b).\end{aligned}$$

## 2.9

$$\begin{aligned}
\frac{d\vec{v}}{dt} &= \frac{d}{dt}(\dot{\rho}\hat{\rho}) + \frac{d}{dt}(\rho\dot{\phi}\hat{\phi}) \\
&= \dot{\rho}\ddot{\rho} + \dot{\rho}\frac{d\hat{\rho}}{dt} + \dot{\rho}\dot{\phi}\hat{\phi} + \rho\ddot{\phi}\hat{\phi} + \rho\dot{\phi}\frac{d\hat{\phi}}{dt} \\
&= \dot{\rho}\ddot{\rho} + \dot{\rho}\dot{\phi}\hat{\phi} + \rho\ddot{\phi}\hat{\phi} + \dot{\rho}\dot{\phi}\hat{\phi} + \left(\frac{\partial\hat{\phi}}{\partial\rho}\frac{d\rho}{dt} + \frac{\partial\hat{\phi}}{\partial\phi}\frac{d\phi}{dt}\right) \\
&= (\ddot{\rho} - \rho\dot{\phi}^2)\hat{\rho} + (\rho\ddot{\phi} + 2\dot{\rho}\dot{\phi})\hat{\phi}.
\end{aligned}$$

## Chapter 3 Problems

For all problems involving  $\arg z$  (or equivalents), I will be using the principle branch,  $\arg z \in (-\pi, \pi]$ .

## 3.5

(a)

$$\begin{aligned}
\sqrt{3} + i &= 2e^{i\frac{\pi}{3}} \\
-\sqrt{3} + i &= 2e^{i\frac{2\pi}{3}}
\end{aligned}$$

(b)

$$\begin{aligned}
\sqrt{2}i &= \sqrt{2}e^{i\frac{\pi}{4} + n\pi} \\
\sqrt{2 + 2\sqrt{3}i} &= 2e^{i\frac{\pi}{6}}
\end{aligned}$$

## 3.6

(a) Real:

$$\begin{aligned}
(-1)^{1/i} &= (e^{i\pi})^{-i} \\
&= e^{\pi}.
\end{aligned}$$

(b) Real:

$$\begin{aligned}
\left(\frac{z}{z^*}\right)^i &= (e^{2i\arg z})^i \\
&= e^{-2\arg z}.
\end{aligned}$$

(c) Imaginary:

$$\begin{aligned}
(z_1 z_2^* - z_1^* z_2)^* &= z_1^* z_2 - z_1 z_2^* \\
&= -(z_1 z_2^* - z_1^* z_2).
\end{aligned}$$

(d) Complex:

$$\sum_{n=0}^N e^{in\theta} = \frac{1 - e^{iN\theta}}{1 - e^{i\theta}}.$$

(e) Real: for each  $a \in \{1, 2, \dots, N\}$ ,  $e^{ia\theta} + e^{-ia\theta} \in \mathbb{R}$ .

**3.9**

(a)

$$\begin{aligned}
 \cos(a+b) + \cos(a-b) &= \frac{1}{2} \left( e^{i(a+b)} + e^{-i(a+b)} \right) + \frac{1}{2} \left( e^{i(a-b)} + e^{-i(a-b)} \right) \\
 &= \frac{1}{2} \left( e^{ia} (e^{ib} + e^{-ib}) + e^{-ia} (e^{ib} + e^{-ib}) \right) \\
 &= \frac{1}{2} (e^{ia} + e^{-ia}) (e^{ib} + e^{-ib}) \\
 &= 2 \cos a \cos b.
 \end{aligned}$$

(b)

$$\begin{aligned}
 \sin(a+b) + \sin(a-b) &= \frac{1}{2i} \left( e^{i(a+b)} - e^{-i(a+b)} \right) + \frac{1}{2i} \left( e^{i(a-b)} - e^{-i(a-b)} \right) \\
 &= \frac{1}{2i} \left( e^{ia} (e^{ib} + e^{-ib}) - e^{-ia} (e^{ib} + e^{-ib}) \right) \\
 &= \frac{1}{2i} (e^{ia} - e^{-ia}) (e^{ib} + e^{-ib}) \\
 &= 2 \sin a \cos b.
 \end{aligned}$$

**3.10**

(a)

$$\begin{aligned}
 e^{i\alpha} + e^{i\beta} &= e^{i\left(\frac{\alpha+\beta}{2} + \frac{\alpha-\beta}{2}\right)} + e^{i\left(\frac{\alpha+\beta}{2} - \frac{\alpha-\beta}{2}\right)} \\
 &= e^{i\frac{\alpha+\beta}{2}} \left( e^{i\frac{\alpha-\beta}{2}} + e^{-i\frac{\alpha-\beta}{2}} \right) \\
 &= 2 \cos\left(\frac{\alpha-\beta}{2}\right) e^{i\frac{\alpha+\beta}{2}}.
 \end{aligned}$$

(b)

$$\begin{aligned}
 e^{i\alpha} - e^{i\beta} &= e^{i\left(\frac{\alpha+\beta}{2} + \frac{\alpha-\beta}{2}\right)} - e^{i\left(\frac{\alpha+\beta}{2} - \frac{\alpha-\beta}{2}\right)} \\
 &= e^{i\frac{\alpha+\beta}{2}} \left( e^{i\frac{\alpha-\beta}{2}} - e^{-i\frac{\alpha-\beta}{2}} \right) \\
 &= 2i \sin\left(\frac{\alpha-\beta}{2}\right) e^{i\frac{\alpha+\beta}{2}}
 \end{aligned}$$

**3.12**

$$\begin{aligned}
 \frac{1}{2i} \ln\left(\frac{a+ib}{a-ib}\right) &= \frac{1}{2i} (\ln(a+ib) - \ln(a-ib)) \\
 &= \frac{1}{2i} \left( \ln|a+ib| + i \arctan\left(\frac{b}{a}\right) - \left( \ln|a+ib| + i \arctan\left(-\frac{b}{a}\right) \right) \right) \\
 &= \arctan\left(\frac{b}{a}\right).
 \end{aligned}$$

## 3.13

$$\begin{aligned}
\frac{d^n}{dt^n} (e^{at} \sin bt) &= \frac{1}{2i} \frac{d^n}{dt^n} (e^{(a+ib)t} - e^{(a-ib)t}) \\
&= \frac{1}{2i} \left( (a+ib)^n e^{(a+ib)t} - (a-ib)^n e^{(a-ib)t} \right) \\
&= e^{at} \frac{1}{2i} (a^2 + b^2)^{n/2} \left( e^{i(b+n \arctan(\frac{b}{a}))t} - e^{i(b+n \arctan(\frac{b}{a}))t} \right) \\
&= e^{at} (a^2 + b^2)^{n/2} \sin \left( bt + n \arctan \left( \frac{b}{a} \right) \right)
\end{aligned}$$

## 3.20

Showing the equivalence between  $C_1 \cos kx + C_2 \sin kx$  and  $A \cos(kx + \alpha)$  and  $B \sin(kx + \beta)$ , we have

$$\begin{aligned}
A \cos(kx + \alpha) &= A \cos kx \cos \alpha - A \sin kx \sin \alpha \\
B \sin(kx + \beta) &= B \cos kx \sin \beta + B \sin kx \cos \beta
\end{aligned}$$

meaning (assuming  $\alpha, \beta \neq \pi n, \pi/2 + \pi n$ )

$$\begin{aligned}
A &= \frac{C_1}{\cos \alpha} \\
&= -\frac{C_2}{\sin \alpha} \\
B &= \frac{C_1}{\sin \beta} \\
&= \frac{C_2}{\cos \beta}.
\end{aligned}$$

Now, we show the equivalence between  $C_1 \cos kx + C_2 \sin kx$  and  $D_1 e^{ikx} + D_2 e^{-ikx}$ .

$$\begin{aligned}
D_1 e^{ikx} + D_2 e^{-ikx} &= \frac{D_1 + D_2}{2} (e^{ikx} + e^{-ikx}) + \frac{D_1 - D_2}{2} (e^{ikx} - e^{-ikx}) \\
&= (D_1 + D_2) \cos kx + i(D_1 - D_2) \sin kx.
\end{aligned}$$

meaning

$$\begin{aligned}
C_1 &= D_1 + D_2 \\
C_2 &= i(D_1 - D_2).
\end{aligned}$$

Finally, we show the equivalence between  $\text{Re}(Fe^{ikx})$  and  $C_1 \cos kx + C_2 \sin kx$ .

$$\begin{aligned}
\text{Re}((a+ib)e^{ikx}) &= \text{Re}(a \cos kx + ia \sin kx + ib \cos kx - b \sin kx) \\
&= a \cos kx - b \sin kx,
\end{aligned}$$

meaning

$$\begin{aligned}
C_1 &= a \\
C_2 &= -b.
\end{aligned}$$