

Amenability: A (Somewhat) Brief Introduction

Avinash Iyer

Occidental College

March 20, 2025

Outline

- 1 Definitions
- 2 Paradoxical Decompositions
- 3 From Paradoxical Decompositions to Amenability
- 4 Hereditary Properties and Equivalent Definitions
- 5 Remarks and Acknowledgments

Contents

- 1 Definitions
- 2 Paradoxical Decompositions
- 3 From Paradoxical Decompositions to Amenability
- 4 Hereditary Properties and Equivalent Definitions
- 5 Remarks and Acknowledgments

Groups

If A is a set, and $\star: A \times A \rightarrow A$ is an operation such that

- $a \star (b \star c) = (a \star b) \star c$;
- there exists e_A such that $a \star e_A = e_A \star a = a$;
- for each a there exists a^{-1} such that $a \star a^{-1} = a^{-1} \star a = e_A$,

then we call the pair (A, \star) a *group*.

Groups

If A is a set, and $\star: A \times A \rightarrow A$ is an operation such that

- $a \star (b \star c) = (a \star b) \star c$;
- there exists e_A such that $a \star e_A = e_A \star a = a$;
- for each a there exists a^{-1} such that $a \star a^{-1} = a^{-1} \star a = e_A$,

then we call the pair (A, \star) a *group*.

We abbreviate $a \star b$ as ab .

Subgroups, Quotient Groups

Let G be a group.

- If $H \subseteq G$ is a subset that satisfies, for all $a, b \in H$, $ab^{-1} \in H$, then we say H is a *subgroup*.

Subgroups, Quotient Groups

Let G be a group.

- If $H \subseteq G$ is a subset that satisfies, for all $a, b \in H$, $ab^{-1} \in H$, then we say H is a *subgroup*.
- If $N \subseteq G$ is a subgroup that satisfies, for all $g \in G$ and $h \in N$, $ghg^{-1} \in N$, then we say N is a *normal subgroup*.

Subgroups, Quotient Groups

Let G be a group.

- If $H \subseteq G$ is a subset that satisfies, for all $a, b \in H$, $ab^{-1} \in H$, then we say H is a *subgroup*.
- If $N \subseteq G$ is a subgroup that satisfies, for all $g \in G$ and $h \in N$, $ghg^{-1} \in N$, then we say N is a *normal subgroup*.
- The equivalence classes under the relation $g \sim_N g'$ if $g^{-1}g' \in N$ form a group $gN := [g]_{\sim}$ known as the *quotient group* G/N .

Some Groups

- The integers \mathbb{Z} are a group under addition.
- The group of invertible $n \times n$ matrices over \mathbb{C} , $GL_n(\mathbb{C})$, is a group under matrix multiplication.
- The subgroup $SO(n) \subseteq GL_n(\mathbb{R})$ consisting of $n \times n$ orthogonal matrices with determinant 1 is a group under multiplication.

Group Actions

Let G be a group, and X a set. Let $\rho: G \times X \rightarrow X$ be a function that satisfies, for all $g, h \in G$ and $x \in X$,

- $\rho(e_G, x) = x$;
- $\rho(g, \rho(h, x)) = \rho(gh, x)$.

Then, we say ρ is an *action* of G on X . We write $\rho(g, x) = g \cdot x$.

σ -Algebras and Measures

If X is a set, then a collection of subsets $\{A_i\}_{i \in I} = \mathcal{A} \subseteq P(X)$ is known as an *algebra* of subsets if

- ① $\emptyset, X \in \mathcal{A}$;
- ② for any $A_i \in \mathcal{A}$, $A_i^c \in \mathcal{A}$;
- ③ for any $A_i, A_j \in \mathcal{A}$, $A_i \cup A_j \in \mathcal{A}$.

σ -Algebras and Measures

If X is a set, then a collection of subsets $\{A_i\}_{i \in I} = \mathcal{A} \subseteq P(X)$ is known as an *algebra* of subsets if

- ① $\emptyset, X \in \mathcal{A}$;
- ② for any $A_i \in \mathcal{A}$, $A_i^c \in \mathcal{A}$;
- ③ for any $A_i, A_j \in \mathcal{A}$, $A_i \cup A_j \in \mathcal{A}$.

If, for any countable collection, $\{A_n\}_{n \geq 1} \subseteq \mathcal{A}$, condition (3) holds, then we say \mathcal{A} is a σ -*algebra* of subsets.

σ -Algebras and Measures, Cont'd

If X is a set and \mathcal{A} is a σ -algebra, then a map $\mu: \mathcal{A} \rightarrow [0, \infty]$ that satisfies:

- $\mu(\emptyset) = 0$;
- for disjoint sets $A, B \in \mathcal{A}$, $\mu(A \sqcup B) = \mu(A) + \mu(B)$,

then we say μ is a *finitely additive* measure.

σ -Algebras and Measures, Cont'd

If X is a set and \mathcal{A} is a σ -algebra, then a map $\mu: \mathcal{A} \rightarrow [0, \infty]$ that satisfies:

- $\mu(\emptyset) = 0$;
- for disjoint sets $A, B \in \mathcal{A}$, $\mu(A \sqcup B) = \mu(A) + \mu(B)$,

then we say μ is a *finitely additive* measure. If $\{A_n\}_{n \geq 1}$ is a countable collection of disjoint sets, then if μ satisfies

- $\mu\left(\bigcup_{n \geq 1} A_n\right) = \sum_{n \geq 1} \mu(A_n)$,

we say μ is a measure.

Contents

- 1 Definitions
- 2 Paradoxical Decompositions**
- 3 From Paradoxical Decompositions to Amenability
- 4 Hereditary Properties and Equivalent Definitions
- 5 Remarks and Acknowledgments

Questions?

- If G is a group, is it possible to reconstruct G by using some subset of G ?
- When may we find a finitely additive probability measure $\mu: P(G) \rightarrow [0, 1]$ such that $\mu(E) = \mu(tE)$ for all $E \subseteq G$?
- Are these questions even related?

Free Groups

- We begin by considering a special group, known as $F(a, b)$ or the *free group on two generators*.

Free Groups

- We begin by considering a special group, known as $F(a, b)$ or the *free group on two generators*.
- We define $F(a, b)$ to be the set of all “words” in the alphabet $\{a, b, a^{-1}, b^{-1}\}$, subject to the condition that, for $w, w' \in F(a, b)$,

$$\begin{aligned} waa^{-1}w' &\sim wa^{-1}aw' \sim ww' \\ wbb^{-1}w' &\sim wb^{-1}bw' \sim ww'. \end{aligned}$$

- Examples: $a^2bab^{-1}, b^{-1}a^2b^2ab \in F(a, b)$.

A Curiosity

Let $W(b) \subseteq F(a, b)$ be all the words that start with b . Then, $b^{-1}W(b)$ consists of

A Curiosity

Let $W(b) \subseteq F(a, b)$ be all the words that start with b . Then, $b^{-1}W(b)$ consists of

- all words that start with a ;
- all words that start with a^{-1} ;
- all words that start with b — think words that start with b^2 before you multiply b^{-1} .

A Curiosity

Let $W(b) \subseteq F(a, b)$ be all the words that start with b . Then, $b^{-1}W(b)$ consists of

- all words that start with a ;
- all words that start with a^{-1} ;
- all words that start with b — think words that start with b^2 before you multiply b^{-1} .

Thus, all we need to do is add back $W(b^{-1})$ to get $F(a, b)$ back.

$$F(a, b) = W(b^{-1}) \cup b^{-1}W(b).$$

A Curiosity

Let $W(b) \subseteq F(a, b)$ be all the words that start with b . Then, $b^{-1}W(b)$ consists of

- all words that start with a ;
- all words that start with a^{-1} ;
- all words that start with b — think words that start with b^2 before you multiply b^{-1} .

Thus, all we need to do is add back $W(b^{-1})$ to get $F(a, b)$ back.

$$F(a, b) = W(b^{-1}) \cup b^{-1}W(b).$$

A Curiosity, Cont'd

Similarly, we can do this for a , giving a decomposition of $F(a, b)$ in two separate ways:

$$\begin{aligned} F(a, b) &= b^{-1} W(b) \cup W(b^{-1}) \\ &= a^{-1} W(a) \cup W(a^{-1}). \end{aligned}$$

A Curiosity, Cont'd

Similarly, we can do this for a , giving a decomposition of $F(a, b)$ in two separate ways:

$$\begin{aligned} F(a, b) &= b^{-1}W(b) \cup W(b^{-1}) \\ &= a^{-1}W(a) \cup W(a^{-1}). \end{aligned}$$

Furthermore, note that $W(a), W(b), W(a^{-1}), W(b^{-1})$ are disjoint.

These decompositions seem to be downright paradoxical — we take a part of the group, translate some of it, and get the whole group back!

Defining Paradoxical Decompositions

Let G be a group. A *paradoxical decomposition* of G consists of

- pairwise disjoint subsets $A_1, \dots, A_n, B_1, \dots, B_m \subseteq G$; and
- elements $g_1, \dots, g_n, h_1, \dots, h_m$;

such that

$$\begin{aligned} G &= \bigcup_{i=1}^n g_i A_i \\ &= \bigcup_{j=1}^m h_j B_j. \end{aligned}$$

Defining Paradoxical Decompositions

Let G be a group. A *paradoxical decomposition* of G consists of

- pairwise disjoint subsets $A_1, \dots, A_n, B_1, \dots, B_m \subseteq G$; and
- elements $g_1, \dots, g_n, h_1, \dots, h_m$;

such that

$$\begin{aligned} G &= \bigcup_{i=1}^n g_i A_i \\ &= \bigcup_{j=1}^m h_j B_j. \end{aligned}$$

If G admits a paradoxical decomposition, we say G is *paradoxical*.

Examples

- The free group $F(a, b)$ is paradoxical.

Examples

- The free group $F(a, b)$ is paradoxical.
- Any group that contains a paradoxical subgroup is paradoxical.
- $F(S)$, where S is any nonempty set with more than two elements, is paradoxical.

A Paradoxical Subgroup of $SO(3)$

The following two matrices (and their inverses) generate a subgroup of $SO(3)$ that is isomorphic to $F(a, b)$.

$$A = \begin{pmatrix} 3/5 & 4/5 & 0 \\ -4/5 & 3/5 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3/5 & -4/5 \\ 0 & 4/5 & 3/5 \end{pmatrix}.$$

A Paradoxical Subgroup of $SO(3)$

The following two matrices (and their inverses) generate a subgroup of $SO(3)$ that is isomorphic to $F(a, b)$.

$$A = \begin{pmatrix} 3/5 & 4/5 & 0 \\ -4/5 & 3/5 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$
$$B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3/5 & -4/5 \\ 0 & 4/5 & 3/5 \end{pmatrix}.$$

This is proven using the Ping-Pong lemma.

Contents

- 1 Definitions
- 2 Paradoxical Decompositions
- 3 From Paradoxical Decompositions to Amenability**
- 4 Hereditary Properties and Equivalent Definitions
- 5 Remarks and Acknowledgments

Contents

- 1 Definitions
- 2 Paradoxical Decompositions
- 3 From Paradoxical Decompositions to Amenability
- 4 Hereditary Properties and Equivalent Definitions**
- 5 Remarks and Acknowledgments

Contents

- 1 Definitions
- 2 Paradoxical Decompositions
- 3 From Paradoxical Decompositions to Amenability
- 4 Hereditary Properties and Equivalent Definitions
- 5 Remarks and Acknowledgments**