

1.8

Problem. Fix a natural number $b \geq 2$. Show that every positive real number x in $[0, 1]$ has a b -adic expansion of the form

$$x = \sum_{n=1}^{\infty} \frac{x_n}{b^n},$$

with each $0 \leq x_n \leq b - 1$.

1.9

Problem. Suppose

$$\sum_{n=1}^{\infty} \frac{x_n}{b^n} = \sum_{n=1}^{\infty} \frac{y_n}{b^n},$$

with $0 \leq x_n \leq b - 1$ and $0 \leq y_n \leq b - 1$ integers. Show that either $x_n = y_n$ for all n , or there is an m such that one of the following two cases occurs:

- $x_m = y_m + 1$ and for $n \geq m + 1$, $y_n = b - 1$ and $x_n = 0$;
- $y_m = x_m + 1$ and for $n \geq m + 1$, $x_n = b - 1$ and $y_n = 0$.

1.10

Problem. Show that a number $x \in [0, 1]$ is rational if and only if its decimal expansion is eventually periodic. Deduce that irrational numbers have unique decimal expansions.

Solution. Let x be rational. Then, $x = \frac{p}{q}$, with $p \in \mathbb{Z}_{\geq 0}$, $q \in \mathbb{Z}_{>0}$, with $\frac{p}{q}$ in lowest terms, with $q > p$.

We write $10x = x_1 + y_1$, with $x_1 = \lfloor 10x \rfloor$ and $y_1 = 10x - \lfloor 10x \rfloor$. Thus, we have

$$\begin{aligned} y_1 &= \frac{10p}{q} - \frac{qx_1}{q} \\ &= \frac{10p - qx_1}{q} \\ &= \frac{m_1}{q}. \end{aligned}$$

We want to show that $0 \leq m_1 < q$.

Now, we take $10y_1 = x_2 + y_2$, with

$$\begin{aligned} y_2 &= \frac{10m_1}{q} - \frac{qx_2}{q} \\ &= \frac{m_2}{q}. \end{aligned}$$

Repeatedly, we get $y_n = \frac{m_n}{q}$.

We have $0 \leq x_i < 10$, and $0 \leq m_i < q$. Thus, looking at the set of pairs $(x_1, m_1), (x_2, m_2), \dots$. Since x_i and m_i are limited, there cannot be infinitely many distinct pairs; thus, there will necessarily be a value of n such that $(x_k, m_k) = (x_{k+n}, m_{k+n})$.

1.11

Problem. Show that the collection of polynomials with rational coefficients is a countably infinite set.

Solution. Let $\mathcal{P}_n(\mathbb{Q})$ denote the set of polynomials with degree n with coefficients in \mathbb{Q} . We construct a bijection

$$\mathcal{P}_n(\mathbb{Q}) \rightarrow \prod_{k=0}^n \mathbb{Q},$$

where \prod denotes the Cartesian product, by taking

$$a_0 + a_1x + \cdots + a_nx^n \mapsto (a_0, a_1, \dots, a_n).$$

Since $\prod_{k=0}^n \mathbb{Q}$ is a countable Cartesian product of countable sets, this means $\mathcal{P}_n(\mathbb{Q})$ is countable.

Finally, we have $\mathbb{Q}[x]$, the set of all polynomials with rational coefficients, is

$$\mathbb{Q}[x] = \bigcup_{k=0}^{\infty} \mathcal{P}_k(\mathbb{Q}),$$

meaning $\mathbb{Q}[x]$ is countable.

1.12

1.13

Extra Problem 1

Extra Problem 2

Problem. If $|A| \leq |B|$, then $|P(A)| \leq |P(B)|$.

Solution. Let $f : A \hookrightarrow B$ be an injection. Given $S \subseteq A$, we have $f(S) \subseteq B$, meaning $S \in P(A)$ implies $f(S) \in P(B)$. We let $g : P(A) \rightarrow P(B)$ be induced by f , with

$$\begin{aligned} g(S) &= f(S) \\ &= \{f(x) \mid x \in S\}. \end{aligned}$$

Extra Problem 3

Extra Problem 4